




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 215/54, A61K 31/47, A61P 43/00, C07D 405/12, 401/12, 417/12, 413/12, 409/12		A1	(11) International Publication Number: WO 00/68201
			(43) International Publication Date: 16 November 2000 (16.11.00)
(21) International Application Number: PCT/GB00/01697 (22) International Filing Date: 3 May 2000 (03.05.00) (30) Priority Data: 9910577.7 8 May 1999 (08.05.99) GB (71) Applicant (for all designated States except US): AS-TRAZENECA AB [SE/SE]; S-151 85 Södertälje (SE). (72) Inventors; and (75) Inventors/Applicants (for US only): BOYLE, Francis, Thomas [GB/GB]; Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). GIBSON, Keith, Hopkinson [GB/GB]; Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). POYSER, Jeffrey, Philip [GB/GB]; Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). TURNER, Paul [GB/GB]; Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). (74) Agent: GILES, Allen, Frank; AstraZeneca, Global Intellectual Property – Patents, Mereside, Alderley Park, P.O. Box 272, Macclesfield, Cheshire SK10 4TG (GB).		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	

(54) Title: QUINOLINE DERIVATIVES AS INHIBITORS OF MEK ENZYMES

(57) Abstract

A compound of formula (I) or a pharmaceutically acceptable salt thereof wherein: n is 0–1; X and Y are independently selected from NH–, –O–, –S–, or NR⁸– where R⁸ is alkyl of 1–6 carbon atoms and X may additionally comprise a CH₂ group; R⁷ is a group (CH₂)_mR⁹ where m is 0, or an integer of from 1–3 and R⁹ is a substituted aryl group, an optionally substituted cycloalkyl ring of up to 10 carbon atoms, or an optionally substituted heterocyclic ring or an N–oxide of any nitrogen containing ring; R⁶ is a divalent cycloalkyl of 3 to 7 carbon atoms, which may be optionally further substituted with one or more alkyl of 1 to 6 carbon atom groups; or is a divalent pyridinyl, pyrimidinyl, or phenyl ring; wherein the pyridinyl, pyrimidinyl, or phenyl ring may be optionally further substituted with one or more specified groups; R₁, R₂, R₃ and R₄ are each independently selected from hydrogen or various specified organic groups. Compounds are useful as pharmaceuticals for the inhibition of MEK activity.



Compounds are useful as pharmaceuticals for the inhibition of MEK activity.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

QUINOLINE DERIVATIVES AS INHIBITORS OF MEK ENZYMES

The present invention relates to certain novel quinoline derivatives as well as to their use as pharmaceuticals, in particular as inhibitors of specific kinase enzymes, such as MEK enzymes. Further aspects of the invention include pharmaceutical compositions and methods of treatment of proliferative disease such as cancer using said compounds.

Cancer is a disease in which cells grow and divide in an uncontrolled fashion. This uncontrolled growth arises from abnormalities in signal transduction pathways that are used by normal cells to regulate cell growth and division in response to various signalling molecules. Normal cells do not proliferate unless stimulated to do so by specific signal molecules located outside the cell derived from nearby cells or tissues. Growth factors bind to the cell membrane via specific receptors which have intrinsic enzyme activity. These receptors relay the growth signal to the cell nucleus via a series of signalling proteins. In cancer, a number of defects in signal pathways are apparent. For example, cancer cells may produce their own growth factors which bind to their cognate receptors, resulting in an autocrine loop, or receptors may be mutated or overexpressed leading to an increased, continuous signal to proliferate. In addition, negative regulators of cell growth may be lost.

Oncogenes are cancer related genes which often encode abnormal versions of signal pathway components, such as receptor tyrosine kinases, serine-threonine kinases, or downstream signaling molecules such as the ras genes, which code for closely related small guanine nucleotide binding proteins which hydrolyse bound guanosine triphosphate (GTP) to guanosine diphosphate (GDP). Ras proteins are active in promoting cell growth and transformation when they are bound to GTP and inactive when they are bound to GDP. Transforming mutants of p21ras are defective in their GTPase activity and hence remain in the active GTP bound state. The ras oncogene is known to play an integral role in certain cancers, and has been found to contribute to the formation of over 20% of all cases of human cancer.

When activated by ligand, cell surface receptors which are coupled to the mitogenic response, such as growth factor receptors, initiate a chain of reactions which leads to the activation of guanine nucleotide exchange activity on ras. When in its active GTP-bound state, a number of proteins interact directly with ras at the plasma membrane

resulting in signal transmission through several distinct pathways. The best characterised effector protein is the product of the raf proto-oncogene. The interaction of raf and ras is a key regulatory step in the control of cell proliferation. Ras-mediated activation of the raf serine-threonine kinase in turn activates the dual-specificity MEK (MEK1 and MEK2), which is the immediate upstream activator of mitogen activated protein kinase (MAPKs known as extracellular signal regulated protein kinases or ERK1 and ERK2). To date, no substrates of MEK other than MAPK have been identified, though recent reports indicate that MEK may also be activated by other upstream signal proteins such as MEK kinase or MEKK1 and PKC. Activated MAPK translocates and accumulates in the nucleus, where it can phosphorylate and activate transcription factors such as Elk-1 and Sap1a, leading to the enhanced expression of genes such as that for c-fos.

The ras-dependent raf-MEK-MAPK cascade is one of the key signalling pathways responsible for transmitting and amplifying mitogenic signals from cell surface to the nucleus resulting in changes in gene expression and cell fate. This ubiquitous pathway appears essential for normal cell proliferation and constitutive activation of this pathway is sufficient to induce cellular transformation. Transforming mutants of p21ras are constitutively active, resulting in raf, MEK and MAPK activity and cell transformation. Inhibition of MEK activity using either antisense raf, a dominant negative MEK mutant or the selective inhibitor PD098059 have been shown to block the growth and morphological transformation of ras-transformed fibroblasts.


The mechanism of activation of raf, MEK and MAPK is through phosphorylation on specific serine, threonine or tyrosine residues. Activated raf and other kinases phosphorylate MEK1 on S218 and S222 and MEK2 on S222 and S226. This results in MEK activation and subsequent phosphorylation and activation of ERK1 on T190 and Y192 and ERK2 on T183 and Y185 by the dual specificity MEKs. Whilst MEK can be activated by a number of protein kinases, and active MAPKs phosphorylate and activate a number of substrate proteins including transcription factors and other protein kinases, MEKs appear specific and sole activators of MAPKs and could act as a focal point for cross-cascade regulation. MEK1 and MEK2 isoforms show unusual specificity and also contain a proline-rich insert between catalytic subdomains IX and X which is not present in any of the other known MEK family members. These differences between MEK and other protein kinases, together with the known role of MEK in proliferative signalling

suggest that it may be possible to discover and employ selective MEK inhibitors as therapeutic agents for use in proliferative disease.

WO 98/43960 discloses a range of 3-cyano quinoline compounds and their use in the treatment of cancer. Certain of the compounds are demonstrated as being inhibitors of Epidermal Growth Factor Receptor Kinase, and to inhibit cancer cell growth. Other quinoline derivatives which inhibit the effect of growth factors such as VEGF are described in WO98/13350.

This invention provides compounds which are inhibitors of the kinase activity of MEK and as a result, can produce therapeutically useful effects in the treatment of proliferative disease and in particular cancer.

According to the present invention there is provided a compound of formula (I)



(I)

or a pharmaceutically acceptable salt thereof.

wherein:

n is 0-1;

X and Y are independently selected from -NH-, -O-, -S-, or -NR⁸- where R⁸ is alkyl of 1-6 carbon atoms and X may additionally comprise a CH₂ group;

R⁷ is a group (CH₂)_mR⁹ where m is 0, or an integer of from 1-3 and R⁹ is a substituted aryl group, an optionally substituted cycloalkyl ring of up to 10 carbon atoms, or an optionally substituted heterocyclic ring or an N-oxide of any nitrogen containing ring;

R⁶ is a divalent cycloalkyl of 3 to 7 carbon atoms, which may be optionally further substituted with one or more alkyl of 1 to 6 carbon atom groups; or is a divalent pyridinyl, pyrimidinyl, or phenyl ring; wherein the pyridinyl, pyrimidinyl, or phenyl ring may be optionally further substituted with one or more groups selected from halogen,

- alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, azido, hydroxyalkyl of 1-6 carbon atoms, halomethyl, alkoxymethyl of 2-7 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkoxy of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, hydroxy, trifluoromethyl, cyano, nitro, carboxy, carboalkoxy of 2-7 carbon atoms, carboalkyl of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzoyl, benzyl, amino, alkylamino of 1-6 carbon atoms, dialkylamino of 2 to 12 carbon atoms, phenylamino, benzylamino, alkanoylamino of 1-6 carbon atoms, alkenoylamino of 3-8 carbon atoms, alkynoylamino of 3-8 carbon atoms, and benzoylamino;
- 10 R_1 , R_2 , R_3 and R_4 are each independently selected from hydrogen, hydroxy, halogeno, cyano, nitro, trifluoromethyl, C_{1-3} alkyl, $-NR^{11}R^{12}$ (wherein R^{11} and R^{12} , which may be the same or different, each represents hydrogen or C_{1-3} alkyl), or a group $R^{13}-X^1-(CH_2)_x$ wherein x is 0 to 3, X^1 represents $-O-$, $-CH_2-$, $-OCO-$, carbonyl, $-S-$, $-SO-$, $-SO_2-$, $-NR^{14}CO-$, $-CONR^{15}-$, $-SO_2NR^{16}-$, $-NR^{17}SO_2-$ or $-NR^{18}-$ (wherein R^{14} , R^{15} , R^{16} , R^{17} and R^{18} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{13} is selected from one of the following sixteen groups:
- 1) C_{1-5} alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro and amino;
- 2) C_{1-5} alkyl IX^2COR^{19} (wherein X^2 represents $-O-$ or $-NR^{20}-$ (wherein R^{20} represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{19} represents $-NR^{21}R^{22}-$ or $-OR^{23}-$ (wherein R^{21} , R^{22} and R^{23} which may be the same or different each represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl));
- 3) C_{1-5} alkyl IX^3R^{24} (wherein X^3 represents $-O-$, $-S-$, $-SO-$, $-SO_2-$, $-OCO-$, $-NR^{25}CO-$, $-CONR^{26}-$, $-SO_2NR^{27}-$, $-NR^{28}SO_2-$ or $-NR^{29}-$ (wherein R^{25} , R^{26} , R^{27} , R^{28} and R^{29} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{24} represents hydrogen, C_{1-3} alkyl, cyclopentyl, cyclohexyl or a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which C_{1-3} alkyl group may bear one or two substituents selected from oxo, hydroxy, halogeno and C_{1-4} alkoxy and which cyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, C_{1-4} alkyl, C_{1-4} hydroxyalkyl and C_{1-4} alkoxy);
- 4) C_{1-5} alkyl IX^4C_{1-5} alkyl IX^5R^{30} (wherein X^4 and X^5 which may be the same or different are each $-O-$, $-S-$, $-SO-$, $-SO_2-$, $-NR^{31}CO-$, $-CONR^{32}-$, $-SO_2NR^{33}-$, $-NR^{34}SO_2-$ or $-NR^{35}-$

(wherein R^{31} , R^{32} , R^{33} , R^{34} and R^{35} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{30} represents hydrogen or C_{1-3} alkyl);

5 5) C_{1-5} alkyl R^{36} (wherein R^{36} is a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which heterocyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, C_{1-4} alkyl, C_{1-4} hydroxyalkyl and C_{1-4} alkoxy);

6) $(CH_2)_qX^6R^{37}$ (wherein q is an integer from 0 to 5, X^6 represents a direct bond, -O-, -S-, -SO-, -SO₂-, -NR³⁸CO-, -CONR³⁹-, -SO₂NR⁴⁰-, -NR⁴¹SO₂- or -NR⁴²- (wherein R^{38} , R^{39} , R^{40} , R^{41} and R^{42} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl)

10 and R^{37} is a phenyl group, a pyridone group or a 5 or 6 membered aromatic heterocyclic group with 1 to 3 heteroatoms selected from O, N and S, which phenyl, pyridone or aromatic heterocyclic group may carry up to 5 substituents selected from hydroxy, halogeno, amino, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} hydroxyalkyl, C_{1-4} hydroxyalkoxy, C_{1-4} aminoalkyl, C_{1-4} alkylamino, carboxy, cyano, -CONR⁴³R⁴⁴ and -NR⁴⁵COR⁴⁶ (wherein R^{43} , R^{44} , R^{45} and R^{46} , which may be the same or different, each represents hydrogen, C_{1-4} alkyl or C_{1-3} alkoxy C_{2-3} alkyl));

7) C_{2-6} alkenyl R^{36} (wherein R^{36} is as defined hereinbefore);

8) C_{2-6} alkynyl R^{36} (wherein R^{36} is as defined hereinbefore);

9) X^7R^{47} (wherein X^7 is -SO₂-, -O- or -CONR⁴⁸R⁴⁹- (wherein R^{48} and R^{49} , which may be the same or different, each represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{47} represents C_{1-5} alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro and amino) with the provisos that when X^7 is -SO₂-, X^1 is -O-, when X^7 is -O-, X^1 is carbonyl, when X^7 is -CONR⁴⁸R⁴⁹-, X^1 is -O- or NR¹⁸ (wherein R^{48} , R^{49} and R^{18} are as defined hereinbefore);

10) C_{2-6} alkenyl R^{37} (wherein R^{37} is as defined hereinbefore);

11) C_{2-6} alkynyl R^{37} (wherein R^{37} is as defined hereinbefore);

12) C_{2-6} alkenyl X^8R^{37} (wherein X^8 represents -O-, -S-, -SO-, -SO₂-, -NR⁵⁰CO-, -CONR⁵¹-, -SO₂NR⁵²-, -NR⁵³SO₂- or -NR⁵⁴- (wherein R^{50} , R^{51} , R^{52} , R^{53} and R^{54} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{37} is as defined hereinbefore);

13) C_{2-6} alkynyl X^9R^{37} (wherein X^9 represents -O-, -S-, -SO-, -SO₂-, -NR⁵⁵CO-, -CONR⁵⁶-, -SO₂NR⁵⁷-, -NR⁵⁸SO₂- or -NR⁵⁹- (wherein R^{55} , R^{56} , R^{57} , R^{58} and R^{59} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{37} is as defined hereinbefore);

14) $C_{1-3}\text{alkyl}X^{10}C_{1-3}\text{alkyl}R^{37}$ (wherein X^{10} represents -O-, -S-, -SO-, -SO₂-, -NR⁶⁰CO-, -CONR⁶¹-, -SO₂NR⁶²-, -NR⁶³SO₂- or -NR⁶⁴- (wherein R⁶⁰, R⁶¹, R⁶², R⁶³ and R⁶⁴ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R³⁷ is as defined hereinbefore);

5 15) R³⁶ (wherein R³⁶ is as defined hereinbefore); and

16) $C_{1-3}\text{alkyl}X^{10}C_{1-3}\text{alkyl}R^{36}$ (wherein X^{10} and R³⁶ are as defined hereinbefore).

Suitable pharmaceutically acceptable salts of compounds of formula (I) include acid addition salts such as methanesulfonate, fumarate, hydrochloride, hydrobromide, citrate, maleate and salts formed with phosphoric and sulphuric acid. A preferred
10 pharmaceutically acceptable salt is a hydrochloride salt.

The alkyl portion of the alkyl, alkoxy, alkanoyloxy, alkoxymethyl, alkanoyloxymethyl, alkylsulphanyl, alkylsulphonyl, alkylsulfonamido, carboalkoxy, carboalkyl, alkanoylamino aminoalkyl, alkylaminoalkyl, N,N-dicycloalkylaminoalkyl, hydroxyalkyl, and alkoxyalkyl substituents include both straight chain as well as branched
15 carbon chains. The cycloalkyl portions of N-cycloalkyl-N-alkylaminoalkyl and N,N-dicycloalkylaminoalkyl substituents include both simple carbocycles as well as carbocycles containing alkyl substituents. The alkenyl portion of the alkenyl, alkenoyloxymethyl, alkenyloxy, alkenylsulfonamido, substituents include both straight chain as well as branched carbon chains and one or more sites of unsaturation. The alkynyl portion of the
20 alkynyl, alkynoyloxymethyl, alkynylsulfonamido, alkynyloxy, substituents include both straight chain as well as branched carbon chains and one or more sites of unsaturation. Carboxy is defined as a -CO₂H radical. Carboalkoxy of 2-7 carbon atoms is defined as a -CO₂R" radical, where R" is an alkyl radical of 1-6 carbon atoms. Carboalkyl is defined as a -COR" radical, where R" is an alkyl radical of 1-6 carbon atoms. Alkanoyloxy is
25 defined as a -OCOR" radical, where R" is an alkyl radical of 1-6 carbon atoms. Alkanoyloxymethyl is defined as R"CO₂CH₂- radical, where R" is an alkyl radical of 1-6 carbon atoms. Alkoxymethyl is defined as R"OCH₂- radical, where R" is an alkyl radical of 1-6 carbon atoms. Alkylsulphanyl is defined as R"SO- radical, where R" is an alkyl radical of 1-6 carbon atoms. Alkylsulphonyl is defined as R"SO₂- radical, where R" is
30 alkyl radical of 1-6 carbon atoms. Alkylsulfonamido, alkenylsulfonamido, alkynylsulfonamido are defined as R"SO₂NH- radical, where R" is an alkyl radical of 1-6 carbon atoms, an alkenyl radical of 2-6 carbon atoms, or an alkynyl radical of 2-6 carbon

atoms, respectively. N-alkylcarbamoyl is defined as $R''\text{NHCO-}$ radical, where R'' is an alkyl radical of 1-6 carbon atoms. N,N-dialkylcarbamoyl is defined as $R'' R'\text{NCO-}$ radical, where R'' is an alkyl radical of 1-6 carbon atoms, R' is an alkyl radical of 1-6 carbon atoms and R'' and R' may be the same or different. When X is substituted, it is preferred that it is mono-, di-, or tri-substituted, with monosubstituted being most preferred. It is preferred that of the substituents, R_1 , R_2 , R_3 and R_4 at least one is hydrogen and it is most preferred that two or three be hydrogen. An azacycloalkyl-N-alkyl substituent refers to a monocyclic heterocycle that contains a nitrogen atom on which is substituted a straight or branched chain alkyl radical. A morpholino-N-alkyl substituent is a morpholine ring substituted on the nitrogen atom with a straight or branch chain alkyl radical. A piperazino-N-alkyl substituent is a piperazine ring substituted on one of the nitrogen atoms with a straight or branch chain alkyl radical. A N-alkyl-piperidino-N-alkyl substituent is a piperidine ring substituted on one of the nitrogen atoms with a straight or branched chain alkyl group and on the other nitrogen atom with a straight or branch chain alkyl radical.

When any group contains an alkyl portion, the alkyl portion contains preferably 1-6 carbon atoms, more preferably 1-4 carbon atoms, particularly methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl or tert-butyl. When any group contains an alkenyl or alkynyl portion, the alkenyl or alkynyl portion contains preferably 2-6 carbon atoms, more preferably 2-4 carbon atoms.


The compounds of this invention may contain an asymmetric carbon; in such cases, the compounds of this invention cover the racemate and the individual R and S enantiomers, and in the case where more than one asymmetric carbon exists, the individual diastereomers, their racemates and individual enantiomers.

Examples of substituents for aryl groups R^9 or optional substituents for carbocyclic or heterocyclic groups R^9 include one or more groups selected from hydroxy; halo; nitro; cyano; carboxy; C_{1-6} alkoxy; C_{1-6} alkyl; C_{2-6} alkenyl; C_{2-6} alkynyl; C_{2-6} alkenyloxy; C_{2-6} alkynyloxy; C_{3-6} cycloalkyl; amino; mono- or di- C_{1-6} alkyl amino; heterocyclyl optionally substituted with C_{1-6} alkyl or oxo; $C(O)R^a$, $C(O)OR^a$, $S(O)_dR^a$, $NR^aC(O)R^b$; $C(O)NR^aS(O)_dR^b$, $C(O)NR^aR^b$; $NR^aC(O)NR^bR^c$; $NR^aS(O)_dR^b$ or $N(S(O)_dR^b)S(O)_dR^c$ where d is 0, 1 or 2 and R^a , R^b and R^c are independently selected from hydrogen, C_{1-6} alkyl, aryl, C_{3-6} cycloalkyl or heterocyclyl, and wherein any alkyl, alkenyl or alkynyl group

or moiety contained within the substituent one R^9 may themselves be optionally substituted with one or more groups selected from hydroxy; cyano; nitro; halo; carboxy; carboalkoxy of 2-7 carbon atoms, C_{3-6} cycloalkyl, heterocyclyl optionally substituted with C_{1-6} alkyl or oxo; $C(O)R^d$, $C(O)OR^d$ NR^dR^e , $S(O)_eR^d$, $NR^dC(O)R^e$; $C(O)NR^dR^e$;

- 5 $NR^dC(O)NR^eR^f$; $NR^dS(O)_eR^e$ where e is 0, 1 or 2 and R^d , R^e and R^f are independently selected from hydrogen or C_{1-6} alkyl optionally substituted with one or more groups selected from hydroxy; cyano; nitro; halo; carboxy; carboalkoxy of 2-7 carbon atoms, C_{3-6} cycloalkyl, heterocyclyl optionally substituted with C_{1-6} alkyl or oxo; $C(O)R^g$, $C(O)OR^g$ NR^gR^h , $S(O)_eR^g$, $NR^hC(O)R^g$; $C(O)NR^gR^h$; $NR^gC(O)NR^hR^i$; $NR^gS(O)_eR^h$ where e is as
10 defined above and R^g , R^h and R^i are independently selected from hydrogen or C_{1-6} alkyl. Alternatively, two substituents on adjacent atoms may be joined to form the second ring of a bicyclic ring system wherein the said second ring is optionally substituted with one or more of the groups listed above for R^9 and optionally contains one or more heteroatoms.

- In some embodiments, the level of substitution on the group R^9 is a chain
15 substituted with complex. Thus, for example, a substituent may comprise an substituted alkyl chain which is optionally interposed with heteroatoms such as groups of sub-formula (i)



- where X^a , X^b and X^c are independently selected from any of the groups listed
20 above for X^1 ,

R^{70} and R^{71} are independently selected from C_{1-6} alkylene, C_{2-6} alkenylene or C_{2-6} alkynylene groups any of which may be optionally substituted with hydroxy; cyano; nitro; halo; carboxy, carboalkoxy of 2-7 carbon atoms or C_{3-6} cycloalkyl;

- R^{72} is hydrogen or an C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl group any of which may
25 be optionally substituted with hydroxy; cyano; nitro; halo; carboxy or C_{3-6} cycloalkyl;
and q and s are independently 0 or 1.

Preferably R^9 is an optionally substituted alkoxy group and most preferably, R^9 is a substituted alkoxy group.

- A particular example of compounds of formula (I) are compounds of formula (IA)
30 which are compounds of formula (I) as defined above provided that R^7 is a group $(CH_2)_mR^9$ where m is 0, or an integer of from 1-3 and R^9 is a substituted aryl or substituted cycloalkyl ring of up to 10 carbon atoms, wherein the substituents comprise at

least one alkoxy group of 1-6 carbon atoms and optionally one or more further substituents, or R^9 is a heterocyclic ring containing 1 or 2 oxygen atoms and optionally one or more substituents, and where R^1 , R^2 , R^3 or R^4 are a group $R^{13}-X^1-(CH_2)_x$ wherein x is 0 to 3, X^1 represents -O-, -CH₂-, -OCO-, carbonyl, -S-, -SO-, -SO₂-, -NR¹⁴CO-,
 5 SO₂NR¹⁶-, -NR¹⁷SO₂- or -NR¹⁸- (wherein R^{14} , R^{15} , R^{16} , R^{17} , R^{18} and R^{13} are as defined above).


Suitable examples of groups Y are -NH-. Suitably X is oxygen.

Preferably n is 0.

Particular examples of groups R^9 include phenyl or cycloalkyl of from 3-8 and
 10 preferably of 6 carbon atoms which are substituted at the position alpha with a alkoxy group, in particular methoxy.

When R^9 is substituted phenyl or cycloalkyl, m is preferably 0.

Examples of heterocyclic rings R^9 include 3- 7 membered rings, up to two of which may be oxygen atoms. Such groups include:




15 where each R^{65} is independently selected from hydrogen or C₁₋₆alkyl and especially methyl. In such compounds, m is suitably 1, 2 or 3.

Other examples of heterocyclic groups R^9 include pyridyl, thiazolyl, pyrazinyl, pyrimidinyl, oxadiazole.

20 Suitable further substituents for R^7 include those listed above for pyridyl, pyrimidinyl and phenyl groups R^6 .

Thus a preferred sub-group of compounds of formula (I) are compounds of formula (II)



(II)

where R^1 , R^2 , R^3 and R^4 are as defined above and R^{66} is C_{1-6} alkyl in particular methyl and R^{67} is selected from hydrogen, halogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, azido, hydroxyalkyl of 1-6 carbon atoms, halomethyl, alkoxyethyl of 2-7 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkoxy of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, hydroxy, trifluoromethyl, cyano, nitro, carboxy, carboalkoxy of 2-7 carbon atoms, carboalkyl of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzoyl, benzyl, amino, alkylamino of 1-6 carbon atoms, dialkylamino of 2 to 12 carbon atoms, phenylamino, benzylamino, alkanoylamino of 1-6 carbon atoms, alkenoylamino of 3-8 carbon atoms, alkynoylamino of 3-8 carbon atoms, and benzoylamino.

Suitably R^{66} is C_{1-6} alkyl such as methyl. Preferably however it is a substituted C_{1-4} alkyl group, wherein the substituents are selected from hydroxy, NR^dR^e , $S(O)_eR^d$, $NR^dC(O)R^e$; $C(O)NR^dR^e$; $NR^dC(O)NR^eR^f$; $NR^dS(O)_eR^e$ where e , R^d , R^e and R^f are as defined above.

Preferably R^{67} is hydrogen.

Examples of preferred groups for R^1 , R^2 , R^3 and R^4 are set out in WO 98/43960. Preferably x is 0. Conveniently R^{13} is selected from one of the following sixteen groups:

- 1) C_{1-5} alkyl which may be unsubstituted or substituted with one or more fluorine atoms, or
- C_{2-5} alkyl which may be unsubstituted or substituted with one or more groups selected from hydroxy and amino;

- 2) $C_{2-3}alkylX^2COR^{19}$ (wherein X^2 is as defined hereinbefore and R^{19} represents $-NR^{21}R^{22}-$ or $-OR^{23}-$ (wherein R^{21} , R^{22} and R^{23} which may be the same or different each represents hydrogen, $C_{1-2}alkyl$ or $C_{1-2}alkoxyethyl$));
- 3) $C_{2-4}alkylX^3R^{24}$ (wherein X^3 is as defined hereinbefore and R^{24} represents hydrogen, $C_{1-3}alkyl$, cyclopentyl, cyclohexyl or a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which $C_{1-3}alkyl$ group may bear one or two substituents selected from oxo, hydroxy, halogeno and $C_{1-3}alkoxy$ and which cyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, $C_{1-3}alkyl$, $C_{1-3}hydroxyalkyl$ and $C_{1-3}alkoxy$);
- 4) $C_{2-3}alkylX^4C_{2-3}alkylX^5R^{30}$ (wherein X^4 and X^5 are as defined hereinbefore and R^{30} represents hydrogen or $C_{1-3}alkyl$);
- 5) $C_{1-5}alkylR^{70}$ (wherein R^{70} is a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which heterocyclic group is linked to $C_{1-5}alkyl$ through a carbon atom and which heterocyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, $C_{1-3}alkyl$, $C_{1-3}hydroxyalkyl$ and $C_{1-3}alkoxy$) or $C_{2-5}alkylR^{71}$ (wherein R^{71} is a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms of which one is N and the other is selected independently from O, S and N, which heterocyclic group is linked to $C_{2-5}alkyl$ through a nitrogen atom and which heterocyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, $C_{1-3}alkyl$, $C_{1-3}hydroxyalkyl$ and $C_{1-3}alkoxy$);
- 6) $(CH_2)_qX^6R^{37}$ (wherein X^6 is as defined hereinbefore; q is an integer from 0 to 4 if X^6 is a direct bond and q is 0, 2 or 3 if X^6 is other than a direct bond; and R^{37} is a phenyl group, a pyridone group or a 5 or 6 membered aromatic heterocyclic group with 1 to 3 heteroatoms selected from O, N and S, of which preferably one is N, which phenyl group, pyridone group or aromatic heterocyclic group may be substituted as hereinbefore defined, advantageously substituted with up to 2 substituents as hereinbefore defined, more preferably substituted with one substituent selected from the group of substituents as hereinbefore defined);
- 7) $C_{4-5}alkenylR^{72}$ (wherein R^{72} represents R^{70} or R^{71} as defined hereinbefore);
- 8) $C_{4-5}alkynylR^{72}$ (wherein R^{72} represents R^{70} or R^{71} as defined hereinbefore);

- 9) X^7R^{47} (wherein X^7 is as defined hereinbefore and R^{47} represents C_{1-3} alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro and amino);
- 10) C_{3-5} alkenyl R^{37} (wherein R^{37} is as defined hereinbefore);
- 5 11) C_{3-5} alkynyl R^{37} (wherein R^{37} is as defined hereinbefore);
- 12) C_{4-5} alkenyl X^8R^{37} (wherein X^8 and R^{37} are as defined hereinbefore);
- 13) C_{4-5} alkynyl X^9R^{30} (wherein X^9 and R^{30} are as defined hereinbefore);
- 14) C_{1-3} alkyl $X^{10}C_{1-3}$ alkyl R^{37} (wherein X^{10} and R^{37} are as defined hereinbefore);
- 15) R^{36} (wherein R^{36} is as defined hereinbefore); and
- 10 16) C_{1-3} alkyl $X^{11}C_{1-3}$ alkyl R^{36} (wherein X^{11} and R^{36} are as defined hereinbefore).

Advantageously R^{13} is selected from one of the following eleven groups:

- 1) C_{1-4} alkyl which may be unsubstituted or substituted with one or more fluorine atoms, or
- C_{2-4} alkyl which may be unsubstituted or substituted with one or two groups selected from
- 15 hydroxy and amino;
- 2) C_{2-3} alkyl X^2COR^{19} (wherein X^2 is as defined hereinbefore and R^{19} represents $-NR^{21}R^{22}-$ or $-OR^{23}-$ (wherein R^{21} , R^{22} and R^{23} which may be the same or different each represents hydrogen, C_{1-2} alkyl or C_{1-2} alkoxyethyl));
- 3) C_{2-3} alkyl X^3R^{24} (wherein X^3 is as defined hereinbefore and R^{24} is a group selected from
- 20 C_{1-3} alkyl, cyclopentyl, cyclohexyl, pyrrolidinyl and piperidinyl which group is linked to X^3 through a carbon atom and which C_{1-3} alkyl group may bear one or two substituents selected from oxo, hydroxy, halogeno and C_{1-2} alkoxy and which cyclopentyl, cyclohexyl, pyrrolidinyl or piperidinyl group may carry one substituent selected from oxo, hydroxy, halogeno, C_{1-2} alkyl, C_{1-2} hydroxyalkyl and C_{1-2} alkoxy);
- 25 4) C_{2-3} alkyl X^4C_{2-3} alkyl X^5R^{30} (wherein X^4 and X^5 are as defined hereinbefore) and R^{30} represents hydrogen or C_{1-2} alkyl);
- 5) C_{1-4} alkyl R^{70} (wherein R^{70} is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, 1,3-dioxolan-2-yl, 1,3-dioxan-2-yl, 1,3-dithiolan-2-yl and 1,3-dithian-2-yl, which group is linked to C_{1-4} alkyl through a carbon atom and which group may carry one or two
- 30 substituents selected from oxo, hydroxy, halogeno, C_{1-2} alkyl, C_{1-2} hydroxyalkyl and C_{1-2} alkoxy) or C_{2-4} alkyl R^{71} (wherein R^{71} is a group selected from morpholino, thiomorpholino, pyrrolidin-1-yl, piperazin-1-yl and piperidino which group may carry one

or two substituents selected from oxo, hydroxy, halogeno, C₁₋₂alkyl, C₁₋₂hydroxyalkyl and C₁₋₂alkoxy); and

6) (CH₂)_qX⁶R³⁷ (wherein X⁶ is as defined hereinbefore; q is an integer from 1 to 3 if X⁶ is a direct bond and q is 2 or 3 if X⁶ is other than a direct bond; and R³⁷ is a phenyl group, a pyridone group or a 5 or 6 membered aromatic heterocyclic group with 1 to 2 heteroatoms selected from O, N and S, of which preferably one is N, which phenyl group, pyridone group or aromatic heterocyclic group may be substituted as hereinbefore defined, preferably substituted with one substituent selected from hydroxy, halogeno, C₁₋₂alkyl, C₁₋₂alkoxy, C₁₋₂hydroxyalkyl, C₁₋₂hydroxyalkoxy, carboxy, cyano, -CONR⁴³R⁴⁴ and - NR⁴⁵COR⁴⁶ (wherein R⁴³, R⁴⁴, R⁴⁵ and R⁴⁶, which may be the same or different, each represents hydrogen or C₁₋₂alkyl));

7) C₄₋₅alkenylR⁷¹ (wherein R⁷¹ is as defined hereinbefore);

8) C₄₋₅alkynylR⁷¹ (wherein R⁷¹ is as defined hereinbefore);

9) C₁₋₃alkylX¹⁰C₁₋₃alkylR³⁷ (wherein X¹⁰ and R³⁷ are as defined hereinbefore);

10) R³⁶ (wherein R³⁶ is as defined hereinbefore); and

11) C₁₋₃alkylX¹¹C₁₋₃alkylR³⁶ (wherein X¹¹ and R³⁶ are as defined hereinbefore).

Preferably R¹³ is selected from one of the following nine groups:

1) C₁₋₃alkyl which may be unsubstituted or substituted with one or more fluorine atoms, or

2) C₂₋₃alkyl which may be unsubstituted or substituted with one or two groups selected from hydroxy and amino;

2) 2-(3,3-dimethylureido)ethyl, 3-(3,3-dimethylureido)propyl, 2-(3-methylureido)ethyl, 3-(3-methylureido)propyl, 2-ureidoethyl, 3-ureidopropyl, 2-(N,N-dimethylcarbamoyloxy)ethyl, 3-(N,N-dimethylcarbamoyloxy)propyl, 2-(N-methylcarbamoyloxy)ethyl, 3-(N-methylcarbamoyloxy)propyl, 2-(carbamoyloxy)ethyl, 3-(carbamoyloxy)propyl;

3) C₂₋₃alkylX³R²⁴ (wherein X³ is as defined hereinbefore and R²⁴ is a group selected from C₁₋₂alkyl, cyclopentyl, cyclohexyl, pyrrolidinyl and piperidinyl which group is linked to X³ through a carbon atom and which C₁₋₂alkyl group may bear one or two substituents selected from oxo, hydroxy, halogeno and C₁₋₂alkoxy and which cyclopentyl, cyclohexyl, pyrrolidinyl or piperidinyl group may carry one substituent selected from oxo, hydroxy, halogeno, C₁₋₂alkyl, C₁₋₂hydroxyalkyl and C₁₋₂alkoxy);

- 4) $C_{2-3}alkylX^4C_{2-3}alkylX^5R^{32}$ (wherein X^4 and X^5 are as defined hereinbefore) and R^{30} represents hydrogen or $C_{1-2}alkyl$);
- 5) $C_{1-2}alkylR^{70}$ (wherein R^{70} is a group selected from pyrrolidinyl, piperazinyl, piperidinyl, 1,3-dioxolan-2-yl, 1,3-dioxan-2-yl, 1,3-dithiolan-2-yl and 1,3-dithian-2-yl, which group is linked to $C_{1-2}alkyl$ through a carbon atom and which group may carry one substituent selected from oxo, hydroxy, halogeno, $C_{1-2}alkyl$, $C_{1-2}hydroxyalkyl$ and $C_{1-2}alkoxy$) or $C_{2-3}alkylR^{59}$ (wherein R^{59} is a group selected from morpholino, thiomorpholino, piperidino, piperazin-1-yl and pyrrolidin-1-yl which group may carry one or two substituents selected from oxo, hydroxy, halogeno, $C_{1-2}alkyl$, $C_{1-2}hydroxyalkyl$ and $C_{1-2}alkoxy$);
- 10 6) $(CH_2)_qX^6R^{37}$ (wherein X^6 is as defined hereinbefore; q is an integer from 1 to 3 if X^6 is a direct bond and q is 2 or 3 if X^6 is other than a direct bond; and R^{37} is a group selected from phenyl, a pyridone group, pyridyl, imidazolyl, thiazolyl, thienyl, triazolyl and pyridazinyl, preferably selected from phenyl, a pyridone group, pyridyl, imidazolyl, thiazolyl and triazolyl which group may be substituted with one substituent selected from
- 15 hydroxy, halogeno, $C_{1-2}alkyl$, $C_{1-2}alkoxy$, $C_{1-2}hydroxyalkyl$, $C_{1-2}hydroxyalkoxy$, carboxy, cyano, $-CONR^{43}R^{44}$ and $-NR^{45}COR^{46}$ (wherein R^{43} , R^{44} , R^{45} and R^{46} are as defined hereinbefore);
- 7) $C_{1-3}alkylX^{10}C_{1-3}alkylR^{37}$ (wherein X^{10} and R^{37} are as defined hereinbefore);
- 8) R^{36} (wherein R^{36} is as defined hereinbefore); and
- 20 9) $C_{1-3}alkylX^{11}C_{1-3}alkylR^{36}$ (wherein X^{11} and R^{36} are as defined hereinbefore).
- More preferably R^{13} represents 2-methylthiazol-4-ylmethyl, 2-acetamidothiazol-4-ylmethyl, 1-methylimidazol-2-ylmethyl, 4-pyridylmethyl, 2-(4-pyridyl)ethyl, 3-(4-pyridyl)propyl, 2-((N-(1-methylimidazol-4-ylsulphonyl)-N-methyl)amino)ethyl, 2-((N-(3-morpholinopropylsulphonyl)-N-methyl)amino)ethyl, 2-((N-methyl-N-4-
- 25 pyridyl)amino)ethyl, 2-(4-oxidomorpholino)ethyl, 3-(4-oxidomorpholino)propyl, 2-(4-oxo-1,4-dihydro-1-pyridyl)ethyl, 3-(4-oxo-1,4-dihydro-1-pyridyl)propyl, methyl, ethyl, trifluoromethyl, 2,2,2-trifluoroethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2-(N,N-dimethylsulphamoyl)ethyl, 2-(N-methylsulphamoyl)ethyl, (1,3-dioxolan-2-yl)methyl, 2-(1,3-dioxolan-2-yl)ethyl, 2-(2-methoxyethylamino)ethyl, 2-(2-hydroxyethylamino)ethyl, 3-
- 30 (2-methoxyethylamino)propyl, 3-(2-hydroxyethylamino)propyl, 2-(1,2,4-triazol-1-yl)ethyl, 2-(1,2,4-triazol-4-yl)ethyl, 3-(1,2,4-triazol-1-yl)propyl, 3-(1,2,4-triazol-4-yl)propyl, 2-(4-pyridyloxy)ethyl, 3-(4-pyridyloxy)propyl, 2-(4-pyridylamino)ethyl, 3-(4-

pyridylamino)propyl, 2-(2-methylimidazol-1-yl)ethyl, 3-(2-methylimidazol-1-yl)propyl, 2-(5-methyl-1,2,4-triazol-1-yl)ethyl, 3-(5-methyl-1,2,4-triazol-1-yl)propyl, morpholino, N-methylpiperazinyl, piperazinyl, 2-(N,N-dimethylamino)ethyl, 3-(N,N-dimethylamino)propyl, 2-morpholinoethyl, 3-morpholinopropyl, 2-piperidinoethyl, 3-piperidinopropyl, 2-(piperazin-1-yl)ethyl, 3-(piperazin-1-yl)propyl, 2-(pyrrolidin-1-yl)ethyl, 3-(pyrrolidin-1-yl)propyl, 2-methoxyethyl, 3-methoxypropyl, 2-(imidazol-1-yl)ethyl, 2-(1,2,3-triazol-1-yl)ethyl, 2-(1,2,3-triazol-2-yl)ethyl, 3-(imidazol-1-yl)propyl, 3-(1,2,3-triazol-1-yl)propyl, 3-(1,2,3-triazol-2-yl)propyl, 2-thiomorpholinoethyl, 3-thiomorpholinopropyl, 2-(1,1-dioxothiomorpholino)ethyl, 3-(1,1-dioxothiomorpholino)propyl, 2-(2-methoxyethoxy)ethyl, 2-(4-methylpiperazin-1-yl)ethyl, 3-(4-methylpiperazin-1-yl)propyl, 3-(methylsulphinyl)propyl, 3-(methylsulphonyl)propyl, 2-(methylsulphinyl)ethyl, benzyl, 2-sulphamoyl ethyl or 2-(methylsulphonyl)ethyl.

Especially R¹³ represents methyl, ethyl, trifluoromethyl, 2,2,2-trifluoroethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2-methoxyethyl, 3-methoxypropyl, 2-(methylsulphinyl)ethyl, 2-(methylsulphonyl)ethyl, 2-(N,N-dimethylsulphamoyl)ethyl, 2-(N-methylsulphamoyl)ethyl, 2-sulphamoyl ethyl, 2-(N,N-dimethylamino)ethyl, 3-(N,N-dimethylamino)propyl, 2-morpholinoethyl, 3-morpholinopropyl, 2-piperidinoethyl, 3-piperidinopropyl, 2-(piperazin-1-yl)ethyl, 3-(piperazin-1-yl)propyl, 2-(pyrrolidin-1-yl)ethyl, 3-(pyrrolidin-1-yl)propyl, (1,3-dioxolan-2-yl)methyl, 2-(1,3-dioxolan-2-yl)ethyl, 2-(2-methoxyethylamino)ethyl, 2-(2-hydroxyethylamino)ethyl, 3-(2-methoxyethylamino)propyl, 3-(2-hydroxyethylamino)propyl, 2-methylthiazol-4-ylmethyl, 2-acetamidothiazol-4-ylmethyl, 1-methylimidazol-2-ylmethyl, 2-(imidazol-1-yl)ethyl, 2-(1,2,3-triazol-1-yl)ethyl, 2-(1,2,3-triazol-2-yl)ethyl, 2-(1,2,4-triazol-1-yl)ethyl, 2-(1,2,4-triazol-4-yl)ethyl, 4-pyridylmethyl, 2-(4-pyridyl)ethyl, 3-(4-pyridyl)propyl, 3-(3-pyridyl)propyl, benzyl, 2-(4-pyridyloxy)ethyl, 2-(4-pyridylamino)ethyl, or 2-(4-oxo-1,4-dihydro-1-pyridyl)ethyl.

More especially R¹³ represents methyl, ethyl, trifluoromethyl, 2,2,2-trifluoroethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2-methoxyethyl, 3-methoxypropyl, 2-(methylsulphinyl)ethyl, 2-(methylsulphonyl)ethyl, 2-(N,N-dimethylsulphamoyl)ethyl, 2-(N-methylsulphamoyl)ethyl, 2-sulphamoyl ethyl, 2-(N,N-dimethylamino)ethyl, 3-(N,N-dimethylamino)propyl, 2-morpholinoethyl, 3-morpholinopropyl, 2-piperidinoethyl, 3-piperidinopropyl, 2-(piperazin-1-yl)ethyl, 3-(piperazin-1-yl)propyl, 2-(pyrrolidin-1-yl)ethyl,

3-(pyrrolidin-1-yl)propyl, (1,3-dioxolan-2-yl)methyl, 2-(1,3-dioxolan-2-yl)ethyl, 2-(2-methoxyethylamino)ethyl, 2-(2-hydroxyethylamino)ethyl, 3-(2-methoxyethylamino)propyl, 3-(2-hydroxyethylamino)propyl, 2-methylthiazol-4-ylmethyl, 2-acetamidothiazol-4-ylmethyl, 1-methylimidazol-2-ylmethyl, 2-(imidazol-1-yl)ethyl, 2-(1,2,3-triazol-1-yl)ethyl,
 5 2-(1,2,3-triazol-2-yl)ethyl, 2-(1,2,4-triazol-1-yl)ethyl, 2-(1,2,4-triazol-4-yl)ethyl, 4-pyridylmethyl, 2-(4-pyridyl)ethyl, 3-(4-pyridyl)propyl, benzyl, 2-(4-pyridyloxy)ethyl, 2-(4-pyridylamino)ethyl, or 2-(4-oxo-1,4-dihydro-1-pyridyl)ethyl.

In particular R^1 and R^4 are suitably hydrogen.

Examples of preferred groups for R^2 include C_{1-6} alkoxy such as methoxy.


10 The group R^3 is suitably selected from hydrogen or C_{1-6} alkoxy.

Preferably both R^2 and R^3 are C_{1-6} alkoxy and are preferably methoxy.

A further preferred group for R^2 or R^3 is 3-morpholinopropoxy.


Particular examples of compounds of formula (I) are listed in Tables 1, 2 and 3.

In these tables "DMMPO" indicates a 1,6-dimethylmorpholinopropoxy group of formula:




15

"MPO" is morpholinopropoxy group of formula:



20


"MEO" is a morpholinoethoxy group of formula:









and Me is CH_3









25





Table 1






No.	R ²	R ³	X	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸³	R ⁸⁴	R ⁸⁵	R ⁸⁶	R ⁸⁷
1	OMe	OMe	O	OMe	H	H	H	H	H	H	H
2	OMe	OMe	NH	H	H	OMe	H	H	H	H	H
3	OMe	OMe	O	H	OMe	H	H	H	H	Cl	H
4	OMe	OMe	O	OMe	H	H	H	H	H	Cl	H
5	OMe	OMe	O	OMe	H	H	H	OMe	H	H	H
6	OMe	OMe	O	OMe	H	H	H	Me	H	H	H
7	OMe	OMe	O	H	OMe	H	H	H	H	H	H
8	OMe	OMe	O	H	H	OMe	H	H	H	H	H
9	OMe	OCH ₂ C ₆ H ₅	O	OMe	H	H	H	H	H	H	H
10	OMe	OMe	O	OMe	OMe	H	H	H	H	H	H
11	OMe	OMe	O	H	OMe	H	OMe	H	H	H	H
12	OMe	OMe	O	OCH ₂ (Me) ₂	H	H	H	H	H	H	H
13	OMe	OMe	O	CO ₂ Me	H	H	OMe	H	H	H	H
14	MPO	OMe	O	OMe	H	H	H	H	H	H	H
15	OMe	OMe	O	H	OMe	H	Cl	H	H	H	H




No.	R ²	R ³	X	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸³	R ⁸⁴	R ⁸⁵	R ⁸⁶	R ⁸⁷
16	OMe	MPO	O	OMe	H	H	H	H	H	H	H
17	MPO	OMe	O	OMe	H	H	H	H	H	H	H
18		OMe	O	OMe	H	H	H	H	H	H	H
19	MPO	OMe	O	OMe	H	H	H	H	H	H	H
20	$O(CH_2)_3N(Me)_2$	OMe	O	OMe	H	H	H	H	H	H	H
21	MPO	OMe	O	OMe	H	H	H	H	H	H	H
22		OMe	O	OMe	H	H	H	H	H	H	H
23		OMe	O	OMe	H	H	H	H	H	H	H
24	MPO	OMe	O	OMe	H	H	H	H	H	H	H
25	$O(CH_2)_2N(Me)_2$	OMe	O	OMe	H	H	H	H	H	H	H
26	OH	OMe	O	OMe	H	H	H	H	H	H	H
27	OMe	OH	O	OMe	H	H	H	H	H	H	H
28		OMe	O	OMe	H	H	H	H	H	H	H
29	2-thiazolyloxy	OMe	O	OMe	H	H	H	H	H	H	H
30	2-pyrimidinylloxy	OMe	O	OMe	H	H	H	H	H	H	H
31	2-pyridylloxy	OMe	O	OMe	H	H	H	H	H	H	H
32	OMe	OMe	O	OMe	H	H	H	H	H	H	H
33	OMe		O	OMe	H	H	H	H	H	H	H
34		OMe	O	OMe	H	H	H	H	H	H	H


No.	R ²	R ³	X	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸³	R ⁸⁴	R ⁸⁵	R ⁸⁶	R ⁸⁷
35	OMe		O	OMe	H	H	H	H	H	H	H
36	OMe		O	OMe	H	H	H	H	H	H	H
37	OMe		O	OMe	H	H	H	H	H	H	H
38	OMe		O	OMe	H	H	H	H	H	H	H
39	OMe		O	OMe	H	H	H	H	H	H	H
40	OMe		O	OMe	H	H	H	H	H	H	H
41	OMe		O	OMe	H	H	H	H	H	H	H
42	O(CH ₂) ₂ OMe	O(CH ₂) ₂ OMe	O	OMe	H	H	H	H	H	H	H
43	OMe	OMe	O	OMe	H	H	H	H	H	Me	H
44	OMe		O	OMe	H	H	H	H	H	H	H
45	O(CH ₂) ₂ OMe	O(CH ₂) ₂ OMe	O	H	OMe	H	H	H	H	H	H
46	O(CH ₂) ₂ OMe	O(CH ₂) ₂ OMe	O	H	OMe	H	OMe	H	H	H	H
47	O(CH ₂) ₂ OMe	O(CH ₂) ₂ OMe	O	OCH ₂ Me	H	H	H	H	H	H	H





No.	R ²	R ³	X	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸³	R ⁸⁴	R ⁸⁵	R ⁸⁶	R ⁸⁷
48	OMe		O	OMe	H	H	H	H	H	H	H
49	OCH ₂ CO ₂ CH ₂ Me	OMe	O	OMe	H	H	H	H	H	H	H
50	OCH ₂ CF ₃	OMe	O	OMe	H	H	H	H	H	H	H
51	OCH ₂ CH=CH ₂	OMe	O	OMe	H	H	H	H	H	H	H
52	OCH ₂ COOH	OMe	O	OMe	H	H	H	H	H	H	H
53		OMe	O	OMe	H	H	H	H	H	H	H
54	OCH ₂ C≡CH	OMe	O	OMe	H	H	H	H	H	H	H
55	OCH ₂ CH ₂ OMe	OMe	O	OMe	H	H	H	H	H	H	H
56	OMe	OMe	O	OCH ₂ Me	H	H	H	H	H	Me	H
57		OMe	O	OMe	H	H	H	H	H	H	H
58		OMe	O	OMe	H	H	H	H	H	H	H
59	OCH ₂ C(O)NH CH ₂ CH=CH ₂	OMe	O	OMe	H	H	H	H	H	H	H
60	OCH ₂ C(O)NH- Me	OMe	O	OMe	H	H	H	H	H	H	H
61	OCH ₂ C(O)NH- (CH ₂) ₂ OMe	OMe	O	OMe	H	H	H	H	H	H	H
62	OMe	OH	O	OCH ₂ Me	H	H	H	H	H	H	H





No.	R ²	R ³	X	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸³	R ⁸⁴	R ⁸⁵	R ⁸⁶	R ⁸⁷
63	OMe		O	OMe	H	H	H	H	H	H	H
64	OMe		O	OMe	H	H	H	H	H	H	H
65	OMe	OCH ₂ CO ₂ CH ₂ Me	O	OMe	H	H	H	H	H	H	H
66	OMe	OCH ₂ CO ₂ H	O	OMe	H	H	H	H	H	H	H
67	OMe	CCH ₂ C≡CH	O	OMe	H	H	H	H	H	H	H
68	OMe		O	OMe	H	H	H	H	H	H	H
69	OMe	MPO	O	H	OMe	H	H	H	H	H	H
70	OMe	MPO	O	OCH ₂ Me	H	H	H	H	H	H	H
71	NHCO ₂ CH(Me) ₂	OMe	O	OMe	H	H	H	H	H	H	H
72	NH ₂	OMe	O	OMe	H	H	H	H	H	H	H
73	NHSO ₂ Me	OMe	O	OMe	H	H	H	H	H	H	H
74	OMe	OMe	O	F	H	H	H	OMe	H	H	H
75	OMe	OMe	O	H	H	Cl	H	H	H	H	H
76	OMe	OMe	O	H	H	NO ₂	H	H	H	H	H
77	OMe	OMe	O	F	H	F	H	H	H	H	H
78	OMe	OMe	S	Me	H	H	H	H	H	Cl	H
79	OMe	OMe	S	H	H	Cl	H	H	H	H	H
80	OMe	OMe	O	H	H	Me	H	H	H	Cl	H
81	OMe	OMe	O	F	H	H	H	H	H	H	H
82	OMe	OMe	O	Me	H	H	H	H	H	H	H
83	OMe	OMe	O	H	H	Cl	H	H	Cl	H	Cl
84	OMe	OMe	S	H	CO ₂ Me	H	H	H	H	H	H








No.	R ²	R ³	X	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸³	R ⁸⁴	R ⁸⁵	R ⁸⁶	R ⁸⁷
85	OMe	OMe	O	SMe	H	H	H	H	H	H	H
86	OMe	OMe	O	CN	H	H	H	H	H	H	H
87	OMe	OMe	O	H	I	H	H	H	H	H	H
88	OMe	OMe	O	Br	H	H	H	H	H	H	H
89	OMe	OMe	O	H	Br	H	H	H	H	H	H
90	OMe	OMe	O	H	F	H	H	H	H	H	H
91	OMe	OMe	O	H	Cl	H	H	H	H	H	H
92	OMe	OMe	O	I	H	H	H	H	H	H	H
93	OMe	OMe	O	Cl	H	H	Cl	H	H	H	H
94	OMe	OMe	O	Cl	H	H	H	H	H	H	H
95	OMe	OMe	O	H	NHC(O)Me	H	H	H	H	H	H
96	OMe	OMe	O	OH	H	H	H	H	H	H	H
97	OMe	OMe	O	C(O) ₂ CH ₂ C ₆ H ₅	H	H	H	H	H	H	H
98	OMe	OMe	O	OCF ₃	H	H	H	H	H	H	H
99	OMe	OMe	O	H	CF ₃	H	H	H	H	H	H
100	OMe	OMe	O	C(O) ₂ H	H	H	H	H	H	H	H
101	OMe	OMe	O	H	NHCH ₂ Me	H	H	H	H	H	H
102	COOH	OMe	O	OMe	H	H	H	H	H	H	H
103	OMe	OMe	O	C(O) ₂ Me	H	H	H	H	H	H	H
104	OMe	OMe	O	H	N(CH ₂ Me) ₂	H	H	H	H	H	H
105	OMe	OMe	O	H	CN	H	H	H	H	H	H
106	OMe	OMe	O	H	NHC(O)Me	H	H	H	Me	H	H
107	OMe	OMe	O	H	CN	H	H	H	Me	H	H
108	OH	OMe	O	H	OCF ₂ CHF ₂	H	H	H	H	H	H
109	OH	OMe	O	OCH ₂ CCH	H	H	H	H	H	H	H
110	OH	OMe	O	CN	H	H	H	H	H	H	H




No.	R ²	R ³	X	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸³	R ⁸⁴	R ⁸⁵	R ⁸⁶	R ⁸⁷
111	OMe	OMe	O	N(Me) ₂	H	H	H	H	H	H	H
112	OMe	OMe	O	H	N(Me) ₂	H	H	H	H	H	H
113	OMe	OH	O	H	OCF ₂ CF ₂ H	H	H	H	H	H	H
114		OMe	O	OCH ₂ C≡CH	H	H	H	H	H	H	H
115	MPO	OMe	O	H	OCF ₂ CF ₂ H	H	H	H	H	H	H
116	OMe	OMe	O	CONH ₂	H	H	H	H	H	H	H
117	OMe	OMe	O	H	NH(Me)	H	H	H	H	H	H
118		OMe	O	H	OCF ₂ CF ₂ H	H	H	H	H	H	H
119	O(CH ₂) ₃ N(Me) ₂	OMe	O	H	OCF ₂ CF ₂ H	H	H	H	H	H	H
120	MPO	OMe	O	OH	H	H	H	H	H	H	H
121	O(CH ₂) ₃ N(Me) ₂	OMe	O	OH	H	H	H	H	H	H	H
122	OH	OMe	O	OCH ₂ CN	H	H	H	H	H	H	H
123	OH	OMe	O	OCH ₂ CH ₂ OH	H	H	H	H	H	H	H
124	OMe	OH	O	OCH ₂ CN	H	H	H	H	H	H	H
125	OMe	OH	O	OCH ₂ CH ₂ OH	H	H	H	H	H	H	H
126	OMe	OH	O	OCH ₂ CH=CH ₂	H	H	H	H	H	H	H
127		OMe	O	OCH ₂ CH=CH ₂	H	H	H	H	H	H	H
128	O(CH ₂) ₃ N(Me) ₂	OMe	O	OCH ₂ CH=CH ₂	H	H	H	H	H	H	H
129	OH	OMe	O	OCH ₂ CONHMe	H	H	H	H	H	H	H
130	OMe	OH	O	CN	H	H	H	H	H	H	H
131	OCH ₂ C≡CH	OMe	O	CN	H	H	H	H	H	H	H
132	OMe	OCH ₂ C≡CH	O	CN	H	H	H	H	H	H	H










No.	R ²	R ³	X	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸³	R ⁸⁴	R ⁸⁵	R ⁸⁶	R ⁸⁷
133	O(CH ₂) ₂ OMe	O(CH ₂) ₂ OMe	O	H	NHCH ₂ Me	H	H	H	H	H	H
134	OMe	MPO	O	H	NHCH ₂ Me	H	H	H	H	H	H
135	OMe	OMe	O		H	H	H	H	H	H	H
136	OMe	OMe	O	S(O)Me	H	H	H	H	H	H	H
137	MPO	OMe	O	OCH ₂ CH ₂ OH	H	H	H	H	H	H	H
138	OMe	MPO	O	OCH ₂ CH ₂ OH	H	H	H	H	H	H	H
139	OMe	MPO	O	CN	H	H	H	H	H	H	H
140	OMe	OMe	O	S(O) ₂ Me	H	H	H	H	H	H	H
141	OMe	MPO	O	F	H	H	H	H	H	H	H
142	OMe	MPO	O	OCH ₂ CONHMe	H	H	H	H	H	H	H
143	MPO	OMe	O	OCH ₂ CONHMe	H	H	H	H	H	H	H
144	OMe	OMe	O	F	F	H	H	H	H	H	H
145	OMe	MPO	O	H	F	H	H	H	H	H	H
146	OMe	OMe	O	H	OCF ₂ CF ₂ H	H	H	H	H	H	H
147	OMe	OMe	O	F	H	H	H	F	H	H	H
148	OMe	OMe	O	F	H	H	F	H	H	H	H
149	OMe	OMe	O	H	F	H	F	H	H	H	H
150	OMe	OH	O	OCH ₂ CO ₂ (CH ₂) ₂ Me	H	H	H	H	H	H	H
151	OMe	OH	O	OCH ₂ CONH(CH ₂) ₂ Cl	H	H	H	H	H	H	H
152	OMe	OH	O	OCH ₂ CONH(CH ₂) ₂ -OH	H	H	H	H	H	H	H
153	OMe	OMe	O	OCH ₂ CONH(CH ₂) ₂ -OH	H	H	H	H	H	H	H
154	OMe	MPO	O	OCF ₂ CF ₂ H	H	H	H	H	H	H	H








No.	R ²	R ³	X	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸³	R ⁸⁴	R ⁸⁵	R ⁸⁶	R ⁸⁷
155	OMe	OH	O	F	H	H	H	H	Me	H	H
156	OMe	OMe	O	H	NCONH-Me	H	H	H	H	H	H
157	OMe	OMe	O	H	OCF ₃	H	H	H	H	H	H
158	OMe	OMe	O	CO ₂ Me	F	H	H	H	H	H	H
159	OMe	OMe	O	OCH ₂ CH ₂ OH	H	H	H	H	H	H	H
160	OMe	OMe	O	F	F	H	H	F	H	H	H
161	OMe	OMe	O	OCH ₂ CONHMe	H	H	H	H	H	H	H
162	OMe	OMe	O	OCF ₃	H	H	H	H	H	H	H
163	OMe	MPO	O	H	H	F	H	H	H	H	H
164	OMe	MPO	O	F	H	F	H	H	H	H	H
165	OMe	MPO	O	CN	H	H	H	H	Me	H	H
166	OMe	OMe	O		H	H	H	H	H	H	H
167	OMe	OMe	O	H		H	H	H	H	H	H
168	OMe	MPO	O	CH ₂ CONHMe	H	H	H	H	H	H	H
169	OMe	MPO	O	CH ₂ CO ₂ (CH ₂) ₂ Me	H	H	H	H	H	H	H
170	OMe	MPO	O	OCH ₂ CO ₂ H	H	H	H	H	H	H	H
171	OMe	OMe	O		H	H	H	H	H	H	H
172	OMe	OMe	O		H	H	H	H	H	H	H











No.	R ²	R ³	X	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸³	R ⁸⁴	R ⁸⁵	R ⁸⁶	R ⁸⁷
173	OMe	MPO	O	CH ₂ CO ₂ H	H	H	H	H	H	H	H
174	OMe	OMe	O	NHC(O)Me	H	H	H	H	H	H	H
175	OMe	MPO	O	OCH ₂ CONH(CH ₂) ₂ -OH	H	H	H	H	H	H	H
176	OMe		O	OCH ₂ CONH(CH ₂) ₂ -OH	H	H	H	H	H	H	H
177	OMe	DMMPPO	O	OCH ₂ CONH(CH ₂) ₂ -OH	H	H	H	H	H	H	H
178	OMe	MPO	O	OCH ₂ CH ₂ NHS(O) ₂ -Me	H	H	H	H	H	H	H
179	OMe	MPO	O	O(CH ₂) ₂ N(Me)CO N(CH ₂ Me) ₂	H	H	H	H	H	H	H
180	OMe	MPO	O	O(CH ₂) ₃ NHCOMe	H	H	H	H	H	H	H
181	OMe	MPO	O	O(CH ₂) ₃ NHCOCH- (Me) ₂	H	H	H	H	H	H	H
182	OMe	MPO	O		H	H	H	H	H	H	H
183	OMe	MPO	O		H	H	H	H	H	H	H
184	OMe	MPO	O	O(CH ₂) ₂ NHCOCH- (Me) ₂	H	H	H	H	H	H	H
185	OMe	OMe	O		H	H	H	H	H	H	H
186	OMe	OMe	O	OCH ₂ CH ₂ NHSO ₂ Me	H	H	H	H	H	H	H

No.	R ²	R ³	X	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸³	R ⁸⁴	R ⁸⁵	R ⁸⁶	R ⁸⁷
187	OMe	OMe	O	H		H	H	H	H	H	H
188	OMe	OMe	O		H	H	H	H	H	H	H
189	OMe	OMe	O		H	H	H	H	H	H	H
190	OMe	OMe	O		H	H	H	H	H	H	H
191	OMe	OMe	O	O(CH ₂) ₃ NHS(O) ₂ Me	H	H	H	H	H	H	H
192	OMe	OMe	O	O(CH ₂) ₃ NHCOCH-(Me) ₂	H	H	H	H	H	H	H
193	OMe	MPO	O	O(CH ₂) ₃ NHS(O) ₂ Me	H	H	H	H	H	H	H
194	OMe		O	OCH ₂ CONH(CH ₂) ₂ -OH	H	H	H	H	H	H	H
195	OMe	OMe	O		H	H	H	H	H	H	H
196	OMe	OMe	O	H	Me	H	H	H	H	H	H
197	OMe		O	OMe	H	H	H	H	H	H	H













No.	R ²	R ³	X	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸³	R ⁸⁴	R ⁸⁵	R ⁸⁶	R ⁸⁷
198	OMe	OMe	O	NHMe	H	H	H	H	H	H	H
199	OMe	OMe	O	NHCH ₂ Me	H	H	H	H	H	H	H
400	OMe	OMe	O	N(SO ₂ Me) ₂	H	H	H	H	H	H	H
401	OMe	OMe	O	OCH ₂ C(O)NHCH ₂ -C(O)NH ₂	H	H	H	H	H	H	H
402	OMe	OMe	O	OCH ₂ C(O)NHCH-(Me) C(O)NHMe	H	H	H	H	H	H	H
403	OMe	OMe	O	OCH ₂ C(O)NHCH ₂ -C(O)NHMe	H	H	H	H	H	H	H
404	OMe	OMe	O	OCH ₂ C(O)N(CH ₂ Me)C(O)NH(CH ₂) ₃ N(Me) ₂	H	H	H	H	H	H	H
405	OMe	OMe	O		H	H	H	H	H	H	H
406	OMe	OMe	O		H	H	H	H	H	H	H
407	OMe	OMe	O	O(CH ₂) ₂ N(Me)C(O)N(CH ₂ Me) ₂	H	H	H	H	H	H	H
408	OMe	MPO	O	O(CH ₂) ₂ NHCOCH-(Me) ₂	H	H	H	H	H	H	H
409	OMe	OMe	O	O(CH ₂) ₂ NHC(O)Me	H	H	H	H	H	H	H
410	OMe	OMe	O	(CH ₂) ₂ C(O)NHMe	H	H	H	H	H	H	H
411	OMe	OMe	O	(CH ₂) ₂ C(O)NHS(O) ₂ Me	H	H	H	H	H	H	H
412	OMe	OMe	O		H	H	H	H	H	H	H
413	OMe	OMe	O	(CH ₂) ₂ C(O)NHCH ₂ -CHCH ₂	H	H	H	H	H	H	H











No.	R ²	R ³	X	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸³	R ⁸⁴	R ⁸⁵	R ⁸⁶	R ⁸⁷
414	OMe	OMe	O		H	H	H	H	H	H	H
415	OMe	OMe	O		H	H	H	H	H	H	H
416		OMe	O	CN	H	H	H	H	H	H	H
417	OMe	OMe	O		H	H	H	H	H	H	H
418	OMe		O	H	NHCH ₂ Me	H	H	H	H	H	H
419	OMe		O	F	H	H	H	H	H	H	H
420	OMe		O	CN	H	H	H	H	H	H	H
421	OMe		O	OMe	H	H	H	H	H	H	H
422	OMe		O	H	OMe	H	H	H	H	H	H
423	OH	OMe	O	H	OCF ₂ CF ₂ H	H	H	H	H	H	H
424	OCH ₂ C ₆ H ₅	OMe	O	OMe	H	H	H	H	H	H	H
425	COOMe	OMe	O	OMe	H	H	H	H	H	H	H
426	OH	OMe	O	OCH ₂ CH=CH ₂	H	H	H	H	H	H	H
427	OMe	OH	O	OCH ₂ CONHMe	H	H	H	H	H	H	H

No.	R ²	R ³	X	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸³	R ⁸⁴	R ⁸⁵	R ⁸⁶	R ⁸⁷
428	H	OMe	O	H		H	H	H	H	H	H
429	H	OMe	O	OMe	H	H	H	H	H	H	H
430	OMe	OMe	O		H	H	H	H	H	H	H
431	OMe		O	H	OMe	H	H	H	H	H	H
432	OMe	OMe	O		H	H	H	H	H	H	H
433	OMe	OMe	O		H	H	H	H	H	H	H
434	OMe	OMe	O		H	H	H	H	H	H	H
435	OMe	OMe	O		H	H	H	H	H	H	H

No.	R ²	R ³	X	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸³	R ⁸⁴	R ⁸⁵	R ⁸⁶	R ⁸⁷
436	OMe	OMe	O		H	H	H	H	H	H	H
437	OMe	OMe	O		H	H	H	H	H	H	H
438	OMe	OMe	O	H		H	H	H	H	H	H
439	OMe	OMe	O	H		H	H	H	H	H	H
440	OMe	OMe	O	H		H	H	H	H	H	H
441	OMe	OMe	O		H	H	H	H	H	H	H
442	OMe	OMe	O		H	H	H	H	H	H	H
443	OMe	OMe	O		H	H	H	H	H	H	H
444	OMe	OMe	O		H	H	H	H	H	H	H
445	OMe	OMe	O		H	H	H	H	H	H	H

No.	R ²	R ³	X	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸³	R ⁸⁴	R ⁸⁵	R ⁸⁶	R ⁸⁷
446	OMe	OMe	O		H	H	H	H	H	H	H
447	OMe	OMe	O		H	H	H	H	H	H	H
448	OMe	OMe	O		H	H	H	H	H	H	H
449	OMe	DMMP	O	H		H	H	H	H	H	H
450	OMe	DMMP	O	H		H	H	H	H	H	H
451	OMe		O	H		H	H	H	H	H	H
452	OMe		O	H		H	H	H	H	H	H
453	OMe		O	H		H	H	H	H	H	H
454	OMe	DMMP	O	H		H	H	H	H	H	H
455	OMe	DMMP	O	H		H	H	H	H	H	H

No.	R ²	R ³	X	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸³	R ⁸⁴	R ⁸⁵	R ⁸⁶	R ⁸⁷
456	OMe		O	H		H	H	H	H	H	H
457	OMe	DMMPO	O	H	OMe	H	H	H	H	H	H
458	OMe	MPO	O	H		H	H	H	H	H	H
459	OMe	DMMPO	O	H		H	H	H	H	H	H
460	OMe		O	H		H	H	H	H	H	H
461	OMe	MPO	O	H		H	H	H	H	H	H
462	OMe	DMMPO	O	H		H	H	H	H	H	H
463	OMe		O	H		H	H	H	H	H	H
464	OMe	MPO	O	H		H	H	H	H	H	H
465	OMe	DMMPO	O	H		H	H	H	H	H	H

No.	R ²	R ³	X	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸³	R ⁸⁴	R ⁸⁵	R ⁸⁶	R ⁸⁷
466	OMe	DMMP	O		H	H	H	H	H	H	H
467	OMe	MPO	O		H	H	H	H	H	H	H
468	OMe		O		H	H	H	H	H	H	H
469	OMe	DMMP	O		H	H	H	H	H	H	H
470	OMe		O		H	H	H	H	H	H	H
471	OMe	MPO	O		H	H	H	H	H	H	H
472	OMe	OMe	O		H	H	H	H	H	H	H
473	OMe	OMe	O		H	H	H	H	H	H	H




No.	R ²	R ³	X	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸³	R ⁸⁴	R ⁸⁵	R ⁸⁶	R ⁸⁷
474	OMe	OMe	O		H	H	H	H	H	H	H
475	OMe	OMe	O	O(CH ₂) ₂ NHCOO CH ₂ CH=CH ₂	H	H	H	H	H	H	H
476	OMe	OMe	O		H	H	H	H	H	H	H
477	OMe	OMe	O	H	OCH ₂ - C(O)NH- Me	H	H	H	H	H	H
478	OMe	MEO	O	F	H	H	H	H	H	H	H
479	OMe	MEO	O	CN	H	H	H	H	H	H	H
480	OMe	MEO	O	H	NHCH ₂ - Me	H	H	H	H	H	H
481	OMe	DMMPPO	O	H	OCH ₃ C(O)NH- Me	H	H	H	H	H	H
482	OMe	OMe	O	H	OCH ₂ C(O)NH- CH(Me) ₂	H	H	H	H	H	H

Table 2



No.	R ²	R ³	Y'	Y''	Z	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸⁶
200	OMe	OMe	CH	CH	N	H	H	OMe	H
201	OMe	OMe	CH	CH	N	H	OMe	H	H
202	OMe	OMe	CH	CH	N	OMe	H	H	H
203	OMe	OH	N	CH	N	OMe	H	H	H
204	O(CH ₂) ₃ -N(CH ₃)	OMe	CH	CH	N	OMe	H	H	H
205	OMe	O(CH ₂) ₃ -N(CH ₃)	CH	CH	N	OMe	H	H	H
206	O(CH ₂) ₃ N(Me) ₂	OMe	CH	CH	N	OMe	H	H	H
207	OMe	O(CH ₂) ₃ N(Me) ₂	CH	CH	N	OMe	H	H	H
208	MPO	OMe	CH	CH	N	OMe	H	H	H
209	OMe	MPO	CH	CH	N	OMe	H	H	Me
210	OMe	MPO	N	CH	N	OMe	H	H	H
211	OMe	OH	N	CH	CH	OMe	H	H	Me
212	OMe	OMe	CH	CH	N	H	CF ₃	H	H


























No.	R ²	R ³	Y'	Y''	Z	R ⁸⁰	R ⁸¹	R ⁸²	R ⁸⁶
213	OMe	MPO	N	CH	CH	F	H	H	H
214	OMe	MEO	N	CH	CH	H	OMe	H	H
215	OMe	MPO	CH	N	CH	OMe	H	H	H
216	OMe	MPO	CH	N	CH	OCH ₂ CONHMe	H	H	H
217	OMe	MPO	N	CH	CH	H	OMe	H	H
218	OMe	OH	N	CH	CH	F	H	H	Me
219	OMe	OCH ₂ C ₆ H ₅	N	CH	CH	F	H	H	Me
220	OH	OMe	N	CH	CH	OMe	H	H	H
221	OMe	OH	N	CH	CH	OMe	H	H	H
222	OMe	OCH ₂ C ₆ H ₅	N	CH	CH	OMe	H	H	Me
223	OMe	DMMP	N	CH	CH	H		H	H
224	OMe	 O(CH ₂) ₃ -N	N	CH	CH	H		H	H
225	OMe	DMMP	N	CH	CH	H		H	H










Table 3













NO.	R ²	R ³	R ⁶	R ⁷	X
250	OMe	OMe	p-Ph		O
251	OMe	OMe	p-Ph		O
252	OMe	OMe	p-Ph		O
253	OMe	OMe	p-Ph		O
254	OMe	OMe	p-Ph		O
255	OMe	OMe	p-Ph		O
256	OMe	OMe	p-Ph		O
257	OMe	OMe	p-Ph		O
258	OMe	OMe	p-Ph		O
259	OMe	OMe	p-Ph		O
260	OMe	DMMPO	p-Ph	2-thiazole	O
261	OMe	OMe	p-Ph		O


NO.	R ²	R ³	R ⁶	R ⁷	X
262	OMe	OMe	p-Ph		O
263	OMe	OMe	p-Ph		O
264	OMe	OMe	p-Ph		O
265	OMe	OMe	p-Ph		O
266	OMe	OMe			O
267	OMe	OMe	p-Ph		S
268	OMe	OMe	p-Ph	2-thiazole	O
269	OMe	OMe	p-Ph		O
270	OMe	OMe	p-Ph		O
271	OMe	OMe	p-Ph		O
272	OCH ₂ C ₆ H ₅	OMe	p-Ph	2-thiazole	O
273	OH	OMe	p-Ph	2-thiazole	O
274	MPO	OMe	p-Ph	2-thiazole	O
275		OMe	p-Ph	2-thiazole	O
276		OMe	p-Ph	2-thiazole	O
277	MPO	OMe	p-Ph	2-thiazole	O
278	MEO	OMe	p-Ph	2-thiazole	O

NO.	R ²	R ³	R ⁶	R ⁷	X
279		OMe	p-Ph	2-thiazole	O
280		OMe	p-Ph	2-thiazole	O
281	O(CH ₂) ₂ N(Me) ₂	OMe	p-Ph	2-thiazole	O
282	OMe	OH	p-Ph	2-thiazole	O
283	OMe	MPO	p-Ph	2-thiazole	O
284	OMe		p-Ph	2-thiazole	O
285	OMe		p-Ph	2-thiazole	O
286	OMe	O(CH ₂) ₃ N(Me) ₂	p-Ph	2-thiazole	O
287	OMe	OMe			O
288	OMe	OCH ₂ COOCH ₂ Me	p-Ph	2-thiazole	O
289	OMe	OCH ₂ COOH	p-Ph	2-thiazole	O
290	O(CH ₂) ₂ OMe	O(CH ₂) ₂ OMe	p-Ph	2-thiazole	O
291	OMe	OCH ₂ CONHMe	p-Ph	2-thiazole	O
292	OMe	OCH ₂ CONHCH ₂ CHCH ₂	p-Ph	2-thiazole	O
293	NH ₂	OMe	p-Ph	2-thiazole	O
294	OMe	MPO	p-Ph	2-pyridyl	O
295	OMe	OMe	p-Ph	2-thiazole	S
296	OMe	OMe	p-Ph		S
297	OMe	OMe	p-Ph	cyclopentyl	O
298	OMe	OMe	p-Ph	cyclohexyl	O
299	OMe	OMe	p-Ph		O
300	OMe	OCH ₂ C ₆ H ₅	p-Ph	2-thiazole	O
301	NHCO ₂ C (Me) ₃	OMe	p-Ph	2-thiazole	O


NO.	R ²	R ³	R ⁶	R ⁷	X
302	OMe		p-Ph	2-thiazole	O
303	OMe	OMe	p-Ph		O
304	OMe	OMe	p-Ph		CH ₂
305	OMe	OMe	p-Ph		CH ₂
306	OMe	OMe	p-Ph		O
307	OMe	OMe	p-Ph		O
308	OMe	OMe	p-Ph		O
309	OMe	OMe	p-Ph		S
310	OMe	MEO	p-Ph		O

NO.	R ²	R ³	R ⁶	R ⁷	X
311	OMe	OMe	p-Ph		O
312	OMe	OMe	p-Ph		O
313	OMe	OMe	p-Ph		O
314	OMe	OMe	p-Ph		O
315	OMe	OMe	p-Ph		O
316	OMe	OMe	p-Ph		O
317	OMe	OMe	p-Ph		O
318	OMe		p-Ph	2-thiazole	O

NO.	R ²	R ³	R ⁶	R ⁷	X
319	OMe		p-Ph	2-thiazole	O
320	OMe		p-Ph	2-thiazole	O


where p-Ph represents a para-phenylene group of formula 

5 Compounds of formula (I) are suitably prepared by reacting a compound of formula (III)



(III)

10 where R^{1'}, R^{2'}, R^{3'}, R^{4'} represent R¹, R², R³ and R⁴ respectively as defined in relation to formula (I) or a precursor thereof, and Z' is a leaving group, with a compound of formula (IV)



(IV)


15 where R⁶, Y, X, and n are as defined in relation to formula (I), and R^{7'} is a group R⁷ or a precursor thereof; and thereafter if necessary or desired converting precursor groups R^{1'}, R^{2'}, R^{3'}, R^{4'} and R^{7'} to groups of formula R¹, R², R³, R⁴ and R⁷ respectively, or converting a group R¹, R², R³, R⁴ and R⁷ to a different such group.

Suitable leaving groups for Z' include halogen such as bromo or chloro, or a mesylate or tosylate group or a substituted phenoxy group.

The reaction is suitably carried out in an organic solvent such as an alcohol for example propanol or cyclohexanol at elevated temperatures, for example of from 50 to 150°C, for example at about 105°C.

Conversion reactions in which precursor groups R^{1'}, R^{2'}, R^{3'}, R^{4'} are converted to groups of formula R¹, R², R³ and R⁴ respectively, or groups R¹, R², R³ and R⁴ are converted to different such group can be carried out using conventional chemistry as outlined hereinafter. Particular precursor groups R^{1'}, R^{2'}, R^{3'}, R^{4'} are groups of formula R^{13'}-X¹-(CH₂)_x wherein x and X¹ are as defined hereinafter, and R^{13'} is C₁₋₅alkyl which is substituted with halo other than fluoro, and in particular chloro or bromo. The chloro or bromo group may readily be converted into many other groups R¹³ as defined in relation to claim 1. Such compounds are novel and form a further aspect of the invention. They may have activity similar to that of compounds of formula (I) in their own right and therefore may be used in place of a compound of formula (I).

Thus the invention further provides a compound of formula (IB)




(IB)

where Y, n, R⁶, X and R⁷ are as defined in claim 1 and at least one of R^{1''}, R^{2''}, R^{3''} or R^{4''} is a group R^{13'}-X¹-(CH₂)_x wherein X¹ and x are as defined in claim 1 and R^{13'} is alkyl substituted by chloro or bromo; and the remainder are groups R¹, R², R³ and R⁴ respectively.

Similarly conversion reactions involving groups R⁷ may be effected using conventional chemistry. For example substituent groups on a group R⁹ within the group R⁷ may be changed, for example by changing acids to esters or amides etc.

Alternatively, compounds of formula (I) are prepared by reacting a compound of formula (V)



(V)

- 5 where R^1 , R^2 , R^3 , R^4 are as defined in relation to formula (III) R^6 , X, Y and n are as defined in relation to formula (I), with a compound of formula (VI)




where R^7 is as defined in relation to formula (IV) and Z'' is a leaving group;

- 10 and thereafter if necessary or desired converting precursor groups R^1 , R^2 , R^3 , R^4 and R^7 to groups of formula R^1 , R^2 , R^3 , R^4 and R^7 respectively, or converting a group R^1 , R^2 , R^3 , R^4 and R^7 to a different such group. Suitable leaving groups for Z'' include halogen such a bromo or chloro, or a mesylate or tosylate group. Conversion reactions are as described above.

- 15 The reaction is suitably carried out in an organic solvent such as DMF at elevated temperatures, for example of from 40 to 120°C, for example at about 80°C.

- Compounds of formula (III) and (V) are either known compounds or they can be prepared from known compounds by conventional methods, for example as described in WO 98/43960, WO 98/13350. Exemplary preparations of compounds of formula (III) are included hereinafter.

- Compounds of formula (IV) are also known compounds (see for example Rev. Chim. (Bucharest) (1988), 39(6), 477-82 and DD 110651: 74.01.05) or they can be prepared from known compounds using conventional methods. For example, where Y is NH, compounds of formula (IV) are suitably prepared by reduction of a compound of formula (VII)
- 25



(VII)

where X, R⁶, R⁷ and n are as defined above. It may be convenient to convert precursor groups R⁷ to groups R⁷ or groups R⁷ to other such groups at the level of compound of
5 formula (VII) or (IV) using conventional chemistry.

Compounds of formula (VI) are also known compounds or they can be prepared from known compounds by conventional methods.

Compounds of the invention are useful in the inhibition of MEK enzyme activity and can be used in the treatment of proliferative disease. They will suitably be in the form
10 of a pharmaceutical composition, in combination with a pharmaceutically acceptable carrier. Such compositions form a further aspect of the invention.

The compositions of the invention may be in a form suitable for oral use (for example as tablets, lozenges, hard or soft capsules, aqueous or oily suspensions, emulsions, dispersible powders or granules, syrups or elixirs), for topical use (for example
15 as creams, ointments, gels, or aqueous or oily solutions or suspensions), for administration by inhalation (for example as a finely divided powder or a liquid aerosol), for administration by insufflation (for example as a finely divided powder) or for parenteral administration (for example as a sterile aqueous or oily solution for intravenous, subcutaneous, intramuscular or intramuscular dosing or as a suppository for
20 rectal dosing).

The compositions of the invention may be obtained by conventional procedures using conventional pharmaceutical excipients, well known in the art. Thus, compositions intended for oral use may contain, for example, one or more colouring, sweetening, flavouring and/or preservative agents.

25 Suitable pharmaceutically acceptable excipients for a tablet formulation include, for example, inert diluents such as lactose, sodium carbonate, calcium phosphate or calcium carbonate, granulating and disintegrating agents such as corn starch or algenic acid; binding agents such as starch; lubricating agents such as magnesium stearate, stearic acid or talc; preservative agents such as ethyl or propyl p-hydroxybenzoate, and
30 anti-oxidants, such as ascorbic acid. Tablet formulations may be uncoated or coated either to modify their disintegration and the subsequent absorption of the active ingredient

within the gastrointestinal tract, or to improve their stability and/or appearance, in either case, using conventional coating agents and procedures well known in the art.

Compositions for oral use may be in the form of hard gelatin capsules in which the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules in which the active ingredient is mixed with water or an oil such as peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions generally contain the active ingredient in finely powdered form together with one or more suspending agents, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents such as lecithin or condensation products of an alkylene oxide with fatty acids (for example polyoxyethylene stearate), or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives (such as ethyl or propyl p-hydroxybenzoate, anti-oxidants (such as ascorbic acid), colouring agents, flavouring agents, and/or sweetening agents (such as sucrose, saccharine or aspartame).

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil (such as arachis oil, olive oil, sesame oil or coconut oil) or in a mineral oil (such as liquid paraffin). The oily suspensions may also contain a thickening agent such as beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set out above, and flavouring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water generally contain the active ingredient together with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable

dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients such as sweetening, flavouring and colouring agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or arachis oil, or a mineral oil, such as for example liquid paraffin or a mixture of any of these. Suitable emulsifying agents may be, for example, naturally-occurring gums such as gum acacia or gum tragacanth, naturally-occurring phosphatides such as soya bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides (for example sorbitan monooleate) and condensation products of the said partial esters with ethylene oxide such as polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening, flavouring and preservative agents.

Syrups and elixirs may be formulated with sweetening agents such as glycerol, propylene glycol, sorbitol, aspartame or sucrose, and may also contain a demulcent, preservative, flavouring and/or colouring agent.

The pharmaceutical compositions may also be in the form of a sterile injectable aqueous or oily suspension, which may be formulated according to known procedures using one or more of the appropriate dispersing or wetting agents and suspending agents, which have been mentioned above. A sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example a solution in 1,3-butanediol.

Suppository formulations may be prepared by mixing the active ingredient with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Suitable excipients include, for example, cocoa butter and polyethylene glycols.

Topical formulations, such as creams, ointments, gels and aqueous or oily solutions or suspensions, may generally be obtained by formulating an active ingredient with a conventional, topically acceptable, vehicle or diluent using conventional procedure well known in the art.

Compositions for administration by insufflation may be in the form of a finely divided powder containing particles of average diameter of, for example, 30 μ or much less, the powder itself comprising either active ingredient alone or diluted with one or

more physiologically acceptable carriers such as lactose. The powder for insufflation is then conveniently retained in a capsule containing, for example, 1 to 50mg of active ingredient for use with a turbo-inhaler device, such as is used for insufflation of the known agent sodium cromoglycate.

5 Compositions for administration by inhalation may be in the form of a conventional pressurised aerosol arranged to dispense the active ingredient either as an aerosol containing finely divided solid or liquid droplets. Conventional aerosol propellants such as volatile fluorinated hydrocarbons or hydrocarbons may be used and the aerosol device is conveniently arranged to dispense a metered quantity of active ingredient.

10 For further information on Formulation the reader is referred to Chapter 25.2 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

 The amount of active ingredient that is combined with one or more excipients to produce a single dosage form will necessarily vary depending upon the host treated and
15 the particular route of administration. For example, a formulation intended for oral administration to humans will generally contain, for example, from 0.5 mg to 2 g of active agent compounded with an appropriate and convenient amount of excipients which may vary from about 5 to about 98 percent by weight of the total composition. Dosage unit forms will generally contain about 1 mg to about 500 mg of an active ingredient. For
20 further information on Routes of Administration and Dosage Regimes the reader is referred to Chapter 25.3 in Volume 5 of Comprehensive Medicinal Chemistry (Corwin Hansch; Chairman of Editorial Board), Pergamon Press 1990.

 The size of the dose for therapeutic or prophylactic purposes of a compound of the Formula I will naturally vary according to the nature and severity of the conditions,
25 the age and sex of the animal or patient and the route of administration, according to well known principles of medicine. As mentioned above, compounds of the Formula I are useful in treating diseases or medical conditions which are due alone or in part to the effects MEK enzymes.

 In using a compound of the Formula I for therapeutic or prophylactic purposes it
30 will generally be administered so that a daily dose in the range, for example, 0.5 mg to 75 mg per kg body weight is received, given if required in divided doses. In general lower doses will be administered when a parenteral route is employed. Thus, for example, for

intravenous administration, a dose in the range, for example, 0.5 mg to 30 mg per kg body weight will generally be used. Similarly, for administration by inhalation, a dose in the range, for example, 0.5 mg to 25 mg per kg body weight will be used. Oral administration is however preferred.

- 5 In a further aspect, the invention provides a method of treating proliferative disease by administering a compound of formula (I) as described above, or a pharmaceutical composition as described above.


Yet a further aspect of the invention provides the use of a compound of formula (I) as defined above, in the preparation of a medicament for use in the inhibition of MEK
10 enzyme activity and in particular for the treatment of proliferative disease such as cancer.

The invention will now be particularly described by way of Example. The preparation of various intermediates used in the Examples is described in the Preparations.

Preparation 1

Chloroquinoline intermediates

- 15 These can be prepared for example using the following scheme where "Bz" represents benzyl.



- A mixture of **(1)** (10.36g., 45.3 mmole) and diethylethoxymethylene malonate (9mL, 45.3
20 mmole) was heated at 110°C for 1 hour and then allowed to cool overnight. The mixture was evaporated and the product **(2)** used in the next step without further purification.

Mass Spectrum m/e 400 ($M^+ + H$).

Preparation of (3)

- 5 A mixture of (2) (assumed 45.3 mmole) and phosphoryl chloride (83.3mL, 906 mmole) was heated at 115 ° C for 18 hours. After cooling, the solution was evaporated to remove excess phosphoryl chloride. The residue was treated with ice and aqueous ammonia to hydrolyse the remaining phosphoryl chloride. The solid product was filtered off and dried in a vacuum oven to give a cream coloured solid, 9.0g (53% yield).
- 10 Mass Spectrum m/e 372 ($M^+ + H$).

Preparation of (4)

- A mixture of (3) (9.0g, 24.2 mmole) was stirred in ethanol (48.3mL) for 15 minutes at ambient temperature to give a smooth suspension. Aqueous sodium hydroxide solution
- 15 (2.0M, 48.3mL, 96.7 mmole) was added and the mixture stirred for 18 hours at ambient temperature. The ethanol was removed by rotary evaporation and the resulting solution was acidified to pH 2 with hydrochloric acid while stirring. The precipitate was filtered off and dried in a vacuum oven to give an orange solid, 7.19g (86% yield).

Mass Spectrum m/e 344 ($M^+ + H$).

20

Preparation of (5)

- A mixture of (4) (7.18g, 20.9 mmole) and thionyl chloride (90 mL) was refluxed for 2 hours. After cooling the excess thionyl chloride was removed by rotary evaporation and the residue was suspended in acetone (175mL) and the resulting suspension cooled in an
- 25 ice-bath. Aqueous ammonia (S.G. 0.880, 20mL) was added gradually, keeping the temperature below 10 ° C. The resulting suspension was filtered off, washed with water and air-dried to give a solid, 5.15g (75% yield).

Mass Spectrum m/e 343 ($M^+ + H$).

30 Preparation of (6)

A mixture of (5) (20.55g, 60 mmole) and phosphoryl chloride (250mL) was heated and stirred at 120 ° C for 4 hours when the starting material had dissolved. Heating and stirring

was continued at 110 ° C for 18 hours. After cooling, the solution was evaporated to remove excess phosphoryl chloride. Last traces of phosphoryl chloride were removed by azeotrope with toluene. The residue was treated with ice and aqueous ammonia to remove acidity. The solid product was filtered off and dried in a vacuum oven to give a grey solid, 19.23g (99% yield).

(This may also be prepared as described in WO 9843960)

Mass Spectrum m/e 325 ($M^+ + H$).

Preparation of (7)

- 10 A mixture of (6) (19.23g, 60.0 mmole) and trifluoroacetic acid (300 mL) and thioanisole (35mL) was refluxed in a nitrogen atmosphere for 3 hours. After cooling the trifluoroacetic acid was removed by rotary evaporation and the oily residue was stirred with ice and water and basified with aqueous ammonia (S.G. 0.880). The resulting suspension was filtered and the solid was washed successively with water, ethyl acetate
- 15 and diethyl ether and then dried to give a khaki solid, 13.74g (97% yield).

Mass Spectrum m/e 235 ($M^+ + H$).

Preparation of (8)

(4-chloro-6-methoxy-7-[3-(1-morpholino)propoxy]-3-quinolinecarbonitrile)

- 20 A mixture of (7) (2.34g, 10.0 mmole) and 1-(3-chloropropyl)morpholine (2.45g, 15.0 mmole) and anhydrous potassium carbonate (2.07g, 15.0 mmole) suspended in butanone (150mL) was stirred in a oil-bath at 88 ° C for 96 hours. The suspension was filtered hot to remove inorganics and the filtrate was allowed to cool and then evaporated to ca. 100mL. A solid precipitated on standing for 72 hours. The solid was filtered off and
- 25 washed with a little acetone and then dried to give a white solid, 0.54g (15% yield).

Mass Spectrum m/e 362 ($M^+ + H$).

Preparation 2

By similar processes the following analogues were also prepared:-





Table 4

R ¹	R ²	Mass Spectrum
OCH ₂ CH ₂ OMe	OCH ₂ CH ₂ OMe	m/e 337 (M ⁺ +H).
OMe	MPE	m/e 348 (M ⁺ +H)
OMe		m/e 332 (M ⁺ +H).
OCH ₂ C ₆ H ₅	OMe	m/e 324 (M ⁺ +H).
OH	OMe	m/e 234 (M ⁺ +H).
OCH ₂ C(O) ₂ CH ₂ Me	OMe	m/e 321 (M ⁺ +H).
OMe	OCH ₂ C(O) ₂ CH ₂ Me	m/e 321 (M ⁺ +H).
OCH ₂ C(O) ₂ Me	OMe	
OMe	O(CH ₂) ₃ Cl	m/e 310 (M ⁺ +H).

Example 1

A mixture of 4-chloro-3-cyano-6,7-dimethoxyquinoline (1.5 g), prepared as described in WO 9843960, and 4-(2-methoxyphenoxy)-aniline (2.58 g), prepared as described in Rev. Chim. (Bucharest) (1988), 39(6),477-82, in 1-propanol (90 ml) was stirred and heated at 105°C for 6 hours. The mixture was cooled to ambient temperature and then filtered.

The crystals were washed with a small volume of 1-propanol and then dried to give 4-(2-methoxyphenoxy)-anilino-3-cyano-6,7-dimethoxyquinoline (Compound 1 in Table 1)

(2.19 g, 85%).

Mass Spectrum m/e 428 (M⁺+H).

NMR Spectrum (d-6-DMSO, d values) 3.75 (s, 3H), 4.00 (s, 6H), 6.95 (m, 3H), 7.05 (m, 1H), 7.20 (m, 2H), 7.40 (d, 2H), 7.50 (s, 1H), 8.20 (s, 1H), 8.85 (s, 1H), 11.10 (broad, 1H).

Example 2Preparation of Compound 253 in Table 3Step 1

A mixture of 4-chloro-3-cyano-6,7-dimethoxy-quinoline (2.49 g) and 4-aminophenol (2.4 g) in n-propanol (150 ml) was stirred and heated at 110°C for 4 hours. The mixture was

cooled to ambient temperature and then filtered. The crystals were washed with a small volume of diethyl ether and then dried to give 3-cyano-6,7-dimethoxy-4-(4-hydroxy)-anilino-quinoline (2.68 g, 83%).

Mass Spectrum m/e 322 ($M^+ + H$).

- 5 NMR Spectrum (d-6-DMSO, d values) 3.85 (s, 3H), 3.9 (s, 3H), 6.8 (d, 2H), 7.1 (d, 2H), 7.25 (s, 1H), 7.8 (s, 1H), 8.3 (s, 1H), 9.3 (broad s, 1H).

Step 2

- 3-Cyano-6,7-dimethoxy-4-(4-hydroxy)-anilino-quinoline (160.5 mg) was dissolved in DMF (5 ml) and potassium carbonate (138 mg) was added. The mixture was stirred
10 under an atmosphere of nitrogen for 5 minutes and then 2-bromomethyl-tetrahydrofuran (180 ml) was added. The mixture was stirred and heated at 80°C for 18 hours. The mixture was cooled to ambient temperature and then diluted with ethyl acetate and then extracted with water. The aqueous phase was re-extracted with ethyl acetate and the combined organic extracts were washed with brine, dried (Na_2SO_4) and evaporated. The
15 residue was then purified by column chromatography using 2-3% methanol/dichloromethane mixtures as eluent. There was thus obtained 3-cyano-6,7-dimethoxy-4-(2-tetrahydrofuranyl-methoxy)-anilino-quinoline (70 mg, 34%).

Mass Spectrum m/e 406 ($M^+ + H$).

- 20 NMR Spectrum ($CDCl_3$, d values) 1.8 (m, 1H), 1.95 (m, 2H), 2.05 (m, 1H), 3.6 (s, 3H), 3.85 (dd, 1H), 3.9 (m, 1H), 3.95 (m, 1H), 4.0 (s, 3H), 4.25 (m, 1H), 6.8 (broad s, 1H), 6.85 (s, 1H), 6.95 (d, 2H), 7.1 (d, 2H), 7.35 (s, 1H), 8.6 (s, 1H).

Example 3

By an analogous procedure to that described for Example 2, step 2, but using an alternative bromide, the compounds listed in Table 5 were prepared:

Table 5

No	bromide	mass spec	nmr	Notes
250	2-bromo-methyltetrahydropyran	m/e 420 (M ⁺ +H)	(d-6-DMSO, d values) 1.2-1.7 (m, 6H), 3.40 (m, 1H), 3.60 (m, 1H), 3.90 (s, 3H), 3.90 (s, 3H), 3.9 (m, 3H), 6.95 (d, 2H), 7.20 (d, 2H), 7.25 (d, 1H), 7.75 (d, 1H), 8.30 (d, 1H), 9.35 (broad s, 1H).	
251	epibromohydrin	m/e 378 (M ⁺ +H)	(d-6-DMSO, d values) 2.70 (dd, 1H), 2.83 (dd, 1H), 3.35 (m, 1H), 3.85 (dd, 1H), 3.90 (s, 3H), 3.95 (s, 3H), 4.35 (dd, 1H), 7.00 (d, 2H), 7.20 (d, 2H), 7.26 (s, 1H), 7.75 (s, 1H), 8.30 (s, 1H), 9.35 (broad s, 1H).	RT/ 48hrs/ DMF/ K ₂ CO ₃
252	2-bromomethyl-1,3-dioxolane	m/e 408 (M ⁺ +H)	(CDCl ₃ , d values) 3.60 (s, 3H), 3.95 (m, 3H), 4.00 (s, 3H), 4.05 (m, 3H), 5.30 (t, 1H), 6.80 (broad s, 1H), 6.85 (s, 1H), 6.95 (d, 2H), 7.15 (d, 2H), 7.35 (s, 1H), 8.60 (s, 1H).	

Example 4


By an analogous procedure to that described for Example 2, step 2, but using a tosylate instead of a bromide, the following compounds were prepared.

Table 6

No	intermediate	mass	nmr
254	2,2-dimethyl-4-(4-toluenesulphonyloxymethyl)-1,3-dioxolane	m/e 436 (M ⁺ +H)	(CDCl ₃ , d values) 1.4 (s, 3H), 1.45 (s, 3H), 3.65 (s, 3H), 3.90 (dd, 1H), 3.95 (m, 1H), 4.00 (s, 3H), 4.05 (m, 1H), 4.15 (dd, 1H), 4.50 (m, 1H), 6.80 (broad s, 1H), 6.90 (s, 1H), 6.95 (d, 2H), 7.10 (d, 2H), 7.35 (s, 1H), 8.60 (s, 1H).
255	4-(4-toluenesulphonyloxymethyl)-1,3-dioxolane	m/e 408 (M ⁺ +H)	(CDCl ₃ , d values) 3.60 (s, 3H), 3.85 (m, 1H), 3.95 (m, 1H), 4.00 (s, 3H), 4.05 (m, 2H), 4.40 (m, 1H), 4.95 (s, 1H), 5.10 (s, 1H), 6.80 (broad s, 1H), 6.85 (s, 1H), 6.95 (d, 2H), 7.10 (d, 2H), 7.35 (s, 1H), 8.60 (s, 1H).
256	5-bromo-5-(4-toluenesulphonyloxymethyl)-1,3-dioxane	m/e 436 (M ⁺ +H)	(CDCl ₃ , d values) 0.95 (s, 3H), 3.50 (d, 2H), 3.65 (s, 3H), 4.00 (d, 2H), 4.00 (s, 3H), 4.10 (s, 1H), 4.70 (d, 1H), 5.00 (d, 1H), 6.80 (broad s, 1H), 6.85 (s, 1H), 6.95 (d, 2H), 7.15 (d, 2H), 7.35 (s, 1H), 8.60 (s, 1H).

Example 5

Using a method analogous to that described in Example 1 (except that in some instances, intermediates (1) and (2) were modified prior to further reaction as described in Examples 14 and 15 hereinafter) i.e. as set out in the following scheme:



but with the appropriate aniline intermediate (2) (where (R³⁰)_m are substituents R²⁰, R²¹, R²², R²³ and R²⁴ as set out in Table 1) and quinoline where R² and R³ are as defined in Table 1, the following compounds set out in Table 7 were prepared.

Table 7

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
2	m/e 427 (M ⁺ +H)	(d-6-DMSO, d values) 3.72 (s, 3H), 3.96 (s, 3H), 3.98 (s, 3H), 6.87 (d, 2H), 6.98 (d, 2H), 7.10 (d, 2H), 7.18 (d, 2H), 7.46 (s, 1H), 8.04 (s, 1H), 8.67 (s, 1H), 2NH assumed under H ₂ O, (2.5-3.6).	165°C/2.5h/ cyclohexanol				
3	m/e 462/ 464 (M ⁺ +H)		160°C/5h/ cyclohexanol				
4	m/e 462/ 464 (M ⁺ +H)		160°C/5h/ cyclohexanol				
5	m/e 458 (M ⁺ +H)	(d-6-DMSO, d values) 3.70 (s, 6H), 3.90 (s, 3H), 3.95 (s, 3H), 6.80 (d, 2H), 6.85 (d, 2H), 7.10 (t, 1H), 7.25 (d, 1H), 7.40 (s, 1H), 8.05 (s, 1H), 8.85 (s, 1H), 10.80 (broad s, 1H)	110°C/4h/ 1-PrOH	m/e 276 (M ⁺ +H)	KOtBu, MeOH	m/e 246 (M ⁺ +H)	H ₂ , Pd/C, EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
6	m/e 442 (M ⁺ +H)	(d-6-DMSO, d values) 2.05 (s, 3H), 3.65 (s, 3H), 3.95 (s, 3H), 4.00 (s, 3H), 6.80 (d, 2H), 6.90 (d, 1H), 7.00 (d, 1H), 7.15 (t, 1H), 7.35 (d, 2H), 7.40 (s, 1H), 8.05 (s, 1H), 8.80 (s, 1H), 10.90 (broad s, 1H)	110°C/4h/ 1-PrOH	m/e 230 (M ⁺ +H)	KOtBu, MeOH	m/e 260 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
7	m/e 428 (M ⁺ +H)	(d-6-DMSO, d values) 3.70 (s, 3H), 4.00 (s, 3H), 4.00 (s, 3H), 6.55 (s, 1H), 6.60 (m, 1H), 6.65 (dd, 1H), 7.15 (d, 2H), 7.25 (t, 1H), 7.45 (s, 1H), 7.50 (d, 2H), 8.05 (s, 1H), 8.85 (s, 1H), 11.10 (broad s, 1H)	110°C/4h/1- PrOH	m/e 216 (M ⁺ +H)			
8	m/e 428 (M ⁺ +H)	(d-6-DMSO, d values) 3.70 (s, 3H), 4.00 (s, 6H), 6.55 (s, 1H), 6.95 (m, 2H), 7.00 (d, 2H), 7.05 (d, 2H), 7.40 (d, 2H), 7.40 (s, 1H), 8.05 (s, 1H), 8.85 (s, 1H), 10.90 (broad s, 1H)	110°C/4h/1- PrOH	m/e 246 (M ⁺ +H)	KOtBu, MeOH	m/e 216 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
9	m/e 504 (M ⁺ +H)	(d-6-DMSO, d values) 3.73 (s, 3H), 3.97 (s, 3H), 5.32 (s, 2H), 6.95 (m, 3H), 7.05 (d, 1H), 7.18 (m, 2H), 7.38 (m, 5H), 7.51 (d, 2H), 7.58 (s, 1H), 8.17 (s, 1H), 8.87 (s, 1H), 11.13 (broad, 1H)	1-PrOH / 115° / 5 h				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
10	m/e 458 (M ⁺ +H)	(d-6-DMSO, d values) 3.65 (s, 3H), 3.80 (s, 3H), 4.00 (s, 6H), 6.65 (d, 1H), 6.90 (d, 1H), 7.05 (m, 3H), 7.40 (d, 2H), 7.45 (m, 1H), 8.15 (m, 1H), 8.90 (s, 1H)	110°C/18h/1 -PrOH	m/e 276 (M ⁺ +H)	KOtBu, DMA	m/e 246 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
11	m/e 458 (M ⁺ +H)	(d-6-DMSO, d values) 3.70 (s, 6H), 4.00 (s, 6H), 6.20 (d, 2H), 6.25 (t, 1H), 7.20 (d, 2H), 7.45 (s, 1H), 7.50 (d, 2H), 8.15 (s, 1H), 8.90 (s, 1H), 11.10 (broad s, 1H)	110°C/18h/1 -PrOH	m/e 276 (M ⁺ +H)	KOtBu, DMA	m/e 246 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
12	m/e 456 (M ⁺ +H)	(d-6-DMSO, d values) 1.20 (d, 6H), 4.00 (s, 6H), 4.6 (m, 1H), 6.95 (m, 3H), 7.05 (d, 1H), 7.20 (d, 2H), 7.40 (d, 2H), 7.50 (s, 1H), 8.20 (s, 1H), 8.90 (s, 1H), 11.10 (broad s, 1H)	110°C/18h/1 -PrOH	m/e 274 (M ⁺ +H)	KOtBu, DMA	m/e 244 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
13	m/e 486 (M ⁺ +H)	(d-6-DMSO, d values) 3.70 (s, 3H), 3.75 (s, 3H), 4.05 (s, 6H), 6.55 (d, 1H), 6.85 (dd, 1H), 7.15 (d, 2H), 7.50 (s, 1H), 7.55 (d, 2H), 7.85 (d, 1H), 8.20 (s, 1H), 8.95 (s, 1H), 11.20 (broad s, 1H)	110°C/18h/1 -PrOH	m/e 304 (M ⁺ +H)	KOtBu, DMA	m/e 274 (M ⁺ +H)	H ₂ , Pd/C, EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
15	m/e	(d-6-DMSO, d values) 3.75 (s, 3H), 4.00 (s, 6H), 6.55 (t, 1H), 6.60 (t, 1H), 6.80 (t, 1H), 7.20 (d, 2H), 7.50 (s, 1H), 7.55 (d, 2H), 8.20 (s, 1H), 8.95 (s, 1H), 11.20 (broad s, 1H)	110°C/18h/1-PrOH		KOtBu, DMA	m/e 250 (M ⁺ +H)	SnCl ₂ .2H ₂ O, HCl, EtOAc
	462 (M ⁺ +H)						
32	m/e	(d-6-DMSO, d values) 1.20 (t, 3H), 3.95 (s, 6H), 4.00 (q, 2H), 6.95 (m, 3H), 7.05 (m, 1H), 7.15 (m, 2H), 7.35 (d, 2H), 7.45 (s, 1H), 8.10 (s, 1H), 8.85 (s, 1H), 10.95 (broad s, 1H)	110°C/4h/1-PrOH	m/e 260 (M ⁺ +H)	KOtBu, MeOH	m/e 230 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
	442 (M ⁺ +H)						
42	m/e 516 (M ⁺ +H)	(d-6-DMSO, d values), 3.35 (s, 6H), 3.74 (s, 3H), 3.76 (m, 4H), 4.32 (m, 4H), 6.97 (m, 3H), 7.05 (d, 1H), 7.07 (m, 2H), 7.39 (d, 2H), 7.47 (s, 1H), 8.14 (s, 1H), 8.89 (s, 1H), 10.96 (broad, 1H)	1-PrOH / reflux / 18h	m/e 337 (M ⁺ +H)	POCl ₃ / 120° / 2h		
43	m/e	(CDCl ₃ , d values) 2.25 (s, 3H), 3.60 (s, 3H), 3.80 (s, 3H), 4.00 (s, 3H), 6.60 (broad s, 1H), 6.80 (m, 2H), 7.00 (m, 5H), 7.15 (td, 1H), 7.30 (s, 1H), 8.60 (s, 1H)	110°C/36h/1-PrOH	m/e 260 (M ⁺ +H)	KOtBu, DMA	m/e 230 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
	442 (M ⁺ +H)						

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
45	m/e 516 (M ⁺ +H)	(d-6-DMSO, d values), 3.49 (m, 6H), 3.71 (s, 3H), 3.77 (m, 4H), 4.33 (m, 4H), 6.60 (m, 2H), 6.70 (d, 1H), 7.17 (d, 2H), 7.28 (t, 1H), 7.47 (d, 2H), 7.50 (s, 1H), 8.16 (s, 1H), 8.90 (s, 1H), 11.02 (broad, 1H)	1-PrOH / reflux / 18 h				
46	m/e 546 (M ⁺ +H)	(d-6-DMSO, d values), 3.35 (m, 6H), 3.69 (s, 6H), 3.77 (m, 4H), 4.33 (m, 4H), 6.19 (d, 2H), 6.26 (t 1H), 7.19 (m, 2H), 7.49 (m, 3H), 8.19 (s, 1H), 8.91 (s, 1H), 11.12 (broad, 1H)	1-PrOH / reflux / 18 h				
47	m/e 530 (M ⁺ +H)	(d-6-DMSO, d values), 1.21 (t, 3H), 3.35 (m, 6H), 3.77 (m, 4H), 4.03 (q, 2H), 4.32 (m, 4H), 6.97 (m, 3H), 7.05 (d, 1H), 7.18 (m, 2H), 7.39 (d, 2H), 7.47 (s, 1H), 8.14 (s, 1H), 8.89 (s, 1H), 10.95 (broad, 1H)	1-PrOH / reflux / 18 h				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
49	m/e 500 (M+H) ⁺	(d-6-DMSO, d values) 1.21 (t, 3H), 3.72 (s, 3H), 4.01 (s, 3H), 4.17 (q, 2H), 4.98 (s, 2H), 6.96 (m, 3H), 7.05 (m, 1H), 7.19 (m, 2H), 7.42 (m, 3H), 8.06 (s, 1H), 8.89 (s, 1H)	100°C/6h/1- PrOH			m/e 321, 323 (M+H) ⁺	RT/30mins /ethylbrom oacetate/K OtBu/n- Bu ₄ NI/DM A
56	m/e 456 (M ⁺ +H)	(CDCl ₃ , d values) 1.30 (q, 3H), 2.25 (s, 3H), 3.60 (s, 3H), 4.00 (s, 3H), 4.05 (t, 2H), 6.60 (m, 1H), 6.75 (m, 2H), 6.90 (m, 1H), 7.00 (m, 4H), 7.15 (m, 1H), 7.30 (s, 1H), 8.55 (s, 1H)	110°C/36h/1 -PrOH	m/e 274 (M ⁺ +H)	KOtBu, DMA	m/e 244 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
62	m/e 428 (M+H) ⁺	(d-6-DMSO, d values) 1.21 (t, 3H), 3.97 (s, 3H), 4.03 (q, 2H), 6.96 (m, 3H), 7.05 (m, 1H), 7.37 (m, 2H), 7.40 (s, 1H), 8.05 (s, 1H), 8.88 (s, 1H)	100°C/18h/1 -PrOH				
65	m/e 500 (M+H) ⁺	(d-6-DMSO, d values) 1.24 (t, 3H), 3.72 (s, 3H), 3.97 (s, 3H), 4.20 (q, 2H), 5.05 (s, 2H), 6.95 (m, 3H), 7.05 (m, 1H), 7.18 (m, 2H), 7.27 (s, 1H), 7.37 (d, 2H), 8.07 (s, 1H), 8.84 (s, 1H)	100°C/18h/1 -PrOH				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
69	m/e 541 (M ⁺ +H)	(d-6-DMSO, d values) 2.34 (m, 2H), 3.12 (m, 2H), 3.50 (m, 4H), 3.73 (s, 3H), 3.85 (m, 2H), 3.98 (s, 2H), 4.02 (s, 3H), 4.33 (t, 2H), 6.62 (m, 2H), 6.72 (m, 1H), 7.20 (d, 2H), 7.30 (t, 1H), 7.49 (d, 2H), 7.54 (s, 1H), 8.21 (s, 1H), 8.89 (s, 1H), 11.08 (broad, 2H)	1-PrOH / 1.0M ethereal HCl (1 equiv.) / 110deg / 3 h				
74	m/e 446 (M ⁺ +H)	(d-6-DMSO, d values) 3.75 (s, 3H), 4.00 (s, 3H), 4.00 (s, 3H), 6.90 (d, 2H), 7.00 (m, 2H), 7.25 (dd, 1H), 7.40 (d, 2H), 7.45 (s, 1H), 8.15 (s, 1H), 8.90 (s, 1H), 11.10 (broad s, 1H)	110°C/18h/1 -PrOH		KOtBu, DMA	m/e 234 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
75	m/e 432 (M ⁺ +H)	(d-6-DMSO, d values) 3.98 (s, 6H), 7.05 (d, 2H), 7.15 (d, 2H), 7.40 (s, 1H), 7.42 (d, 2H), 7.50 (d, 2H), 8.10 (s, 1H), 8.85 (s, 1H)					
76	m/e 443 (M ⁺ +H)	(d-6-DMSO, d values) 3.99 (s, 6H), 7.15-7.30 (m, 4H), 7.48-7.52 (m, 3H), 8.05 (s, 1H), 8.11 (d, 2H), 8.68 (s, 1H), NH assumed under H ₂ O, (3.2-3.4).	165°C/2.5h/ cyclohexanol				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
77	m/e 434 (M ⁺ +H)	(d-6-DMSO, d values) 3.92 (s, 3H), 3.94 (s, 3H), 6.95 (m, 1H), 7.05 (d, 2H), 7.05 - 7.25 (m, obscured), 7.29 (d, 2H), 7.4 - 7.5 (m, 1H), 7.75 (s, 1H), 8.40 (s, 1H), 9.43 (s, 1H)	150°C/16h/ Dowtherm A				
78	m/e 462/ 464 (M ⁺ +H)		150°C/16h/ Dowtherm A				
79	m/e 448/ 450 (M ⁺ +H)	(d-6-DMSO, d values) 3.96 (s, 3H), 3.98 (s, 3H), 7.30 (d, 2H), 7.37 (d, 4H), 7.45 (m, 3H), 8.04 (s, 1H), 8.7 (s, obscured).	160°C/5h/ cyclohexanol				
80	m/e 446/ 448 (M ⁺ +H)		160°C/5h/ cyclohexanol				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
81	m/e 416 (M ⁺ +H)	(d-6-DMSO, d values) 4.00 (s, 6H), 7.10 (d, 2H), 7.15 (m, 1H), 7.20 (m, 2H), 7.40 (m, 1H), 7.45 (m, 3H), 8.20 (s, 1H), 8.90 (s, 1H), 11.12 (broad s, 1H)	110°C/4h/1- PrOH		KOtBu, MeOH	m/e 204 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
82	m/e 412 (M ⁺ +H)	(d-6-DMSO, d values) 2.10 (s, 3H), 4.00 (s, 6H), 6.95 (m, 3H), 7.10 (t, 1H), 7.20 (t, 1H), 7.40 (d, 2H), 7.45 (s, 1H), 8.15 (s, 1H), 8.85 (s, 1H), 11.05 (broad s, 1H)	110°C/4h/1- PrOH		KOtBu, MeOH	m/e 200 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
83	m/e 514 (M ⁺ +H)	(d-6-DMSO, d values) 4.00 (s, 3H), 4.00 (s, 3H), 5.05 (s, 2H), 7.45 (d, 2H), 7.45 (s, 1H), 7.55 (d, 2H), 7.60 (s, 2H), 8.05 (s, 1H), 8.95 (s, 1H)	110°C/18h/1 -PrOH				
84	m/e 486 (M ⁺ +H)	(d-6-DMSO, d values) 3.80 (s, 3H), 3.95 (s, 3H), 4.00 (s, 3H), 4.35 (s, 2H), 7.35 (d, 2H), 7.45 (m, 4H), 7.60 (d, 1H), 7.80 (d, 1H), 8.00 (s, 1H), 8.05 (s, 1H), 8.90 (s, 1H), 10.90 (broad s, 1H)	110°C/18h/1 -PrOH				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
85	m/e	(d-6-DMSO, d values) 2.4 (s, 3H), 4.00 (s, 6H), 6.90 (dd, 1H), 7.05 (d, 2H), 7.20 (m, 2H), 7.35 (dd, 1H), 7.45 (m, 3H), 8.10 (s, 1H), 8.85 (s, 1H) 10.90 (broad s, 1H)	110°C/5.5h/1 -PrOH		KOtBu,	m/e	H ₂ , Pd/C,
	444 (M ⁺ +H)				MeOH, DMA	232 (M ⁺ +H)	EtOAc
86	m/e	(d-6-DMSO, d values) 4.00 (s, 6H), 6.95 (d, 1H), 7.25 (t, 1H), 7.30 (d, 2H), 7.45 (s, 1H), 7.55 (d, 2H), 7.60 (m, 1H), 7.90 (dd, 1H), 8.15 (s, 1H), 8.90 (s, 1H)	110°C/5.5h/1 -PrOH	m/e	KOtBu,	m/e	H ₂ , Pd/C,
	423 (M ⁺ +H)			239 (M-H)	MeOH, DMA	211 (M ⁺ +H)	EtOAc
87	m/e	(d-6-DMSO, d values) 4.00 (s, 6H), 7.05 (dd, 1H), 7.15 (m, 3H), 7.35 (t, 1H), 7.40 (s, 1H), 7.45 (m, 3H), 8.05 (s, 1H), 8.80 (s, 1H)	110°C/18h/1 -PrOH		KOtBu,	m/e	SnCl ₂ .2H ₂ O
	524 (M ⁺ +H)				DMA	312 (M ⁺ +H)	, EtOAc
88	m/e	(d-6-DMSO, d values) 4.00 (s, 6H), 7.10 (m, 4H), 7.40 (td, 1H), 7.45 (s, 1H), 7.45 (d, 2H), 7.75 (dd, 1H), 8.05 (s, 1H), 8.90 (s, 1H), 11.05 (broad s, 1H)	110°C/18h/1 -PrOH		KOtBu,	m/e	SnCl ₂ .2H ₂ O
	476 (M ⁺ +H)				DMA	264 (M ⁺ +H)	, EtOAc
89	m/e	(d-6-DMSO, d values) 3.95 (s, 3H), 3.95 (s, 3H), 7.00 (m, 1H), 7.20 (m, 3H), 7.30 (m, 3H), 7.40 (d, 2H), 7.90 (s, 1H), 8.60 (s, 1H)	110°C/18h/1 -PrOH		KOtBu,	m/e	SnCl ₂ .
	476 (M ⁺ +H)				DMA	264 (M ⁺ +H)	2H ₂ O, EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
90	m/e	(d-6-DMSO, d values) 3.95 (s, 3H), 3.95 (s, 3H), 7.00 (m, 1H), 7.20 (m, 3H), 7.30 (m, 3H), 7.40 (d, 2H), 7.90 (s, 1H), 8.60 (s, 1H)	110°C/18h/1 -PrOH		KOtBu,	m/e	SnCl ₂ .2H ₂ O
	476 (M ⁺ +H)				DMA	264 (M ⁺ +H)	, EtOAc
91	m/e	(d-6-DMSO, d values) 4.00 (s, 6H), 7.00 (m, 2H), 7.20 (dd, 1H), 7.20 (d, 2H), 7.40 (t, 1H), 7.45 (s, 1H), 7.50 (d, 2H), 8.20 (s, 1H), 8.95 (s, 1H), 11.20 (broad s, 1H)	110°C/18h/1 -PrOH		KOtBu,	m/e	SnCl ₂ .2H ₂ O
	432 (M ⁺ +H)				DMA	220 (M ⁺ +H)	, EtOAc
92	m/e	(d-6-DMSO, d values) 4.00 (s, 6H), 6.95 (d, 2H), 7.05 (m, 2H), 7.40 (m, 1H), 7.45 (s, 1H), 7.45 (d, 2H), 7.90 (d, 1H), 8.15 (s, 1H), 8.90 (s, 1H), 11.05 (broad s, 1H)	110°C/18h/1 -PrOH		KOtBu,	m/e	SnCl ₂ .
	524 (M ⁺ +H)				DMA	312 (M ⁺ +H)	2H ₂ O, EtOAc
93	m/e	(d-6-DMSO, d values) 4.00 (s, 6H), 7.05 (d, 1H), 7.15 (m, 3H), 7.45 (s, 1H), 7.55 (d, 2H), 7.60 (d, 1H), 8.15 (s, 1H), 8.95 (s, 1H), 11.10 (broad s, 1H)	110°C/18h/1 -PrOH				
	466 (M ⁺ +H)						

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
94	m/e	(d-6-DMSO, d values) 4.00 (s, 6H), 7.10 (t, 3H), 7.20 (t, 1H), 7.35 (t, 1H), 7.50 (s, 1H), 7.55 (d, 2H), 7.60 (d, 1H), 8.20 (s, 1H), 8.95 (s, 1H), 11.20 (broad s, 1H)	110°C/18h/1 -PrOH	m/e	KOtBu,	m/e	SnCl ₂ .2H ₂ O
	432 (M ⁺ +H)			243 (M ⁺ +H)	DMA	220 (M ⁺ +H)	, HCl, EtOAc
95	m/e	(d-6-DMSO, d values) 2.05 (s, 3H), 4.00 (s, 6H), 6.65 (m, 1H), 7.15 (d, 2H), 7.30 (d, 2H), 7.45 (m, 4H), 8.20 (s, 1H), 8.95 (s, 1H), 10.10 (broad s, 1H), 11.20 (broad s, 1H)	110°C/18h/1 -PrOH	m/e	KOtBu,		H ₂ , Pd/C,
	455 (M ⁺ +H)			273 (M ⁺ +H)	DMA		EtOAc
96	m/e	(d-6-DMSO, d values) 4.00 (s, 6H), 6.80 (m, 1H), 6.95 (m, 5H), 7.35 (d, 2H), 7.40 (s, 1H), 8.00 (s, 1H), 8.75 (s, 1H), 9.60 (broad s, 1H), 10.50 (broad s, 1H)	110°C/18h/1 -PrOH				H ₂ , Pd/C,
	414 (M ⁺ +H)						EtOAc
97	m/e	(d-6-DMSO, d values) 4.00 (s, 6H), 5.25 (s, 2H), 7.05 (m, 3H), 7.30 (m, 6H), 7.50 (m, 3H), 7.60 (m, 1H), 7.90 (dd, 1H), 8.15 (s, 1H), 8.90 (s, 1H), 11.10 (broad s, 1H)	110°C/18h/1 -PrOH	m/e	KOtBu,	m/e	SnCl ₂ .2H ₂
	532 (M ⁺ +H)			350 (M ⁺ +H)	DMA	320 (M ⁺ +H)	O, HCl, EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
98	m/e 466 (M ⁺ +H)	(d-6-DMSO, d values) 4.00 (s, 6H), 7.00 (d, 1H), 7.20 (d, 2H), 7.30 (t, 1H), 7.50 (s, 1H), 7.55 (d, 2H), 7.60 (t, 1H), 7.80 (d, 1H), 8.20 (s, 1H), 8.95 (s, 1H), 11.20 (broad s, 1H)	110°C/18h/1 -PrOH		KOtBu, DMA	m/e 254 (M ⁺ +H)	SnCl ₂ .2H ₂ O, HCl, EtOAc
99	m/e 466 (M ⁺ +H)	(d-6-DMSO, d values) 4.00 (s, 3H), 4.00 (s, 3H), 7.30 (m, 3H), 7.35 (d, 1H), 7.50 (m, 2H), 7.55 (d, 2H), 7.60 (t, 1H), 8.20 (s, 1H), 8.95 (s, 1H), 11.20 (broad s, 1H)	110°C/18h/1 -PrOH		KOtBu, DMA	m/e 254 (M ⁺ +H)	SnCl ₂ . 2H ₂ O, HCl, EtOAc
100	m/e 442 (M ⁺ +H)	(d-6-DMSO, d values) 4.00 (s, 6H), 6.95 (d, 1H), 7.05 (d, 2H), 7.20 (t, 1H), 7.45 (d, 2H), 7.50 (s, 1H), 7.55 (m, 1H), 7.80 (dd, 1H), 8.20 (s, 1H), 8.95 (s, 1H), 11.20 (broad s, 1H)	110°C/18h/1 -PrOH	m/e 350 (M ⁺ +H)	KOtBu, DMA	m/e 230 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
101	m/e 441 (M ⁺ +H)	(d-6-DMSO, d values) 1.15 (t, 3H), 3.00 (q, 2H), 4.00 (s, 6H), 6.25 (dd, 1H), 6.30 (t, 1H), 6.40 (dd, 1H), 7.10 (m, 3H), 7.45 (d, 2H), 7.50 (s, 1H), 8.15 (s, 1H), 8.85 (s, 1H), 11.00 (broad s, 1H)	110°C/18h/1 -PrOH	m/e 259 (M ⁺ +H)	KOtBu, DMA	m/e 229 (M ⁺ +H)	H ₂ , Pd/C, EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
103	m/e	(d-6-DMSO, d values) 3.75 (s, 3H), 4.00 (s, 6H), 7.10 (s, 1H), 7.10 (d, 2H), 7.30 (t, 1H), 7.50 (m, 3H), 7.60 (t, 1H), 7.90 (d, 1H), 8.20 (s, 1H), 8.95 (s, 1H), 11.20 (broad s, 1H)	110°C/18h/1 -PrOH	m/e	KOtBu,	m/e	H ₂ , Pd/C,
	456 (M ⁺ +H)			274 (M ⁺ +H)	DMA	244 (M ⁺ +H)	EtOAc
104	m/e	(d-6-DMSO@373K, d values) 1.10 (t, 6H), 3.30 (q, 4H), 4.00 (s, 6H), 6.35 (dd, 1H), 6.50 (s, 1H), 6.60 (dd, 1H), 7.10 (d, 2H), 7.20 (t, 1H), 7.40 (d, 2H), 7.50 (s, 1H), 8.05 (s, 1H), 8.65 (s, 1H)	110°C/18h/1 -PrOH	m/e	KOtBu,	m/e	H ₂ , Pd/C,
	469 (M ⁺ +H)			287 (M ⁺ +H)	DMA	257 (M ⁺ +H)	EtOAc
105	m/e	(d-6-DMSO, d values) 4.00 (s, 3H), 4.00 (s, 3H), 7.30 (d, 2H), 7.40 (m, 2H), 7.50 (m, 5H), 8.30 (s, 1H), 8.95 (s, 1H), 11.60 (broad s, 1H)	110°C/18h/1 -PrOH		KOtBu,	m/e	SnCl ₂ .2H ₂
	423 (M ⁺ +H)				DMA	211 (M ⁺ +H)	O, HCl, EtOAc
106	m/e	(CDCl ₃ , d values) 2.10 (s, 3H), 2.25 (s, 3H), 3.80 (s, 3H), 4.00 (s, 3H), 6.80 (dd, 1H), 6.90 (m, 2H), 7.00 (d, 1H), 7.10 (m, 3H), 7.30 (m, 1H), 7.35 (s, 1H), 7.50 (broad s, 1H), 8.55 (s, 1H)	110°C/36h/1 -PrOH	m/e	KOtBu,	m/e	H ₂ , Pd/C,
	469 (M ⁺ +H)			287 (M ⁺ +H)	DMA	257 (M ⁺ +H)	EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
107	m/e 437 (M ⁺ +H)	(d-6-DMSO, d values) 2.25 (s, 3H), 4.00 (s, 3H), 4.00 (s, 3H), 7.00 (d, 1H), 7.15 (dd, 1H), 7.25 (m, 2H), 7.50 (m, 2H), 7.60 (td, 1H), 7.90 (dd, 1H), 8.30 (s, 1H), 8.90 (s, 1H), 11.20 (broad s, 1H)	110°C/60h/1 -PrOH	m/e 255 (M ⁺ +H)	KOtBu, DMA	m/e 225 (M ⁺ +H)	SnCl ₂ .2H ₂ O, HCl, EtOAc
108	m/e 500 (M+H) ⁺	(d-6-DMSO, d values) 4.02 (s, 3H), 6.74 (tt, 1H), 6.89 (m, 1H), 7.03 (m, 2H), 7.22 (d, 2H), 7.46 (m, 3H), 7.50 (1H, s), 7.95 (s, 1H), 8.88 (s, 1H)	100°C/18h/1 -PrOH				
109	m/e 438 (M+H) ⁺	(d-6-DMSO, d values) 3.54 (m, 1H), 4.01 (s, 3H), 4.80 (m, 2H), 6.99 (m, 4H), 7.18 (m, 1H), 7.25 (m, 1H), 7.38 (d, 2H), 7.48 (1H, s), 7.94 (s, 1H), 8.88 (s, 1H)	100°C/18h/1 -PrOH		60°C/1h/ K ₂ CO ₃ / HCCCCH ₂ Br/acetone	m/e 240 (M+H) ⁺	90°C/2h/Sn Cl ₂ .2H ₂ O/ EtOAc
110	m/e 409 (M ⁺ +H)	(d-6-DMSO, d values) 4.0 (s, 3H), 6.97 (d, 1H), 7.23-7.35 (m, 3H), 7.47 (s, 1H), 7.51 (d, 2H), 7.63 (t, 1H), 7.9 (d, 1H), 7.95 (s, 1H), 8.89 (s, 1H), 10.5 (br.s, 1H), 10.85 (br.s, 1H)	82°C/20h/iso -PrOH				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
111	m/e 441	(d-6-DMSO, d values) 2.80 (s, 6H), 4.00 (s, 6H), 6.95 (m, 2H), 7.05 (d, 2H), 7.20 (m, 2H), 7.40 (d, 2H), 7.40 (s, 1H), 8.10 (s, 1H), 8.85 (s, 1H), 10.90 (broad s, 1H)	110°C/18h/1 -PrOH	m/e 259 (M ⁺ +H)	KOtBu, DMA/ HCHO, AcOH, NaBH ₃ C N, EtOH	m/e 229 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
112	m/e 441	(d-6-DMSO, d values) 2.90 (s, 6H), 4.00 (s, 6H), 6.35 (m, 2H), 6.50 (d, 1H), 7.15 (m, 3H), 7.45 (d, 2H), 7.50 (s, 1H), 8.15 (s, 1H), 8.90 (s, 1H), 11.10 (broad s, 1H)	110°C/18h/1 -PrOH	m/e 259 (M ⁺ +H)	HCHO, AcOH, NaBH ₃ C N, EtOH	m/e 229 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
113	m/e 500 (M+H) ⁺	(d-6-DMSO, d values) 3.97 (s, 3H), 6.74 (tt, 1H), 6.89 (m, 1H), 7.03 (m, 2H), 7.24 (d, 2H), 7.34 (s, 1H), 7.45 (d, 1H), 7.51 (d, 2H), 8.04 (s, 1H), 8.87 (s, 1H)	100°C/18h/1 -PrOH				
116	m/e 441 (M ⁺ +H)	(d-6-DMSO, d values) 4.00 (s, 6H), 6.95 (d, 1H), 7.00 (d, 1H), 7.40 (m, 4H), 7.85 (d, 2H), 7.95 (dd, 1H), 8.15 (s, 1H), 8.95 (s, 1H), 10.55 (broad s, 1H), 11.10 (broad s, 1H), 11.70 (broad s, 1H)	110°C/70h/1 -PrOH	m/e 257 (M-1H)	KOtBu, DMA	m/e 229 (M ⁺ +H)	SnCl ₂ .2H ₂ O , HCl, EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
117	m/e 427 (M ⁺ +H)	(d-6-DMSO, d values) 2.63 (s, 3H), 3.97 (d, 6H), 6.24 (m, 3H), 6.38 (d, 1H), 7.10 (m, 3H), 7.45 (t, 3H), 8.19 (s, 1H), 8.90 (s, 1H), 11.17 (broad, 1H)	1-PrOH / 110 deg / 18h			m/e 215 (M ⁺ +H)	H ₂ / Pd/C / EtOAc / RT / ambient pressure
122	m/e 439 (M+H) ⁺	(d-6-DMSO, d values) 4.00 (s, 3H), 5.14 (s, 2H), 7.01 (d, 2H), 7.09 (m, 2H), 7.23 (m, 1H), 7.33 (d, 1H), 7.40 (d, 2H), 7.45 (s, 1H), 7.94 (s, 1H), 8.87 (s, 1H)	100°C/18h/1 -PrOH		60°C/1h/ K ₂ CO ₃ /b romoacet onitrile/a cetone	m/e 241 (M+H) ⁺	90°C/2h/Sn Cl ₂ .2H ₂ O/ EtOAc
123	m/e 444 (M+H) ⁺	(d-6-DMSO, d values) 3.60 (t, 2H), 4.00 (m, 5H), 6.98 (m, 4H), 7.17 (m, 2H), 7.27 (d, 2H), 7.46 (s, 1H), 7.93 (s, 1H), 8.87 (s, 1H)	100°C/18h/1 -PrOH				
124	m/e 439 (M+H) ⁺	(d-6-DMSO, d values) 3.95 (s, 3H), 5.15 (s, 2H), 7.03 (d, 2H), 7.10 (m, 2H), 7.24 (m, 1H), 7.31 (m, 1H), 7.41 (m, 2H), 7.45 (m, 1H), 8.08 (s, 1H), 8.83 (s, 1H)	100°C/18h/1 -PrOH				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
125	m/e 444 (M+H) ⁺	(d-6-DMSO, d values) 3.60 (t, 2H), 3.96 (s, 3H), 3.98 (t, 2H), 7.00 (m, 4H), 7.16 (m, 2H), 7.37 (s, 1H), 7.42 (m, 2H), 8.10 (s, 1H), 8.84 (s, 1H)	100°C/18h/1 -PrOH				
126	m/e 440 (M+H) ⁺	(d-6-DMSO, d values) 3.89 (s, 3H), 4.55 (m, 2H), 5.17 (dd, 1H), 5.29 (dd, 1H), 5.92 (m, 1H), 6.89 (d, 2H), 6.93 (m, 1H), 7.02 (m, 1H), 7.13 (m, 2H), 7.16 (s, 1H), 7.21 (d, 2H), 7.72 (s, 1H), 8.29 (s, 1H), 9.34 (s, 1H)	100°C/18h/1 -PrOH		60°C/1h/ K ₂ CO ₃ / allyl bromide/ acetone	m/e 242 (M+H) ⁺	90°C/3h/Sn Cl ₂ .2H ₂ O/ EtOAc
129	m/e 471 (M+H) ⁺	(d-6-DMSO, d values) 2.61 (d, 3H), 3.98 (s, 3H), 4.46 (s, 2H), 7.00 (m, 4H), 7.04 (m, 1H), 7.12 (m, 2H), 7.33 (d, 2H), 7.41 (s, 1H), 7.49 (bs, 1H), 7.86 (s, 1H), 8.74 (s, 1H)	100°C/18h/1 -PrOH				
130	m/e 409.2 (M ⁺ +H)	(d-6-DMSO, d values) 3.98 (s, 3H), 6.95 (d, 1H), 7.22-7.4 (m, 3H), 7.42 (s, 1H), 7.5-7.7 (m, 3H), 7.9 (d, 1H), 8.09 (s, 1H), 8.89 (s, 1H), 11.1 (br.s, 1H), 11.7 (br.s, 1H)	82°C/20h/iso -PrOH				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
133	m/e 529 (M ⁺ +H)	(d-6-DMSO, d values) 1.19 (t, 3H), 3.12 (q, 2H), 3.37 (s, 6H), 3.79 (m, 4H), 4.36 (m, 4H), 6.66 (m, 3H), 7.18 (d, 2H), 7.26 (m, 1H), 7.51 (d, 2H), 7.56 (s, 1H), 8.31 (s, 1H), 8.99 (s, 1H), 11.39 (s, 1H)	EtOH / reflux / 18 h				
134	m/e 554 (M ⁺ +H)	(d-6-DMSO, d values) 1.13 (t, 3H), 2.30 (m, 2H), 3.12 (q, 2H), 3.16 (broad, 2H), 3.49 (broad, 2H), 3.80 (broad, 4H), 3.95 (s, 3H), 4.31 (t, 2H), 6.32 (m, 2H), 6.48 (m, 1H), 7.13 (m, 3H), 7.42 (m, 3H), 8.07 (s, 1H), 8.90 (s, 1H), 10.80 (broad, 1H), 10.95 (broad, 1H)	1-PrOH / 1.0M ethereal HCl (1 equiv.) / reflux / 48 h				
135	m/e 466	(CDCl ₃ , d values) 3.80 (s, 3H), 4.05 (s, 3H), 7.00 (m, 5H), 7.15 (d, 2H), 7.25 (m, 1H), 7.40 (s, 1H), 7.50 (td, 1H), 8.05 (dd, 1H), 8.45 (s, 1H), 8.60 (s, 1H)	110°C/18h/1 -PrOH	m/e 284 (M ⁺ +H)	KOtBu, DMA	m/e 254 (M ⁺ +H)	H ₂ , Pd/C, EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
141	m/e 529 (M ⁺ +H)	(d-6-DMSO, d values) 2.34 (m, 2H), 3.08 (m, 2H), 3.48 (m, 4H), 3.90 (m, 4H), 4.01 (s, 3H), 4.30 (t, 2H), 7.12 (d, 2H), 7.21 (m, 3H), 7.40 (m, 1H), 7.48 (d, 2H), 7.57 (s, 1H), 8.34 (s, 1H), 8.90 (s, 1H), 11.28 (broad, 2H)	1-PrOH / 1.0M ethereal HCl (1 equiv.) / 60deg / 72 h				
144	m/e 434 (M ⁺ +H)	(d-6-DMSO, d values) 4.00 (s, 6H), 6.95 (m, 1H), 7.20 (m, 4H), 7.50 (d, 2H), 7.50 (s, 1H), 8.20 (s, 1H), 8.95 (s, 1H), 11.25 (broad s, 1H)	110°C/18h/1 -PrOH	KOtBu, DMA	m/e 222 (M ⁺ +H)	H ₂ , Pd/C, EtOAc	
145	m/e 529 (M ⁺ +H)		1-PrOH / 1.0M ethereal HCl (1 equiv.) / 60deg / 72 h				
146	m/e 514 (M ⁺ +H)	(d-6-DMSO, d values) 4.00 (s, 6H), 6.75 (tt, 1H), 6.90 (t, 1H), 7.00 (m, 2H), 7.20 (d, 2H), 7.45 (s, 1H), 7.50 (d, 1H), 7.55 (d, 2H), 8.20 (s, 1H), 8.95 (s, 1H), 11.20 (broad s, 1H)	110°C/18h/1 -PrOH				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
147	m/e 434 (M ⁺ +H)	(d-6-DMSO, d values) 4.00 (s, 6H), 7.05 (d, 2H), 7.35 (m, 3H), 7.45 (d, 2H), 7.50 (s, 1H), 8.20 (s, 1H), 8.95 (s, 1H), 11.25 (broad s, 1H)	110°C/18h/1 -PrOH		KOtBu, DMA	m/e 222 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
148	m/e 434 (M ⁺ +H)	(d-6-DMSO, d values) 4.00 (s, 3H), 4.00 (s, 3H), 7.00 (m, 2H), 7.20 (d, 2H), 7.45 (m, 1H), 7.50 (s, 1H), 7.55 (d, 2H), 8.20 (s, 1H), 8.95 (s, 1H), 11.35 (broad s, 1H)	110°C/18h/1 -PrOH		KOtBu, DMA	m/e 222 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
149	m/e 434 (M ⁺ +H)	(d-6-DMSO, d values) 4.00 (s, 3H), 4.00 (s, 3H), 6.75 (dd, 2H), 6.95 (tt, 1H), 7.30 (d, 2H), 7.50 (s, 1H), 7.55 (d, 2H), 8.25 (s, 1H), 8.95 (s, 1H), 11.45 (broad s, 1H)	110°C/18h/1 -PrOH		KOtBu, DMA	m/e 222 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
150	m/e 500 (M ⁺ +H)	(d-6-DMSO, d values) 0.83 (t, 3H), 1.57 (m, 2H), 3.9 (s, 3H), 4.05(t, 2H), 4.8 (s, 2H), 6.9-7.04 (m, 7H), 7.18 (s, 1H), 7.23 (d, 2H), 7.72 (s, 1H), 8.3 (s, 1H), 9.34 (s, 1H)	100°C/5h/1- PrOH/HCl	m/e 333.51 (M ⁺ +H)	DMA/ KOtBu, /150°C/0 .5h	m/e 303.58 (M ⁺ +H)	Hydrogen/ 5% Pd/C/ EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
151	m/e 519.52 1 (M ⁺ +H)	(d-6-DMSO, d values) 3.43 (q, 2H), 3.6 (t, 2H), 3.9 (s, 3H), 4.5(s, 2H), 6.93-7.15 (m, 6H), 7.16 (s, 1H), 7.24 (d, 2H), 7.73 (s, 1H), 7.89 (t, 1H), 8.3 (s, 1H), 9.35 (s, 1H)	100°C/5h/1- PrOH/HCl	m/e 333.51 (M ⁺ +H)	DMA/ KOtBu, /150°C/0 .5h	m/e 303.58 (M ⁺ +H)	Hydrogen/ 5% Pd/C/ EtOAc
152	m/e 500.52 (M ⁺ +H)	(d-6-DMSO, d values) 3.16 (q, 2H), 3.4 (t, 2H), 3.9 (s, 3H), 4.47(s, 2H), 4.7(t, 1H), 6.94-7.17 (m, 7H), 7.18 (s, 1H), 7.24 (d, 2H), 7.57 (t, 1H), 7.74 (s, 1H), 8.31 (s, 1H), 9.34 (s, 1H)	100°C/5h/1- PrOH/HCl	m/e 333.51 (M ⁺ +H)	DMA/K- butoxide/ 150°C/0. 5h	m/e 303.58 (M ⁺ +H)	Hydrogen/ 5% Pd/C/EtOA c
153	m/e 515.44 (M ⁺ +H)	(d-6-DMSO, d values) 3.16 (q, 2H), 3.39 (t, 2H), 3.98 (s, 6H), 3.95 (v.br. s, 1H), 4.48(s, 2H), 6.95-7.22 (m, 6H), 7.41 (s, 1H), 7.44 (d, 2H), 7.6 (t, 1H), 8.13 (s, 1H), 8.9 (s, 1H), 11.07 (br.s, 1H)	100°C/2h/1- PrOH				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
156	m/e 470 (M ⁺ +H)	(d-6-DMSO, d values) 2.60 (s, 3H), 4.00 (s, 6H), 6.20 (broad s, 1H), 6.50 (dd, 1H), 7.00 (d, 1H), 7.10 (d, 2H), 7.20 (t, 1H), 7.35 (t, 1H), 7.45 (d, 2H), 7.45 (s, 1H), 8.15 (s, 1H), 8.80 (broad s, 1H), 8.90 (s, 1H), 11.10 (broad s, 1H)	110°C/18h/1 -PrOH			m/e 256 (M-H)	H ₂ , Pd/C, EtOAc
157	m/e 482 (M ⁺ +H)	(d-6-DMSO, d values) 4.00 (s, 3H), 4.00 (s, 3H), 7.00 (broad s, 1H), 7.05 (m, 2H), 7.25 (d, 2H), 7.50 (m, 4H), 8.25 (s, 1H), 8.95 (s, 1H), 11.40 (broad s, 1H)	110°C/18h/1 -PrOH		KOtBu, DMA	m/e 270 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
158	m/e 474 (M ⁺ +H)	(d-6-DMSO, d values) 3.80 (s, 3H), 4.00 (s, 3H), 4.00 (s, 3H), 6.80 (d, 1H), 7.15 (t, 1H), 7.20 (d, 2H), 7.50 (m, 4H), 8.20 (s, 1H), 8.95 (s, 1H), 11.25 (broad s, 1H)	110°C/18h/1 -PrOH		KOtBu, DMA	m/e 262 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
159	m/e 458 (M+H) ⁺	(d-6-DMSO, d values) 3.61 (m, 2H), 4.00 (bs, 8H), 6.98 (m, 4H), 7.17 (m, 2H), 7.42 (m, 3H), 8.13 (s, 1H), 8.90 (s, 1H)	100°C/18h/1 -PrOH				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
160	m/e 452 (M ⁺ +H)	(CDCl ₃ , d values) 3.80 (s, 3H), 4.00 (s, 3H), 6.75 (s, 1H), 6.80 (broad s, 1H), 6.95 (m, 4H), 7.10 (d, 2H), 7.35 (s, 1H), 8.60 (s, 1H)	110°C/18h/1 -PrOH		KOtBu, DMA	m/e 240 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
161	m/e 485 (M ⁺ +H) ⁺	(d-6-DMSO, d values) 2.62 (d, 3H), 3.97 (s, 6H), 4.33 (s, 2H), 7.08 (m, 6H), 7.42 (m, 3H), 7.52 (m, 1H), 8.13 (s, 1H), 8.92 (s, 1H)	100°C/18h/1 -PrOH				
162	m/e 482 (M ⁺ +H)	(d-6-DMSO, d values) 4.00 (s, 3H), 4.00 (s, 3H), 7.10 (d, 1H), 7.15 (d, 2H), 7.25 (m, 1H), 7.40 (td, 1H), 7.50 (m, 4H), 8.20 (s, 1H), 8.95 (s, 1H), 11.30 (broad s, 1H)	110°C/18h/1 -PrOH		KOtBu, DMA	m/e 270 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
163	m/e 529 (M ⁺ +H)	(d-6-DMSO, d values) 2.33 (m, 2H), 3.12 (m, 2H), 3.50 (m, 4H), 3.83 (t, 2H), 3.99 (s, 2H), 4.02 (s, 3H), 4.36 (t, 2H), 7.12 (m, 4H), 7.26 (m, 2H), 7.48 (d, 2H), 7.52 (s, 1H), 8.18 (s, 1H), 8.88 (s, 1H), 10.92 (broad, 2H)	1-PrOH / 1.0M ethereal HCl (1 equiv.) / 110deg / 48 h				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
164	m/e 529 (M ⁺ +H)	(d-6-DMSO, d values) 2.34 (m, 2H), 3.12 (m, 2H), 3.49 (m, 4H), 3.83 (t, 2H), 4.00 (m, 5H), 4.32 (t, 2H), 7.15 (m, 3H), 7.27 (m, 1H), 7.50 (m, 4H), 8.16 (s, 1H), 8.88 (s, 1H), 10.94 (broad, 2H)	1-PrOH / 1.0M ethereal HCl (1 equiv.) / 110deg / 48h				
165	m/e 550 (M ⁺ +H)	(d-6-DMSO, d values) 2.28 (s, 3H), 2.34 (m, 2H), 3.12 (m, 2H), 3.29 (m, 2H), 3.50 (m, 2H), 3.84 (t, 2H), 4.02 (m, 5H), 4.33 (t, 2H), 7.02 (d, 1H), 7.18 (m, 1H), 7.29 (m, 2H), 7.53 (d, 2H), 7.64 (m, 1H), 7.92 (m, 1H), 8.27 (s, 1H), 8.88 (s, 1H), 11.00 (broad, 2H)	1-PrOH / 1.0M ethereal HCl (1 equiv.) / 110deg / 48h				
166	m/e 480 (M ⁺ +H)	(d-6-DMSO@373K, d values) 2.60 (s, 3H), 4.00 (s, 6H), 7.05 (d, 2H), 7.10 (d, 1H), 7.35 (t, 1H), 7.40 (d, 2H), 7.55 (s, 1H), 7.55 (t, 1H), 7.95 (dd, 1H), 8.15 (s, 1H), 8.70 (s, 1H)	110°C/18h/1 -PrOH		KOtBu, DMF	m/e 268 (M ⁺ +H)	Na ₂ S ₂ O ₄ , EtOH, H ₂ O

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
167	m/e 466 (M ⁺ +H)	(CDCl ₃ , d values) 3.80 (s, 3H), 4.00 (s, 3H), 7.00 (s, 1H), 7.05 (d, 2H), 7.05 (s, 1H), 7.20 (d, 2H), 7.20 (d, 1H), 7.40 (s, 1H), 7.50 (t, 1H), 7.70 (t, 1H), 7.80 (d, 1H), 8.45 (s, 1H), 8.60 (s, 1H)	110°C/18h/1 -PrOH		KOtBu, DMA	m/e 254 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
169	m/e 611 (M+H) ⁺	(d-6-DMSO, d values) 0.91 (t, 3H), 1.53 (m, 2H), 2.33 (m, 2H), 3.08 (m, 2H), 3.26 (m, 2H), 3.35-3.50 (m, 2H (under H ₂ O signal)), 3.68 (s, 2H), 3.81 (m, 2H), 3.95 (m, 4H), 3.99 (s, 3H), 4.29 (m, 2H), 6.87 (d, 1H), 7.04 (d, 2H), 7.10 (m, 1H), 7.26 (m, 1H), 7.37 (d, 1H), 7.46 (d, 2H), 7.54 (s, 1H), 8.20 (s, 1H), 8.89 (s, 1H)	100°C/18h/1 -PrOH				
171	m/e 480 (M ⁺ +H)	(d-6-DMSO, d values) 2.40 (s, 3H), 4.00 (s, 3H), 4.00 (s, 3H), 7.05 (d, 1H), 7.10 (d, 2H), 7.35 (t, 1H), 7.50 (d, 2H), 7.50 (s, 1H), 7.60 (t, 1H), 8.10 (d, 1H), 8.20 (s, 1H), 8.90 (s, 1H), 11.25 (broad s, 1H)	110°C/18h/1 -PrOH.		KOtBu, DMF	m/e 268 (M ⁺ +H)	Na ₂ S ₂ O ₄ , EtOH, H ₂ O

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
172	m/e 483 (M ⁺ +H)	(d-6-DMSO, d values) 3.05 (m, 4H), 3.65 (m, 4H), 4.00 (s, 3H), 4.00 (s, 3H), 6.45 (dd, 1H), 6.55 (d, 1H), 6.65 (dd, 1H), 7.15 (d, 2H), 7.20 (t, 1H), 7.45 (d, 2H), 7.45 (s, 1H), 8.20 (s, 1H), 8.95 (s, 1H), 11.25 (broad s, 1H)	110°C/5h/1- PrOH	m/e 301 (M ⁺ +H)	KOtBu, DMA	m/e 271 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
173	m/e 569 (M+H) ⁺	(d-6-DMSO, d values) 1.18 (t, 3H), 2.31 (m, 2H), 3.05 (m, 4H), 3.29 (m, 2H), 3.35-3.50 (m, 2H (under H ₂ O signal)), 3.63 (s, 2H), 3.81 (m, 2H), 3.97 (m, 5H), 4.28 (m, 2H), 6.86 (d, 1H), 7.06 (d, 2H), 7.12 (m, 1H), 7.24 (m, 1H), 7.37 (m, 1H), 7.43 (d, 2H), 7.46 (s, 1H), 8.10 (s, 1H), 8.82 (bs, 1H), 10.80 (bs, 1H)	100°C/18h/1 -PrOH				
174	m/e 455 (M ⁺ +H)	(d-6-DMSO, d values) 2.00 (s, 3H), 4.00 (s, 3H), 4.00 (s, 3H), 6.90 (dd, 1H), 7.05 (m, 2H), 7.10 (d, 2H), 7.45 (d, 2H), 7.50 (s, 1H), 7.95 (d, 1H), 8.20 (s, 1H), 8.90 (s, 1H), 9.40 (broad s, 1H), 11.30 (broad s, 1H)	110°C/18h/1 -PrOH.	m/e 273 (M ⁺ +H)	Ac ₂ O, DMA	m/e 243 (M ⁺ +H)	H ₂ , Pd/C, EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
178	m/e 648.5 (M-H) ⁺	(d-6-DMSO, d values) 2.32 (m, 2H), 2.89 (s, 3H), 3.09 (m, 2H), 3.28 (m, 4H), 3.50 (m, 2H), 3.82 (m, 2H), 3.96 (m, 2H), 4.00 (s, 3H), 4.05 (m, 2H), 4.30 (m, 2H), 6.99 (m, 4H), 7.18 (m, 3H), 7.39 (d, 2H), 7.50 (s, 1H), 8.16 (s, 1H), 8.86 (s, 1H)	100°C/18h/1 -PrOH		RT/18h/ MeSO ₂ C 1/ iPr ₂ NEt/ DCM	m/e 323 (M+H) ⁺	RT/18h/H ₂ /5% Pd/C/EtOA c
179	m/e 683 (M+H) ⁺	(d-6-DMSO, d values) 0.95 (t, 6H), 2.32 (m, 2H), 2.74 (s, 3H), 3.01 (q, 4H), 3.08 (m, 2H), 3.26 (m, 2H), 3.33 (t, 2H), 3.47 (m, 2H), 3.79 (m, 2H), 3.95 (m, 2H), 3.99 (s, 3H), 4.10 (t, 2H), 4.29 (m, 2H), 6.95 (m, 3H), 7.03 (m, 1H), 7.18 (m, 2H), 7.39 (d, 2H), 7.51 (s, 1H), 8.18 (s, 1H), 8.92 (s, 1H)	100°C/18h/1 -PrOH	m/e 388 M+H) ⁺	RT/18h/ DEAD/P Ph ₃ / DCM	m/e 358 (M+H) ⁺	RT/18h/5 %Pd/C/H ₂ / EtOAc
180	m/e 626 (M+H) ⁺	(d-6-DMSO, d values) 1.69 (m, 2H), 1.78 (s, 3H), 2.34 (m, 2H), 3.02 (m, 2H), 3.08 (m, 2H), 3.26 (m, 2H), 3.47 (m, 2H), 3.79 (m, 2H), 3.95 (m, 2H), 3.97 (m, 2H), 4.00 (s, 3H), 4.30 (m, 2H), 6.98 (m, 3H), 7.05 (m, 1H), 7.39 (d, 2H), 7.53 (s, 1H), 7.84 (m, 1H), 8.24 (s, 1H), 8.95 (s, 1H)	100°C/18h/1 -PrOH	m/e 331 M+H) ⁺	RT/2h/ acetyl chloride/ iPr ₂ NEt/ DCM	m/e 301 (M+H) ⁺	RT/18h/H ₂ /5%Pd/C/E tOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
181	m/e 654 (M+H) ⁺	(d-6-DMSO, d values) 0.95 (d, 6H), 1.71 (m, 2H), 2.31 (m, 3H), 3.04 (m, 4H), 3.28 (m, 2H), 3.47 (m, 2H), 3.81 (m, 2H), 3.95 (m, 2H), 3.99 (m, 5H), 4.29 (m, 2H), 6.99 (m, 3H), 7.04 (m, 1H), 7.14 (m, 2H), 7.40 (d, 2H), 7.53 (s, 1H), 7.71 (m, 1H), 8.26 (s, 1H), 8.95 (s, 1H)	100°C/18h/1 -PrOH	m/e 359 M+H) ⁺	RT/2h/ iso-butyl chloride/ -Pr ₂ NEt DCM	m/e 329 (M+H) ⁺	RT/18h/H ₂ /5%Pd/C/E tOAc
182	m/e 639 (M+H) ⁺	(d-6-DMSO, d values) 2.31 (m, 2H), 3.06 (m, 2H), 3.12 (m, 2H), 3.26 (m, 4H), 3.47 (m, 2H), 3.80 (m, 2H), 3.95 (m, 2H), 3.99 (s, 3H), 4.04 (t, 2H), 4.30 (m, 2H), 6.97 (m, 3H), 7.08 (m, 1H), 7.18 (d, 2H), 7.38 (d, 2H), 7.50 (s, 1H), 8.17 (s, 1H), 8.87 (s, 1H)	100°C/18h/1 -PrOH	m/e 344 M+H) ⁺	RT/18h/ DEAD/P Ph ₃ / DCM	m/e 314 (M+H) ⁺	RT/18h/H ₂ /5%Pd/C/E tOAc
184	m/e 640.6 (M+H) ⁺	(d-6-DMSO, d values) 0.96 (d, 6H), 2.34 (m, 3H), 3.11 (m, 2H), 3.29 (m, 4H), 3.50 (m, 2H), 3.80 (m, 2H), 3.97 (m, 7H), 4.29 (m, 2H), 6.99 (m, 4H), 7.17 (m, 2H), 7.59 (d, 2H), 7.49 (s, 1H), 7.79 (s, 1H), 8.13 (s, 1H), 8.86 (s, 1H)	100°C/18h/1 -PrOH		RT/18h/ so butyl chloride/ Pr ₂ NEt/ DCM	m/e 315.5 (M+H) ⁺	RT/18h/H ₂ /5% Pd/C/EtOA c

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
185	m/e 601.5 (M+H) ⁺	(d-6-DMSO, d values) 3.09 (m, 2H), 3.67 (s, 3H), 4.97 (m, 8H), 7.00 (m, 4H), 7.14 (m, 2H), 7.40 (m, 3H), 7.50 (m, 1H), 7.72 (d, 2H), 8.05 (s, 1H), 8.88 (s, 1H)	100°C/18h/1 -PrOH		RT/18h/ Methyl imid- azole MeSO ₂ C i ^t Pr ₂ NEt/ DCM	m/e 357.5 (M-H) ⁺	RT/18h/H ₂ /5% Pd/C/EtOA c
186	m/e 535.5 (M+H) ⁺	(d-6-DMSO, d values) 2.89 (s, 3H), 3.26 (m, 2H), 3.97 (m, 6H), 4.05 (m, 2H), 7.00 (m, 4H), 7.17 (m, 3H), 7.41 (m, 3H), 8.09 (s, 1H), 8.89 (s, 1H)	100°C/18h/1 -PrOH				
187	m/e 483 (M ⁺ +H)	(d-6-DMSO, d values) 3.05 (m, 4H), 3.65 (m, 4H), 4.00 (s, 3H), 4.00 (s, 3H), 6.45 (dd, 1H), 6.55 (t, 1H), 6.70 (dd, 1H), 7.15 (d, 2H), 7.20 (t, 1H), 7.45 (d, 2H), 7.45 (s, 1H), 8.20 (s, 1H), 8.90 (s, 1H), 11.30 (broad s, 1H)	110°C/5h/1- PrOH	m/e 301 (M ⁺ +H)	KOtBu, DMA	m/e 271 (M ⁺ +H)	H ₂ , Pd/C, EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
188	m/e 481 (M ⁺ +H)	(d-6-DMSO, d values) 1.40 (broad s, 2H), 1.55 (broad s, 4H), 3.00 (broad s, 4H), 4.00 (s, 3H), 4.00 (s, 3H), 7.00 (m, 4H), 7.20 (m, 2H), 7.40 (d, 2H), 7.45 (s, 1H), 8.20 (s, 1H), 8.85 (s, 1H), 11.10 (broad s, 1H)	110°C/5h/1- PrOH	m/e 299 (M ⁺ +H)	KOtBu, DMA	m/e 269 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
				m/e 285 (M ⁺ +H)	KOtBu, DMA	m/e 255 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
189	m/e 467 (M ⁺ +H)	(d-6-DMSO, d values) 1.80 (m, 4H), 3.25 (m, 4H), 3.95 (s, 6H), 6.75 (t, 1H), 6.90 (m, 4H), 7.05 (t, 1H), 7.40 (d, 2H), 7.45 (s, 1H), 8.20 (s, 1H), 8.85 (s, 1H), 11.15 (broad s, 1H)	110°C/5h/1- PrOH	m/e 525 (M ⁺ +H)	100°C/18h/1 -PrOH		
190	m/e 525 (M ⁺ +H)	(d-6-DMSO, d values) 3.13 (m, 2H), 3.30 (m, 4H), 3.97 (d, 6H), 4.04 (m, 2H), 6.98 (m, 3H), 7.06 (m, 1H), 7.18 (m, 2H), 7.37 (m, 2H), 8.06 (s, 1H), 8.89 (s, 1H)	100°C/18h/1 -PrOH				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
191	m/e 548 (M+H) ⁺	(d-6-DMSO, d values) 1.79 (m, 2H), 2.84 (s, 3H), 2.97 (m, 2H), 3.97 (s, 6H), 4.03 (m, 2H), 6.97 (m, 4H), 7.05 (m, 1H), 7.18 (m, 2H), 7.42 (m, 3H), 8.13 (s, 1H), 8.91 (s, 1H)	100°C/18h/1 -PrOH	m/e 367 M+H) ⁺	RT/2h/ MeSO ₂ - Cl /iPr ₂ NEt/ DCM	m/e 336 (M+H) ⁺	RT/18h/H ₂ /5%Pd/C/E tOAc
192	m/e 541 (M+H) ⁺	(d-6-DMSO, d values) 0.96 (d, 6H), 1.71 (m, 2H), 2.31 (m, 1H), 3.05 (m, 2H), 3.97 (s, 8H), 6.97 (m, 3H), 7.04 (m, 1H), 7.16 (m, 2H), 7.40 (m, 3H), 7.71 (bs, 1H), 8.11 (s, 1H), 8.89 (s, 1H)	100°C/18h/1 -PrOH				
193	m/e 660 (M+H) ⁺	(d-6-DMSO, d values) 1.79 (m, 2H), 2.31 (m, 2H), 2.84 (s, 3H), 2.98 (m, 2H), 3.10 (m, 2H), 3.28 (m, 2H), 3.4-3.6 (m, 2H (under H ₂ O peak)), 3.78 (m, 2H), 3.98 (bs, 5H), 4.02 (m, 2H), 4.28 (m, 2H), 6.97 (m, 4H), 7.05 (m, 1H), 7.16 (m, 2H), 7.37 (d, 2H), 7.46 (s, 1H), 8.10 (s, 1H), 8.84 (s, 1H)	100°C/18h/1 -PrOH				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
195	m/e 630 (M+H) ⁺	(d-6-DMSO, d values) 1.75 (t, 2H), 2.27 (s, 3H), 2.53 (s, 3H), 2.87 (m, 2H), 3.98 (m, 8H), 6.95 (m, 3H), 7.01 (m, 1H), 7.13 (m, 2H), 7.38 (m, 3H), 7.87 (m, 1H), 8.06 (s, 1H), 8.85 (s, 1H)	100°C/18h/1 -PrOH	m/e 448 (M+H) ⁺	RT/18h/ DMSO chloride/ iPr ₂ NEt/ DCM	m/e 418 (M+H) ⁺	80°C/18h/ SnCl ₂ .2H ₂ O/EtOAc
196	m/e 412 (M ⁺ +H)	(d-6-DMSO, d values) 2.30 (s, 3H), 4.00 (s, 6H), 6.80 (d, 1H), 6.80 (s, 1H), 6.95 (d, 1H), 7.15 (d, 2H), 7.25 (t, 1H), 7.45 (d, 2H), 7.45 (s, 1H), 8.15 (s, 1H), 8.90 (s, 1H) 11.10 (broad s, 1H)	110°C/18h/1 -PrOH	m/e 230 (M ⁺ +H)	KOtBu, DMA	m/e 200 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
198	m/e 427	(d-6-DMSO, d values) 2.65 (s, 3H), 4.00 (s, 3H), 4.00 (s, 3H), 6.60 (t, 1H), 6.75 (m, 2H), 7.00 (m, 1H), 7.05 (d, 2H), 7.40 (d, 2H), 7.50 (s, 1H), 8.20 (s, 1H), 8.90 (s, 1H), 11.20 (broad s, 1H)	110°C/18h/1 -PrOH	m/e 243 (M-H)	HCHO, AcOH, BH ₃ .SM e ₂ , THF	m/e 215 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
199	m/e 441 (M ⁺ +H)	(d-6-DMSO, d values) 1.15 (t, 3H), 3.10 (q, 2H), 4.00 (s, 3H), 4.00 (s, 3H), 6.60 (t, 1H), 6.80 (m, 2H), 7.00 (m, 1H), 7.05 (d, 2H), 7.40 (d, 2H), 7.50 (s, 1H), 8.20 (s, 1H), 8.85 (s, 1H), 11.15 (broad s, 1H)	110°C/12h/1 -PrOH	m/e 259 (M ⁺ +H)	BH ₃ . SMe ₂ , THF	m/e 229 (M ⁺ +H)	H ₂ , Pd/C, EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
200	m/e	(d-6-DMSO, δ values) 3.76 (s, 3H), 3.98 (s, 3H), 4.00 (s, 3H), 6.94 - 7.00 (m, 2H), 7.03 - 7.09 (m, 2H), 7.14 (d, 1H), 7.47 (s, 1H), 7.94 (dd, 1H), 8.21 (s, 1H), 8.27 (d, 1H), 8.93 (s, 1H), 11.23 (bs, 1H)	95°C/16h/1- PrOH	m/e	115°C/	m/e	10% Pd
	429.4 (M+H) ⁺			247.2 M+H) ⁺	2h/ K ₂ CO ₃ / DMA	217.2 (M+H) ⁺	on C/EtOAc
201	m/e	(d-6-DMSO, δ values) 3.74 (s, 3H), 3.98 (s, 3H), 4.00 (s, 3H), 6.66 - 6.72 (m, 2H), 6.77 (dd, 1H), 7.19 (d, 1H), 7.31 (t, 1H), 7.48 (s, 1H), 7.98 (dd, 1H), 8.21 (s, 1H), 8.32 (d, 1H), 8.94 (s, 1H), 11.24 (bs, 1H)	95°C/16h/ 1-PrOH	m/e	115°C/	m/e	10% Pd
	429.4 (M+H) ⁺			247.2 M+H) ⁺	2h/ K ₂ CO ₃ / DMA	217.2 (M+H) ⁺	on C/EtOAc
202	m/e	(d-6-DMSO, δ values) 3.68 (s, 3H), 3.98 (s, 3H), 3.99 (s, 3H), 6.98 (m, 1H), 7.09 - 7.16 (m, 3H), 7.21 (m, 1H), 7.48 (s, 1H), 7.92 (dd, 1H), 8.17 - 8.22 (m, 2H), 8.94 (s, 1H), 11.14 (bs, 1H)	95°C/16h/ 1- PrOH		115°C/		10% Pd
	429.4 (M+H) ⁺				2h/ K ₂ CO ₃ / DMA		on C/EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
203		(d-6-DMSO, δ values) 3.67 (s, 3H), 3.99 (s, 3H), 7.00 (t, 1H), 7.12 - 7.29 (m, 3H), 7.42 (s, 1H), 8.16 (s, 1H), 8.77 (s, 2H), 8.95 (s, 1H)	100°C/16h/ 1-PrOH	m/e 247 $M+H)^+$	RT/1h/ KOtBu/ MeO- phenol/ DMA 135°C/ 5h/	m/e 217.9 $(M+H)^+$	RT/4h/5% Pd on C/H ₂ / EtOAc
212	m/e 467.4 $(M+H)^+$	(d-6-DMSO, δ values) 3.99 (s, 3H), 4.00 (s, 3H), 7.32 (d, 1H), 7.44 - 7.49 (m, 2H), 7.57 (d, 1H), 7.68 (t, 1H), 8.03 (dd, 1H), 8.19 (s, 1H), 8.35 (d, 1H), 8.94 (s, 1H)	100°C/7h/1- PrOH				
217	m/e 542 (M^++H)	(d-6-DMSO, δ values) 2.34 (m, 2H), 3.14 (m, 2H), 3.50 (m, 4H), 3.76 (s, 3H), 3.82 (m, 2H), 3.99 (s, 2H), 4.02 (s, 3H), 4.32 (t, 2H), 6.71 (m, 2H), 6.80 (m, 1H), 7.20 (d, 2H), 7.33 (t, 1H), 7.50 (s, 1H), 7.96 (m, 1H), 8.16 (s, 1H), 8.32 (d, 1H), 8.81 (s, 1H), 10.86 (broad, 2H)	1-PrOH / 1.0M ethereal HCl (1 equiv.) / 110deg / 3 h				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
219	m/e 507.4 (M+H) ⁺		RT/15min/ NaH/ DMA RT2h		100°C/ 3h/ K ₂ CO ₃ / DMA	m/e 219.3 (M+H) ⁺	RT/5h/10 %Pd on C/H ₂ / EtOAc
220		(d-6-DMSO, δ values) 3.68 (s, 3H), 4.00 (s, 3H), 6.98 (t, 1H), 7.08 - 7.16 (m, 3H), 7.22 (m, 1H), 7.52 (s, 1H), 7.88 (dd, 1H), 7.96 (s, 1H), 8.17 (dd, 1H), 8.91 (s, 1H), 10.80 (bs, 1H)	100°C/16h/1 -PrOH				
222	m/e 519.4 (M ⁺ +H)		RT/15min/ NaH//DMA then ii) RT2h		100°C/ 3h/ K ₂ CO ₃ / DMA	m/e 230.6 (M ⁺ +H)	RT/5h/10 %Pd on C/H ₂ / EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
226	m/e 528 (M ⁺ +H)	(d-6-DMSO, d values) 3.58 (m, 4H), 3.70 (m, 2H), 3.76 (s, 3H), 3.86 (m, 2H), 4.00 (m, 2H), 4.03 (s, 3H), 4.70 (t, 2H), 6.71 (m, 3H), 6.80 (m, 1H), 7.20 (d, 1H), 7.34 (t, 1H), 7.54 (s, 1H), 7.97 (m, 1H), 8.21 (s, 1H), 8.33 (d, 1H), 8.86 (s, 1H), 10.95 (broad, 1H), 11.28 (broad, 1H)	1-PrOH / 1.0M ethereal HCl (1 equiv.) / 110deg / 6 h				
258	m/e 392 (M ⁺ +H)	(CDCl ₃ , d values) 2.10 (m, 2H), 3.65 (s, 3H), 3.95 (m, 4H), 4.00 (s, 3H), 4.95 (m, 1H), 6.90 (d, 2H), 6.90 (s, 1H), 7.15 (d, 2H), 7.25 (s, 1H), 7.35 (s, 1H), 8.60 (s, 1H)	110°C/5h/1- PrOH		KOtBu, DMA	m/e 180 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
259	m/e 406 (M ⁺ +H)	(d-6-DMSO, d values) 1.60 (m, 2H), 2.00 (m, 2H), 3.50 (m, 2H), 3.85 (m, 2H), 4.00 (s, 6H), 4.65 (m, 1H), 7.05 (d, 2H), 7.35 (d, 2H), 7.50 (s, 1H), 8.20 (s, 1H), 8.90 (s, 1H), 11.20 (broad s, 1H)	110°C/3h/1- PrOH		KOtBu, DMA	m/e 194 (M ⁺ +H)	H ₂ , Pd/C, EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
261	m/e 433, 435 (M+H) ⁺	(d-6-DMSO, d values) 3.98 (d, 6H), 7.2 (m, 2H), 7.28 (m, 2H) 7.42 (m, 3H), 8.10 (m, 3H), 8.95 (s, 1H)	85°C/18h/ DME			m/e 220 (M+H) ⁺	iKF- Al ₂ O ₃ , 18- C-6, DMSO then TFA, Et ₃ SiH
262	m/e 397 (M+H) ⁺	(d-6-DMSO, d values) 3.90 (s, 3H), 3.95 (s, 3H), 6.98 (d, 1H), 7.16 (m, 1H) 7.19 (d, 1H), 7.28 (d, 1H), 7.31 (m, 1H), 7.74 (s, 1H), 7.82 (m, 1H), 8.19 (m, 1H), 8.41 (s, 1H), 9.42 (s, 1H)	100°C/24h/1 -PrOH		m/e 187 (M+H) ⁺	TFA, Et ₃ SiH	
263	m/e 424 (M+H) ⁺	(d-6-DMSO, d values) 3.98 (d, 6H), 7.31 (m, 2H), 7.38 (d, 2H) 7.42 (s, 1H), 7.51 (d, 2H), 8.11 (s, 1H), 8.4 (m, 2H), 8.95 (1H, s).	100°C/18h/1 -PrOH			m/e (M+H) ⁺	TFA, Et ₃ SiH
264	m/e 424 (M+H) ⁺	(d-6-DMSO, d values) 3.98 (d, 6H), 7.32 (m, 2H), 7.41 (s, 1H) 7.50 (m, 2H), 7.61 (d, 1H), 8.12 (s, 1H), 8.42 (d, 1H), 8.96 (s, 1H)	100°C/18h/1 -PrOH			m/e 212 (M+H) ⁺	TFA, Et ₃ SiH

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
265	m/e 415 (M+H) ⁺	(d-6-DMSO, d values) 4.00 (d, 6H), 7.18 (m, 2H), 7.22 (m, 2H) 7.36 (m, 1H), 7.46 (d, 2H), 7.50 (s, 1H), 8.10 (s, 1H), 8.38 (dd, 1H), 8.90 (s, 1H)	100°C/18h/1 -PrOH			m/e (M+H) ⁺	TFA, Et ₃ SiH
266	m/e 400.3 (M+H) ⁺	(d-6-DMSO, δ values) 3.98 (s, 3H), 4.00 (s, 3H), 7.34 (d, 1H), 7.50 (s, 1H), 7.54 (dd, 1H), 7.68 (dd, 1H), 8.02 (dd, 1H), 8.26 (s, 1H), 8.31 (d, 1H), 8.46 (d, 1H), 8.50 (d, 1H), 8.92 (s, 1H)	100°C/7h/1- PrOH				
267	m/e 440 (M+H) ⁺	(d-6-DMSO, d values) 3.99 (ap.d, 6H), 7.08 (d, 1H), 7.42 (s, 1H) 7.52 (d, 2H), 7.70 (d, 2H), 8.00 (m, 2H), 8.80 (m, 1H), 8.90 (s, 1H)	100°C/18h/1 -PrOH				
268	m/e 405 (M+H) ⁺	(d-6-DMSO, d values) 3.99 (s, 6H), 7.22 (d, 1H), 7.32 (d, 1H) 7.46 (m, 3H), 7.52 (d, 2H), 8.15 (s, 1H), 8.95 (s, 1H)	100°C/18h/1 -PrOH				
269	m/e 434, 436 (M+H) ⁺	(d-6-DMSO, d values) 3.98 (ap.d, 6H), 7.40 (m, 3H), 7.53 (d, 2H) 8.12 (s, 1H), 8.20 (d, 1H), 8.25 (d, 1H), 8.96 (s, 1H)	100°C/18h/1 -PrOH		K ₂ CO ₃ , DMA		SnCl ₂ .2H ₂ O, EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
270	m/e 400 (M+H) ⁺	(d-6-DMSO, d values) 4.00 (s, 6H), 7.30 (d, 1H), 7.33 (d, 2H), 7.45 (m, 2H), 7.52 (s, 1H), 8.18 (s, 1H), 8.66 (d, 2H), 8.96 (s, 1H)	100°C/18h/1 -PrOH	m/e 218 M+H ⁺	K ₂ CO ₃ , DMA		10%Pd/C, EtOAc
271	m/e 446 (M+H) ⁺	(d-6-DMSO, d values) 2.40 (s, 3H), 4.00 (s, 6H), 6.78 (d, 1H), 7.40 (bd, 2H), 7.51 (s, 1H), 7.57 (d, 2H), 8.19 (s, 1H), 8.53 (d, 1H), 8.98 (s, 1H)	100°C/18h/1 -PrOH	m/e 264 M+H ⁺	K ₂ CO ₃ , DMA		SnCl ₂ .2H ₂ O, EtOAc
272	m/e 481 (M+H) ⁺	(d-6-DMSO, d values) 3.97 (s, 3H), 5.29 (s, 2H), 7.29 (d, 1H), 7.33 (d, 1H), 7.35 (m, 2H), 7.42 (m, 2H), 7.43-7.54 (m, 6H), 8.41 (s, 1H), 8.95 (s, 1H)	100°C/18h/1 -PrOH			m/e 193 (M+H) ⁺	120°C/18 h/KOH/D MA
287	m/e 446 (M ⁺ +H)	(d-6-DMSO, d values) 3.60 (s, 3H), 3.95 (s, 3H), 4.00 (s, 3H), 6.55 (dd, 1H), 6.95 (td, 1H), 7.00 (d, 1H), 7.05 (d, 1H), 7.10 (td, 1H), 7.15 (td, 1H), 7.45 (s, 1H), 7.60 (dd, 1H), 8.00 (s, 1H), 9.00 (s, 1H), 10.90 (broad s, 1H)	110°C/60h/1 -PrOH	m/e 264 M ⁺ +H)	KOtBu, DMA	m/e 234 (M ⁺ +H)	SnCl ₂ .2H ₂ O, EtOAc
288	m/e 477 (M+H) ⁺	(d-6-DMSO, d values) 1.23 (t, 3H), 4.00 (s, 3H), 4.20 (q, 2H), 5.06 (s, 2H), 7.26 (d, 1H), 7.33 (m, 3H), 7.50 (m, 4H), 8.16 (s, 1H), 8.89 (s, 1H)	100°C/18h/1 -PrOH				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
290	m/e 493 (M ⁺ +H)	(d-6-DMSO, d values) 3.36 (m, 6H), 3.77 (m, 4H), 4.33 (m, 4H), 7.27 (d, 1H), 7.33 (d, 1H), 7.48 (m, 2H), 7.52 (m, 3H), 8.21 (s, 1H), 8.91 (s, 1H), 11.12 (broad, 1H)	EtOH / reflux / 18 h				
294	m/e 511 (M+H) ⁺	(d-6-DMSO, d values) 2.33 (m, 2H), 3.08 (m, 2H), 3.28 (m, 2H), 3.47 (m, 2H), 3.81 (m, 2H), 3.93 (m, 2H), 3.99 (s, 3H), 4.29 (m, 2H), 7.01 (d, 1H), 7.14 (m, 1H), 7.26 (d, 2H), 7.34 (d, 2H), 7.54 (s, 1H), 7.85 (m, 1H), 8.18 (s, 1H), 8.91 (s, 1H)	100°C/18h/1 -PrOH				
295	m/e 421 (M ⁺ +H)	(d-6-DMSO, d values) 3.90 (s, 3H), 3.95 (s, 3H), 7.25 (d, 2H), 7.40 (s, 1H), 7.65 (m, 4H), 7.75 (d, 1H), 8.60 (s, 1H), 9.60 (broad s, 1H)	110°C/18h/1 -PrOH		KOtBu, DMA	m/e 209 (M ⁺ +H)	SnCl ₂ .2H ₂ 0, HCl, MeOH,
296	m/e 434 (M-H)	(d-6-DMSO, d values) 4.00 (s, 3H), 4.00 (s, 3H), 7.35 (d, 2H), 7.40 (d, 2H), 7.50 (s, 1H), 7.55 (s, 1H), 8.25 (s, 1H), 8.95 (s, 1H), 11.40 (broad s, 1H)	110°C/18h/1 -PrOH/HCl			m/e 224 (M ⁺ +H)	SnCl ₂ .2H ₂ 0HCl, MeOH,

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
297	m/e 390 (M ⁺ +H)	(d-6-DMSO, d values) 1.60 (m, 2H), 1.70 (m, 4H), 1.90 (m, 2H), 4.00 (s, 3H), 4.00 (s, 3H), 4.85 (m, 1H), 7.00 (d, 2H), 7.35 (d, 2H), 7.50 (s, 1H), 8.20 (s, 1H), 8.90 (s, 1H), 11.20 (broad s, 1H)	110°C/5h/1-PrOH	m/e 208 (M ⁺ +H)	KOtBu, DMA	m/e 178 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
298	m/e 404 (M ⁺ +H)	(d-6-DMSO, d values) 1.40 (m, 6H), 1.70 (m, 2H), 1.95 (m, 2H), 4.00 (s, 6H), 4.40 (m, 1H), 7.00 (d, 2H), 7.35 (d, 2H), 7.45 (s, 1H), 8.20 (s, 1H), 8.90 (s, 1H), 11.15 (broad s, 1H)	110°C/3h/1-PrOH		KOtBu, DMA	m/e 192 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
299	m/e 500 (M+H) ⁺	(d-6-DMSO, d values) 2.83 (s, 3H), 2.99 (s, 3H), 3.98 (s, 6H), 4.96 (s, 2H), 7.10 (m, 1H), 7.20 (d, 2H), 7.42 (m, 1H), 7.48 (m, 3H), 7.69 (m, 1H), 8.16 (s, 1H), 8.95 (s, 1H)	100°C/18h/1-PrOH				
300	m/e 391 (M+H) ⁺	(d-6-DMSO, d values) 3.90 (s, 3H), 7.21 (d, 2H), 7.30 (m, 3H), 7.37 (m, 2H), 7.69 (s, 1H), 8.40 (s, 1H)	75°C/2h/TF A thioanisole				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
303	m/e 505 (M ⁺ +H)	(d-6-DMSO, d values) 1.40 (s, 9H), 1.55 (m, 2H), 1.90 (m, 2H), 3.2 (m, 2H), 3.65 (m, 2H), 4.00 (s, 3H), 4.00 (s, 3H), 4.60 (m, 1H), 7.05 (d, 2H), 7.35 (d, 2H), 7.50 (s, 1H), 8.20 (s, 1H), 8.85 (s, 1H), 11.25 (broad s, 1H)	110°C/18h/1 -PrOH				
304	m/e 432 (M ⁺ +H)	(d-6-DMSO, d values) 2.45 (s, 3H), 3.85 (s, 3H), 3.95 (s, 3H), 5.15 (s, 2H), 6.95 (s, 1H), 7.20 (s, 4H), 7.30 (s, 1H), 7.35 (s, 1H), 7.65 (s, 1H), 8.45 (s, 1H), 9.40 (broad s, 1H)	110°C/18h/1 -PrOH/HCl	m/e 220 (M ⁺ +H)		SnCl ₂ .2H ₂ 0 HCl, MeOH	
305	m/e 386 (M ⁺ +H)	(d-6-DMSO, d values) 3.85 (s, 3H), 3.95 (s, 3H), 5.20 (s, 2H), 6.90 (s, 1H), 7.15 (s, 1H), 7.20 (d, 2H), 7.25 (s, 1H), 7.30 (s, 1H), 7.35 (s, 1H), 7.70 (d, 2H), 8.45 (s, 1H), 9.40 (broad s, 1H)	110°C/18h/1 -PrOH/HCl	m/e 174 (M ⁺ +H)		SnCl ₂ .2H ₂ 0HCl, MeOH	
306	m/e 454 (M ⁺ +H)	(d-6-DMSO, d values) 4.00 (s, 3H), 4.00 (s, 3H), 5.30 (s, 2H), 7.25 (d, 2H), 7.30 (t, 1H), 7.55 (m, 5H), 8.25 (s, 1H), 8.95 (s, 1H), 11.35 (broad s, 1H)	110°C/18h/1 -PrOH		KOtBu, DMA	m/e 242 (M ⁺ +H)	H ₂ , Pd/C, EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
307	m/e	(d-6-DMSO, d values) 3.95 (s, 3H), 4.00 (s, 3H), 6.35 (d, 1H), 7.40 (d, 2H), 7.50 (s, 1H), 7.55 (d, 2H), 8.25 (s, 1H), 8.80 (d, 1H), 8.95 (s, 1H), 11.40 (broad s, 1H)	90°C/18h/1-PrOH		KOtBu, DMA		SnCl ₂ , 2H ₂ O, HCl, MeOH
	389 (M ⁺ +H)						
308	m/e	(d-6-DMSO, d values) 2.00 (m, 2H), 2.75 (t, 2H), 2.90 (t, 2H), 4.00 (s, 3H), 4.00 (s, 3H), 6.80 (d, 1H), 7.00 (d, 2H), 7.05 (d, 1H), 7.15 (t, 1H), 7.40 (d, 2H), 7.50 (s, 1H), 8.20 (s, 1H), 8.90 (s, 1H), 11.20 (broad s, 1H)	110°C/18h/1-PrOH		KOtBu, DMA	m/e 226 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
	438 (M ⁺ +H)						
309	m/e	(d-6-DMSO, d values) 3.95 (s, 3H), 4.00 (s, 3H), 7.45 (d, 2H), 7.55 (d, 2H), 7.60 (s, 1H), 7.80 (s, 2H), 8.40 (s, 1H), 8.95 (s, 1H), 11.70 (broad s, 1H)	110°C/18h/1-PrOH, HCl	m/e 222 (M ⁺ +H)	KOtBu, DMA	m/e 192 (M ⁺ +H)	SnCl ₂ , 2H ₂ O, HCl, MeOH,
	404 (M ⁺ +H)						

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
310	m/e 611 (M+H) ⁺	(d-6-DMSO, d values) 2.31 (m, 2H), 2.84 (s, 3H), 2.99 (s, 3H), 3.10 (m, 2H), 3.25-3.55 (m, 4H (under H ₂ O signal)), 3.80 (s, 2H), 3.96 (m, 2H), 3.98 (s, 3H), 4.31 (m, 2H), 4.95 (s, 2H), 7.09 (m, 1H), 7.17 (d, 2H), 7.41 (m, 3H), 7.50 (s, 1H), 7.68 (m, 1H), 8.16 (s, 1H), 8.87 (s, 1H).	100°C/18h/1- -PrOH				
311	m/e 468 (M ⁺ +H)	(d-6-DMSO, d values) 1.40 (s, 6H), 3.05 (s, 2H), 3.95 (s, 6H), 6.80 (m, 2H), 7.00 (d, 2H), 7.05 (t, 1H), 7.40 (d, 2H), 7.50 (s, 1H), 8.20 (s, 1H), 8.90 (s, 1H), 11.20 (broad s, 1H)	110°C/18h/1- -PrOH			m/e 256 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
316	m/e 419.4 (M+H) ⁺	(d-6-DMSO, δ values) 1.82 - 1.90 (m, 1H), 2.09 - 2.31 (m, 3H), 3.86 - 4.04 (m, 9H), 7.05 (d, 2H), 7.37 (d, 2H), 7.45 (s, 1H), 7.82 (s, 1H), 8.14 (s, 1H), 8.90 (s, 1H)	100°C/3h/1- PrOH	m/e 237.1 M+H ⁺	RT/18h/ PPh ₃ /DE AD/THF	m/e 207.4 (M+H) ⁺	RT/18h/1 0% Pd on C/EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
317	m/e 419.4 (M+H) ⁺	(d-6-DMSO, δ values) 1.80 - 1.92 (m, 1H), 2.08 - 2.30 (m, 3H), 3.85 - 4.04 (m, 9H), 7.06 (d, 2H), 7.38 (d, 2H), 7.46 (s, 1H), 7.84 (s, 1H), 8.14 (s, 1H), 8.90 (s, 1H)	100°C/3h/ 1-PrOH	m/e 237.1 M+H ⁺	RT/18h/ PPh ₃ / DEAD/ THF	m/e 207.4 (M+H) ⁺	RT/4h/10 % Pd on C/EtOAc
318	m/e 488 (M ⁺ +H)	(d-6-DMSO, δ values) 1.89 (m, 2H), 2.03 (m, 2H), 3.14 (m, 2H), 3.61 (m, 2H), 3.71 (m, 2H), 4.03 (s, 3H), 4.62 (t, 2H), 7.27 (d, 1H), 7.33 (d, 1H), 7.47 (d, 1H), 7.55 (d, 1H), 7.60 (s, 1H), 8.34 (s, 1H), 8.93 (s, 1H), 11.29 (broad, 1H), 11.44 (broad, 1H)	1-PrOH / 1.0M ethereal HCl (1 equiv.) / 105 °C / 20 h				
320	m/e 504 (M ⁺ +H)	(d-6-DMSO, δ values) 3.57 (m, 4H), 3.70 (m, 2H), 3.85 (m, 2H), 4.00 (m, 2H), 4.02 (s, 3H), 4.71 (t, 2H), 7.30 (m, 1H), 7.36 (m, 1H), 7.50 (m, 5H), 8.19 (s, 1H), 8.90 (s, 1H), 10.96 (broad, 1H), 11.38 (broad, 1H)	1-PrOH / 1.0M ethereal HCl (1 equiv.) / 110°/ 6 h				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
400	m/e 569 (M ⁺ +H)	(d-6-DMSO, d values) 3.55 (s, 6H), 3.95 (s, 3H), 4.00 (s, 3H), 6.90 (d, 1H), 7.10 (t, 1H), 7.15 (d, 2H), 7.40 (t, 1H), 7.50 (s, 1H), 7.55 (d, 2H), 7.60 (d, 1H), 8.20 (s, 1H), 8.90 (s, 1H), 11.40 (broad s, 1H)	110°C/3h/1- PrOH.		MsCl, NEt ₃ , CH ₂ Cl ₂	m/e 357 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
401	m/e 528.32 (M ⁺ +H)	(d-6-DMSO, d values) 3.68 (d, 2H), 3.98 (d, 6H), 4.53(s, 2H), 6.94-7.2 (m, 7H), 7.33 (br.s, 1H), 7.4 (s, 1H), 7.42 (d, 2H), 7.95 (br.t, 1H), 8.09 (s, 1H), 8.92(s, 1H), 10.99(br.s, 1H)	100°C/2h/1- PrOH				
402	m/e 556.38 (M ⁺ +H)	(d-6-DMSO, d values) 1.2 (d, 3H), 2.56 (d, 3H), 3.98 (d, 6H), 4.28(m, 1H), 4.52 (s, 2H), 6.96-7.2 (m, 6H), 7.4 (s, 1H), 7.42 (s, 2H), 7.85 (br.d, 1H), 7.92 (br.q, 1H), 8.08 (s, 1H), 8.9(s, 1H), 10.98(br.s, 1H)	100°C/2h/1- PrOH	m/e 374.15 M ⁺ +H)	EDC/D MAP/H OBT/D MA	m/e 344.24 (M ⁺ +H)	Hydrogen/ 5% Pd/C/EtO Ac
403	m/e 542.35 (M ⁺ +H)	(d-6-DMSO, d values) 2.57 (d, 3H), 3.7 (d, 2H), 3.98 (s, 6H), 4.54(s, 2H), 6.94-7.2 (m, 6H), 7.4 (s, 1H), 7.43 (s, 2H), 7.8 (br.q, 1H), 7.92 (br.t, 1H), 8.09 (s, 1H), 8.9(s, 1H), 11.0(br.s, 1H)	100°C/2h/1- PrOH		EDC/D MAP/H OBT/D MA	m/e 330.22 (M ⁺ +H)	Hydrogen/ 5% Pd/C

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
404	m/e 627.49 (M ⁺ +H)	(d-6-DMSO, d values) 1.06 (t, 3H), 1.7 (t, 2H), 3.0 (q, 1H), 3.12 (m, 2H), 3.28 (s, 6H), 3.36 (q, 1H), 3.6 (t, 2H), 3.92 (d, 6H), 5.05(s, 2H), 6.85-7.03 (m, 6H), 7.25 (d, 2H), 7.3 (s, 1H), 7.78 (s, 1H), 8.36 (s, 1H), 8.72 (br.s, 1H) 9.52 (s, 1H)	100°C/2h/1-PrOH	m/e 445.35 M ⁺ +H	EDC/N-Methyl morpho- line/ DCM	m/e 415.32 (M ⁺ +H)	Hydrogen/ 5% Pd/C
405	m/e 582.42 (M ⁺ +H)	(d-6-DMSO, d values) 1.25-1.45 (m, 1H), 1.6-1.8 (m, 5H), 2.74-2.94 (m, 2H), 3.0-3.14 (m, 2H), 3.27-3.56 (m, 4H), 3.97 (d, 6H), 4.55(s, 2H), 6.97-7.2 (m, 6H), 7.42 (d, 2H), 7.48 (s, 1H), 8.08 (t, 1H), 8.22 (s, 1H), 8.95 (s, 1H), 10.13 (br.s, 1H), 11.2 (br.s, 1H)	100°C/2h/1-PrOH	m/e 400.33 M ⁺ +H	EDC/ NMM/ DCM	m/e 370.2 (M ⁺ +H)	Hydrogen/ 5% Pd/C
406	m/e 584.42 (M ⁺ +H)	(d-6-DMSO, d values) 2.96-3.7 (m, 8H), 3.7-3.97 (m, 4H), 3.99 (s, 6H), 4.5(s, 2H), 6.95-7.2 (m, 6H), 7.41 (d, 2H), 7.44 (s, 1H), 8.1 (t, 1H), 8.18 (s, 1H), 8.89 (s, 1H)	100°C/2h/1-PrOH	m/e 402.27 (M ⁺ +H)	EDC/ NMM/ DCM	m/e 372.25 (M ⁺ +H)	Hydrogen/ 5% Pd/C

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
407	m/e 570 (M+H) ⁺	(d-6-DMSO, d values) 0.95 (t, 6H), 2.74 (s, 3H), 3.03 (q, 4H), 3.96 (m, 6H), 4.11 (t, 2H), 6.98 (m, 4H), 7.05 (m, 1H), 7.18 (m, 2H), 7.38 (m, 4H), 8.06 (bs, 1H), 8.87 (bs, 1H)	100°C/18h/1 -PrOH				
409	m/e 513 (M+H) ⁺	(d-6-DMSO, d values) 1.68 (m, 2H), 1.76 (s, 3H), 3.00 (m, 2H), 3.97 (s, 8H), 6.99 (m, 3H), 7.05 (m, 1H), 7.16 (m, 2H), 7.42 (m, 3H), 7.83 (bs, 1H), 8.14 (s, 1H), 8.96 (s, 1H)	100°C/18h/1 -PrOH				
410	m/e 483 (M+H) ⁺	(d-6-DMSO, d values) 2.34 (t, 2H), 2.53 (m, 3H), 2.80 (t, 2H), 3.96 (m, 6H), 6.85 (d, 1H), 7.05 (m, 3H), 7.19 (m, 1H), 7.31 (d, 1H), 7.39 (s, 1H), 7.45 (d, 2H), 7.68 (bs, 1H), 8.05 (s, 1H), 8.89 (s, 1H)	100°C/18h/1 -PrOH	m/e 301 M+H ⁺	RT/18h/ methyla mine.HC I/EDC/ DMAP/ NMM/ DCM	m/e 271 (M+H) ⁺	RT/18h/5 %PdC/H ₂ / EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
411	m/e 547 (M+H) ⁺	(d-6-DMSO, d values) 2.60 (t, 2H), 2.85 (t, 3H), 3.18 (s, 3H), 3.98 (s, 6H), 6.86 (d, 1H), 7.08 (m, 3H), 7.20 (m, 1H), 7.31 (m, 1H), 7.45 (m, 3H), 8.18 (s, 1H), 8' (s, 1H)	100°C/18h/1 -PrOH	m/e 363 (M-H) ⁺	RT/18h/ methane sulphon- amide/ EDC/ DMAP/ NMM/ DCM	m/e 335 (M+H) ⁺	RT/18h/5 %PdC/H ₂ /EtOAc
412	m/e 539 (M+H) ⁺	(d-6-DMSO, d values) 2.58 (m, 2H), 2.83 (m, 2H), 3.47 (m, 4H), 3.95 (m, 6H), 6.88 (d, 1H), 7.08 (d, 2H), 7.11 (m, 1H), 7.20 (m, 1H), 7.35 (m, 2H), 7.43 (d, 2H), 8.02 (s, 1H), 8.94 (s, 1H)	100°C/18h/1 -PrOH	m/e 357 M+H) ⁺	RT/18h/ morpholi ne/EDC/ DMAP/ NMM/D CM	m/e 327 (M+H) ⁺	RT/18h/5 %PdC/H ₂ /EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
413	m/e 509 (M+H) ⁺	(d-6-DMSO, d values) 2.43 (t, 2H), 2.81 (t, 2H), 3.66 (m, 4H), 3.99 (s, 6H), 5.00 (m, 2H), 5.74 (m, 1H), 6.89 (d, 1H), 7.08 (m, 3H), 7.19 (m, 1H), 7.31 (m, 1H), 7.47 (m, 3H), 7.92 (bs, 1H), 8.13 (s, 1H), 8.92 (s, 1H)	100°C/18h/1 -PrOH	m/e 327 M+H) ⁺	RT/18h/ allyl amine EDC/D MAP/ NMM/ DCM	m/e 297 (M+H) ⁺	80°C/18h /SnCl ₂ .2 H ₂ O/EtO Ac
414	m/e 509 (M+H) ⁺	(d-6-DMSO, d values) 3.97 (s, 6H), 4.37 (m, 2H), 4.65 (m, 2H), 6.93 (d, 2H), 7.00 (m, 1H), 7.06 (m, 1H), 7.14 (m, 2H), 7.41 (d, 2H), 7.46 (s, 1H), 7.67 (s, 1H), 7.87 (s, 1H), 8.17 (bs, 1H), 8.91 (s, 1H)	100°C/18h/1 -PrOH	m/e 327 M+H) ⁺	RT/18h/ DEAD/ PPh ₃ / DCM	m/e 297 (M+H) ⁺	RT/18h/5 %Pd/C/H ₂ / EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
415	m/e 510 (M+H) ⁺	(d-6-DMSO, d values) 1.81 (m, 2H), 1.91 (m, 2H), 2.95 (m, 2H), 3.97 (m, 6H), 4.35 (m, 2H), 6.97 (d, 2H), 7.05 (m, 1H), 7.10 (m, 1H), 7.24 (m, 2H), 7.40 (d, 2H), 7.47 (s, 1H), 8.24 (s, 1H), 8.87 (s, 1H)	100°C/18h/1 -PrOH	m/e 329 M+H) ⁺	RT/18h/ b-OH- ethylpyrr olidine/ DEAD/ PPh ₃ / DCM	m/e 299 (M+H) ⁺	RT/18h/5 %Pd/C/ H ₂ / EtOAc
416	m/e 475 (M+H) ⁺		80°C/18h/D ME				
417	m/e 509 (M+H) ⁺	(d-6-DMSO, d values) 3.98 (s, 6H), 4.31 (m, 2H), 4.42 (m, 2H), 6.95 (d, 2H), 7.00 (m, 1H), 7.04 (m, 2H), 7.14 (m, 2H), 7.40 (m, 4H), 7.95 (s, 1H), 8.11 (s, 1H), 8.28 (s, 1H), 8.89 (s, 1H)	100°C/18h/1 -PrOH	m/e 327 M+H) ⁺	RT/18h/ DEAD/ PPh ₃ / DCM	m/e 297 (M+H) ⁺	RT/18h/ 5%Pd/C/ H ₂ / EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
418	m/e 524 (M ⁺ +H)	(d-6-DMSO, d values) 1.12 (t, 3H), 1.88 (m, 2H), 2.04 (m, 2H), 2.97 (q, 2H), 3.16 (m, 2H), 3.68 (m, 4H), 3.95 (s, 3H), 4.54 (t, 2H), 5.66 (broad, 1H), 6.14 (q, 1H), 6.21 (t, 1H), 6.33 (q, 1H), 7.05 (m, 3H), 7.30 (d, 2H), 7.43 (s, 1H), 7.89 (s, 1H), 8.48 (s, 1H), 9.73 (broad, 1H), 10.33 (broad, 1H)	1-PrOH / 1.0M ethereal HCl (1 equiv.) / 105 °C / 20 h				
419	m/e 499 (M ⁺ +H)	(d-6-DMSO, d values) 1.90 (m, 2H), 2.04 (m, 2H), 3.15 (m, 2H), 3.62 (m, 2H), 3.71 (m, 2H), 3.99 (s, 3H), 4.59 (t, 2H), 7.17 (m, 5H), 7.44 (m, 3H), 7.52 (s, 1H), 8.16 (s, 1H), 8.86 (s, 1H), 10.91 (broad, 2H)	1-PrOH / 1.0M ethereal HCl (1 equiv.) 105 °C / 20h				
420	m/e 506 (M ⁺ +H)	(d-6-DMSO, d values) 1.89 (m, 2H), 2.04 (m, 2H), 3.17 (m, 2H), 3.64 (m, 2H), 3.71 (m, 2H), 4.01 (s, 3H), 4.59 (t, 2H), 6.96 (d, 2H), 7.31 (m, 3H), 7.52 (m, 3H), 7.64 (m, 1H), 7.91 (m, 1H), 8.13 (s, 1H), 8.82 (s, 1H), 10.74 (broad, 2H)	1-PrOH / 1.0M ethereal HCl (1 equiv.) / 105 °C / 20h				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
421	m/e 527 (M ⁺ +H)	(d-6-DMSO, d values) 3.55 (m, 4H), 3.70 (m, 2H), 3.76 (s, 3H), 3.85 (m, 2H), 4.00 (m, 2H), 4.01 (s, 3H), 4.70 (t, 2H), 6.99 (m, 2H), 7.07 (d, 1H), 7.21 (m, 2H), 7.40 (d, 2H), 7.54 (s, 1H), 8.18 (s, 1H), 8.88 (s, 1H), 10.94 (broad, 1H), 11.41 (broad, 1H)	1-PrOH / 1.0M ethereal HCl (1 equiv.) / 110deg / 3 h				
422	m/e 511 (M ⁺ +H)	(d-6-DMSO, d values) 1.89 (m, 2H), 2.04 (m, 2H), 3.15 (m, 2H), 3.63 (m, 4H), 3.71 (m, 2H), 3.74 (s, 3H), 3.99 (s, 3H), 4.59 (t, 2H), 6.97 (m, 3H), 7.05 (m, 1H), 7.19 (m, 2H), 7.37 (d, 2H), 7.50 (s, 1H), 8.13 (s, 1H), 8.83 (s, 1H), 10.89 (broad, 1H)	1-PrOH / 1.0M ethereal HCl (1 equiv.) / 105 °C / 20h				
423	m/e 568(M ⁺ +H)		100°C/18h/ N-PrOH				
424	m/e 504 (M ⁺ +H)		100°C/18h/ 1-PrOH				
425	m/e 456 (M ⁺ +H)		100°C/18h/ 1-PrOH				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
427	m/e 471 (M ⁺ +H)		100°C/18h/ 1-PrOH	m/e 303 M+H) ⁺	150°C/2. 5h/ DMA/ KO ^t Bu	m/e 273 (M+H) ⁺	RT/18/ H ₂ /10% Pd/C/ EtOAc
428	m/e 481.4 (M+H) ⁺	(d-6-DMSO, δ values) 0.47 (m, 2H), 0.61 (m, 2H), 2.69 (m, 1H), 3.98 (s, 3H), 4.40 (s, 2H), 6.59 - 6.65 (m, 2H), 6.71 (d, 1H), 7.15 (d, 2H), 7.28 (t, 1H), 7.43 (s, 1H), 7.49 (d, 3H), 8.08 (m, 1H), 8.70 (d, 1H), 8.99 (s, 1H)	100°C/5h/ 1-PrOH				
429	m/e 398.3 (M+H) ⁺	(d-6-DMSO, δ values) 3.74 (s, 3H), 3.98 (s, 3H), 6.92 - 6.98 (m, 3H), 7.06 (d, 1H), 7.16 - 7.26 (m, 2H), 7.36 - 7.44 (m, 3H), 7.47 (d, 1H), 8.70 (d, 1H), 9.00 (s, 1H), 11.28 (s, 1H)	100°C/3h/ 1-PrOH				Rev. Chim. (1988), 39 (6), 477-82

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
431	m/e 512 (M ⁺ +H)	(d-6-DMSO, d values) 1.89 (m, 2H), 2.03 (m, 2H), 3.13 (m, 2H), 3.63 (m, 2H), 3.71 (m, 2H), 3.73 (s, 3H), 4.04 (s, 3H), 4.60 (m, 2H), 6.68 (m, 2H), 6.77 (d, 1H), 7.17 (d, 1H), 7.30 (t, 1H), 7.57 (s, 1H), 7.96 (m, 1H), 8.31 (d, 1H), 8.39 (s, 1H), 8.91 (s, 1H), 11.22 (broad, 1H), 11.47 (broad, 1H)	1-PrOH / 1.0M ethereal HCl (1 equiv.) / 105 °C / 20 h				
432	m/e 554 (M ⁺ +H)	(d-6-DMSO, d values) 2.95 (t, 2H), 3.05 (m, 2H), 3.15 (m, 4H), 3.80 (m, 2H), 3.90 (m, 2H), 3.95 (s, 3H), 4.00 (s, 3H), 6.80 (m, 1H), 7.10 (d, 4H), 7.45 (d, 2H), 7.50 (s, 1H), 7.85 (m, 1H), 8.30 (s, 1H), 8.90 (s, 1H), 9.80 (broad s, 1H), 11.20 (broad s, 1H), 11.40 (broad s, 1H)	110°C/18h/ 1-PrOH/ HCl	m/e 342 (M ⁺ +H)		H ₂ , Pd/C, EtOAc	
433	m/e 582 (M ⁺ +H)	(d-6-DMSO, d values) 1.10 (s, 3H), 1.15 (s, 3H), 2.60 (m, 2H), 2.95 (t, 2H), 3.35 (m, 4H), 4.00 (s, 3H), 4.00 (s, 3H), 6.90 (m, 1H), 7.10 (d, 4H), 7.45 (d, 2H), 7.55 (s, 1H), 7.90 (m, 1H), 8.35 (s, 1H), 8.90 (s, 1H), 9.80 (broad s, 1H), 11.45 (broad s, 2H)	110°C/18h/ 1-PrOH/ HCl	m/e 370 (M ⁺ +H)		H ₂ , Pd/C, EtOAc	

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
434	m/e 552 (M ⁺ +H)	(d-6-DMSO, d values) 1.35 (m, 1H), 1.70 (m, 5H), 2.90 (m, 4H), 3.20 (m, 2H), 3.30 (m, 2H), 4.00 (s, 3H), 4.00 (s, 3H), 6.90 (m, 1H), 7.10 (m, 4H), 7.45 (d, 2H), 7.55 (s, 1H), 7.85 (m, 1H), 8.30 (s, 1H), 8.90 (s, 1H), 9.80 (broad s, 1H), 10.35 (broad s, 1H), 11.40 (broad s, 1H)	110°C/2h/1- PrOH/HCl			m/e 340	H ₂ , Pd/C, EtOAc
435	m/e 498 (M ⁺ +H)	<u>NMR Spectrum</u> (d-6-DMSO@373K, d values) 2.55 (s, 3H), 3.10 (m, 2H), 3.70 (m, 2H), 4.00 (s, 3H), 4.00 (s, 3H), 6.95 (m, 1H), 7.05 (d, 2H), 7.10 (m, 2H), 7.40 (d, 2H), 7.55 (s, 1H), 7.85 (m, 1H), 8.25 (s, 1H), 8.65 (s, 1H), 8.90 (broad s, 1H), 9.45 (broad s, 1H)	110°C/2h/1- PrOH/HCl			m/e 286 (M ⁺ +H)	H ₂ , Pd/C, EtOAc
436	m/e 512 (M ⁺ +H)	(d-6-DMSO@373K, d values) 2.75 (s, 6H), 2.90 (t, 2H), 3.30 (t, 2H), 4.00 (s, 3H), 4.00 (s, 3H), 6.95 (m, 1H), 7.05 (d, 2H), 7.10 (m, 2H), 7.40 (d, 2H), 7.55 (s, 1H), 7.85 (m, 1H), 8.20 (s, 1H), 8.65 (s, 1H), 9.50 (broad s, 1H)	110°C/2h/1- PrOH/HCl			m/e 300 (M ⁺ +H)	H ₂ , Pd/C, EtOAc

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
437	m/e 497 (M+H) ⁺	(d-6-DMSO, d values) 0.69 (m, 2H), 0.87 (m, 2H), 2.71 (m, 1H), 3.28 (s, 2H), 3.96 (m, 6H), 7.02 (m, 4H), 7.21 (m, 2H), 7.40 (d, 2H), 7.47 (s, 1H), 8.21 (s, 1H), 8.87 (s, 1H), 9.35 (bs, 2H)	100°C/18h/1 -PrOH				
438	m/e 509 (M+H) ⁺	(d-6-DMSO, d values) 0.38 (m, 2H), 0.59 (m, 2H), 2.53 (m, 1H), 3.54 (s, 2H), 3.97 (s, 6H), 6.18 (m, 1H), 6.26 (m, 1H), 6.33 (m, 1H), 7.08 (m, 3H), 7.45 (m, 3H), 7.95 (m, 1H), 8.18 (s, 1H), 8.95 (s, 1H)	100°C/18h/1 -PrOH				
439	m/e 484 (M+H) ⁺	(d-6-DMSO, d values) 2.58 (d, 3H), 3.57 (s, 2H), 3.96 (s, 6H), 6.20 (m, 1H), 6.23 (m, 1H), 6.31 (m, 1H), 7.08 (m, 3H), 7.43 (m, 3H), 7.79 (m, 1H), 8.08 (s, 1H), 8.87 (s, 1H)	100°C/18h/1 -PrOH				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
440	m/e 582.54 (M ⁺ +H)	(d-6-DMSO, d values) 1.56-1.74 (m, 2H), 2.00 (m, 2H), 2.12(m, 1H), 2.64 (d, 3H), 2.72 (d, 3H), 2.96 (m, 2H), 3.44 (m, 2H), 4.0 (s, 3H), 4.06 (d, 2H), 4.40(s, 2H), 6.60(m, 2H), 6.73 (d, 1H), 7.15 (d, 2H), 7.28 (t, 1H), 7.49 (d, 2H), 7.54 (s, 1H), 8.0(br.s, 1H), 8.2(s, 1H), 8.89 (s, 1H), 10.17 (br.s, 1H), 11.16 (br.s, 1H)	100°C/2.5h/ 1-PrOH/ ethereal HCl				
441	m/e 629.52 (M ⁺ +H)	(d-6-DMSO, d values) 1.99 (m, 1H), 2.01 (m, 1H), 2.35(t, 2H), 3.54 (s, 3H), 3.6 (s, 3H), 3.96 (2s, 6H), 4.35 (m, 1H), 4.55 (m, 2H), 6.95-7.21 (m, 6H), 7.4(s, 1H), 7.42(s, 2H), 8.08 (s, 1H), 8.28 (d, 1H), 8.9 (s, 1H), 10.96 (br.s, 1H)	100°C/2h/1- PrOH			m/e 417.26 (M ⁺ +H)	Hydroge n/ 5% Pd/C
442	m/e 571.47 (M ⁺ +H)	(d-6-DMSO, d values) 1.13(t, 2H), 2.45 (t, 2H), 3.32 (t, 2H), 3.96 (2s, 6H), 4.0 (q, 2H), 4.46 (s, 2H), 6.96-7.20 (m, 6H), 7.4(s, 2H), 7.42(s, 1H), 7.75 (t, 1H), 8.06 (s, 1H), 8.89 (s, 1H)	100°C/2h/1- PrOH			m/e 359.22(M ⁺ +H)	Hydroge n/5% Pd/C

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
443	m/e 543.42 (M ⁺ +H)	(d-6-DMSO, d values) 3.60 (s, 3H), 3.90 (d, 2H), 3.96 (2s, 6H), 4.55 (s, 2H), 6.96-7.2 (m, 6H), 7.4 (s, 1H), 7.42 (s, 2H), 8.05 (s, 1H), 8.16 (t, 1H), 8.9 (s, 1H), 10.99 (br.s, 1H)	100°C/2h/1- PrOH			m/e 331.14(M ⁺ +H)	Hydroge n/5% Pd/C
444	m/e 629.52 (M ⁺ +H)	(d-6-DMSO, d values) 1.70 (m, 1H), 1.86 (m, 1H), 2.0 (t, 2H), 2.45 (d, 3H), 2.56 (d, 3H), 3.96 (2s, 6H), 4.2 (m, 1H), 4.52 (s, 2H), 6.94-7.21 (m, 6H), 7.39 (s, 1H), 7.41 (s, 2H), 7.7 (q, 1H), 7.81 (d, 2H), 7.92 (q, 1H), 8.08 (s, 1H), 8.9 (s, 1H), 10.92 (br.s, 1H)	100°C/2h/1- PrOH				
445	m/e 556.45 (M ⁺ +H)	(d-6-DMSO, d values) 2.23 (t, 2H), 2.5 (d, 3H), 3.29 (t, 2H), 3.97 (2s, 6H), 4.45 (s, 2H), 6.96-7.2 (m, 6H), 7.41 (s, 1H), 7.44 (s, 2H), 7.62 (t, 1H), 7.8 (q, 1H), 8.13 (s, 1H), 8.9 (s, 1H), 11.03 (br.s, 1H)	100°C/2h/1- PrOH				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
446	m/e 568.45 (M ⁺ +H)	(d-6-DMSO, d values) 0.4 (m, 2H), 0.56 (m, 2H), 2.47 (m, 1H), 3.66 (d, 2H), 3.98 (d, 6H), 4.54(s, 2H), 6.94-7.2 (m, 6H), 7.4 (s, 1H), 7.42 (s, 2H), 7.85 (br.t, 1H), 7.95 (d, 1H), 8.10 (s, 1H), 8.88 (s, 1H), 11.09(br.s, 1H)	100°C/2h/1- PrOH				
447		(d-6-DMSO, d values) 0.4 (m, 2H), 0.56 (m, 2H), 1.24 (d, 3H), 2.47 (m, 1H), 3.98 (2s, 6H), 4.23 (m, 1H), 4.54(s, 2H), 6.94-7.2 (m, 6H), 7.4 (s, 1H), 7.42 (s, 2H), 7.90 (d, 1H), 8.03 (d, 1H), 8.10 (s, 1H), 8.88 (s, 1H), 10.94(br.s, 1H)	100°C/2h/1- PrOH				
471	m/e 598.5 (M+H) ⁺	(d-6-DMSO D4 Acetic, δ values) 2.24 - 2.35 (m, 2H), 2.62 (s, 3H), 3.03 - 3.10 (m, 4H), 3.29 (t, 2H), 3.73 - 3.78 (m, 4H), 3.98 (s, 3H), 4.28 (t, 2H), 4.41 (s, 2H), 6.59 - 6.65 (m, 2H), 6.73 (dd, 1H), 7.15 (d, 2H), 7.28 (t, 1H), 7.46 (s, 2H), 7.49 (s, 1H), 8.08 (s, 1H), 8.84 (s, 1H)	RT/48h/Nal/ Morpholine				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
472	m/e 538.5 (M+H) ⁺	(d-6-DMSO, d values) 2.42 (m, 2H), 2.58 (m, 2H), 3.34 (m, 2H), 3.97 (m, 8H), 6.99 (m, 4H), 7.17 (m, 2H), 7.38 (m, 2H), 7.41 (s, 1H), 8.08 (s, 1H), 8.11 (m, 1H), 8.87 (s, 1H)	100°C/18h/ 1-PrOH				
473	m/e 527.5 (M+H) ⁺	(d-6-DMSO, d values) 0.97 (d, 6H), 2.34 (m, 1H), 3.30 (m, 2H), 3.97 (m, 8H), 7.00 (m, 4H), 7.30 (m, 2H), 7.41 (m, 3H), 7.78 (m, 1H), 8.13 (s, 1H), 8.96 (s, 1H)	100°C/18h/1 -PrOH				
474	m/e 499.5 (M+H) ⁺	(d-6-DMSO, d values) 1.79 (s, 3H), 3.29 (m, 2H), 3.96 (m, 8H), 6.99 (m, 4H), 7.17 (m, 2H), 7.41 (m, 3H), 7.89 (m, 1H), 8.12 (s, 1H), 8.92 (s, 1H)	100°C/18h/1 -PrOH				
475	m/e 541.5 (M+H) ⁺	(d-6-DMSO, d values) 3.26 (m, 2H), 4.97 (m, 8H), 4.45 (m, 2H), 5.17 (m, 2H), 5.87 (m, 1H), 7.00 (m, 4H), 7.18 (m, 3H), 7.60 (m, 3H), 8.08 (s, 1H), 8.89 (s, 1H)	100°C/18h/1 -PrOH				

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
476	m/e 610.7 (M-H ⁺)	(d-6-DMSO, d values) 2.32 (m, 2H), 2.82 (s, 3H), 2.93 (s, 3H), 3.10 (m, 2H), 3.22-3.53 (m, 4H, under H ₂ O peak), 3.78 (m, 2H), 3.95 (m, 5H), 4.29 (m, 2H), 4.81 (s, 2H), 7.04 (m, 7H), 7.36 (m, 2H), 7.43 (s, 1H), 8.07 (s, 1H), 8.81 (s, 1H)	RT/18h/ HNMe ₂ .HCl/ DMAP/EDC /NMM/DCM				
477		(d-6-DMSO, δ values) 2.64 (d, 3H), 3.99 (s, 6H), 4.42 (s, 2H), 6.60 - 6.67 (m, 2H), 6.74 (dd, 1H), 7.17 (d, 2H), 7.28 (t, 1H), 7.45 - 7.53 (m, 3H), 7.99 (m, 1H), 8.16 (s, 1H), 8.92 (s, 1H), 11.14 (bs, 1H)	100°C/2h/ 1-PrOH			m/e 273.2 (M+H) ⁺	5% Pd on C/H ₂ / EtOAc
478	m/e 515 (M ⁺ +H)	(d-6-DMSO, d values) 3.56 (m, 4H), 3.70 (m, 2H), 3.86 (m, 2H), 4.00 (m, 2H), 4.02 (s, 3H), 4.71 (t, 2H), 7.16 (d, 2H), 7.25 (m, 3H), 7.43 (m, 1H), 7.48 (d, 2H), 7.57 (s, 1H), 8.23 (s, 1H), 8.91 (s, 1H), 11.10 (broad, 1H), 11.52 (broad, 1H)	1-PrOH / 1.0M ethereal HCl (1 equiv.) / 110deg / 6h				

120

No.	mass spec	n.m.r.	reaction conditions	Intermediate 1		Intermediate 2	
				Mass	Reaction	Mass	Reaction
479	m/e 522 (M ⁺ +H)	(d-6-DMSO, d values) 3.57 (m, 4H), 3.71 (m, 2H), 3.85 (m, 2H), 4.00 (m, 2H), 4.04 (s, 3H), 4.71 (t, 2H), 6.99 (d, 2H), 7.32 (m, 3H), 7.57 (m, 3H), 7.67 (m, 1H), 7.93 (m, 1H), 8.23 (s, 1H), 8.91 (s, 1H), 11.11 (broad, 1H), 11.45 (broad, 1H)	1-PrOH / 1.0M ethereal HCl (1 equiv.) / 110deg / 6h				
480	m/e 540 (M ⁺ +H)	(d-6-DMSO, d values) 1.66 (t, 3H), 3.06 (q, 2H), 3.56 (m, 4H), 3.71 (m, 2H), 3.87 (m, 2H), 4.00 (m, 2H), 4.03 (s, 3H), 4.71 (t, 2H), 6.44 (m, 3H), 7.16 (m, 3H), 7.48 (d, 2H), 7.57 (s, 1H), 8.28 (s, 1H), 8.94 (s, 1H), 11.24 (broad, 1H), 11.55 (broad, 1H)	1-PrOH / 1.0M ethereal HCl (1 equiv.) / 110deg / 6 h				
482	m/e 513.5 (M+H) ⁺	(d-6-DMSO, d values) 1.05 (d, 6H), 3.87 (m, 1H), 3.97 (m, 6H), 4.43 (s, 2H), 7.05 (m, 6H), 7.42 (m, 4H), 8.08 (s, 1H), 8.89 (s, 1H)	100°C/18h/1 -PrOH			m/e 301.5 (M+H)	RT/18h/ H ₂ /5% Pd/C/ EtOAc

In the above and other Examples, the following abbreviations have been used:

- ^1H NMR data is quoted and is in the form of delta values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as an internal standard;
 - nitrogen atoms which are shown as less than trivalent are H substituted to complete the
- 5 trivalency;
- the following abbreviations are used:

	DMSO	dimethyl sulphoxide;
	DMF	<i>N,N</i> -dimethylformamide;
	DCM	dichloromethane;
10	EtOAc	ethyl acetate;
	HOBT	N-hydroxybenzotriazole hydrate ;
	NMM	<i>N</i> -Methylmorpholine;
	TFA	Trifluoroacetic acid;
	1-Pr-OH	propan-1-ol;
15	MeOH	methanol;
	EtOH	ethanol;
	KOtBu	potassium tert-butoxide;
	RT	room/ambient temperature.

Example 6


- 20 Compounds of formula (I) were also converted to different such compounds by reacting appropriate derivatisation reactions, either directly or by way of certain chloro substituted intermediates. These can be summarised in the following Table 8 with the Intermediates listed in the Intermediate Table 9 below.

Table 8

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
I8	morpholine	RT/2hrs	14	m/e 541 (M+H) ⁺	(d-6-DMSO, d values) 2.30 (m, 2H), 3.18 (m, 2H), 3.40 (m, 4H), 3.75 (s, 3H), 3.81 (m, 2H), 3.95 (m, 2H), 3.98 (s, 3H), 4.30 (m, 2H), 6.94 (m, 3H), 7.08 (m, 1H), 7.19 (m, 2H), 7.38 (d, 2H), 7.47 (s, 1H), 8.25 (s, 1H), 8.86 (s, 1H).
I18	N-methyl piperazine	EtOH / 80deg / 3.5 hours	16	m/e 554 (M ⁺ +H)	(d-6-DMSO, d values) 2.31 (m, 2H), 2.83 (s, 3H), 3.30 (m, 2H), 3.54 (broad, 8H), 3.73 (s, 3H), 3.99 (s, 3H), 4.31 (m, 2H), 6.95 (m, 3H), 7.04 (d, 1H), 7.17 (m, 2H), 7.39 (d, 2H), 7.48 (s, 1H), 8.15 (s, 1H), 8.88 (s, 1H), 11.12 (broad, 1H).
I9	N-methyl piperazine	RT/18hrs/NaI	17	m/e 554 (M+H) ⁺	(d-6-DMSO, d values) 2.25 (m, 2H), 2.80 (m, 3H), 3.38 (m, 2H), 3.60 (m, 8H), 3.78 (s, 3H), 3.99 (s, 3H), 4.35 (m, 2H), 6.96 (m, 3H), 7.08 (m, 1H), 7.19 (m, 2H), 7.39 (d, 2H), 7.50 (s, 1H), 8.25 (bs, 1H), 8.91 (s, 1H).
I9	pyrrolidine	RT/18hrs/NaI	18	m/e 525 (M+H) ⁺	(d-6-DMSO, d values) 1.88 (m, 2H), 2.04 (m, 2H), 2.26 (m, 2H), 3.32 (m, 2H), 3.60 (m, 4H), 3.75 (s, 3H), 4.00 (s, 3H), 4.28 (m, 2H), 6.95 (m, 3H), 7.05 (m, 1H), 7.10 (m, 2H), 7.38 (m, 3H), 8.15 (s, 1H), 8.60 (bs, 1H), 8.93 (s, 1H).


Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
I9	piperidine	RT/18hrs/NaI	19	m/e 539 (M+H) ⁺	(d-6-DMSO, d values) 1.40 (m, 2H), 1.6-1.8 (m, 4H), 2.28 (m, 2H), 2.95 (m, 2H), 3.21 (m, 2H), 3.45 (m, 2H), 3.72 (s, 3H), 3.97 (s, 3H), 4.28 (m, 2H), 6.94 (m, 3H), 7.07 (m, 1H), 7.20 (m, 2H), 7.39 (d, 2H), 7.45 (s, 1H), 8.24 (s, 1H), 8.92 (s, 1H).
I9	dimethyl-amine	RT/18hrs/NaI/ EtOH	20	m/e 499 (M+H) ⁺	(d-6-DMSO, d values) 2.23 (m, 2H), 2.81 (d, 6H), 3.24 (m, 2H), 3.73 (s, 3H), 3.99 (s, 3H), 4.29 (m, 2H), 6.95 (m, 3H), 7.06 (m, 1H), 7.18 (m, 2H), 7.37 (d, 2H), 7.39 (s, 1H), 8.13 (s, 1H), 8.85 (s, 1H).
I10	morpholine	RT/18hrs/NaI	21	m/e 527 (M+H) ⁺	(d-6-DMSO, d values) 3.06 (m, 2H), 3.39 (m, 2H), 3.64 (m, 2H), 3.71 (s, 3H), 3.75 (m, 2H), 3.90 (m, 2H), 4.00 (s, 3H), 4.68 (m, 2H), 6.94 (m, 3H), 7.05 (m, 1H), 7.19 (m, 2H), 7.37 (d, 2H), 7.50 (s, 1H), 8.38 (s, 1H), 8.87 (s, 1H).
I10	N-methyl piperazine	RT/18hrs/NaI	22	m/e 540 (M+H) ⁺	(d-6-DMSO, d values) 2.80 (s, 3H), 3.24-3.65 (m, 10H), 3.72 (s, 3H), 3.99 (s, 3H), 4.58 (m, 2H), 6.95 (m, 3H), 7.06 (m, 1H), 7.19 (m, 2H), 7.39 (d, 2H), 7.50 (s, 1H), 8.36 (s, 1H), 8.85 (s, 1H).
I10	pyrrolidine	RT/18hrs/NaI	23	m/e 511 (M+H) ⁺	(d-6-DMSO, d values) 1.84 (m, 2H), 2.04 (m, 2H), 3.05 (m, 2H), 3.65-3.72 (m, 4H), 3.75 (s, 3H), 3.98 (s, 3H), 4.60 (m, 2H), 6.96 (m, 3H), 7.07 (m, 1H), 7.18 (m, 2H), 7.39 (d, 2H), 7.47 (s, 1H), 8.32 (s, 1H), 8.89 (s, 1H).



Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
110	piperidine	RT/18hrs/Nal	24	m/e 525 (M+H) ⁺	(d-6-DMSO, d values) 1.5-1.85 (m, 6H), 3.02 (m, 2H), 3.4-3.6 (m, 4H), 3.73 (s, 3H), 3.99 (s, 3H), 4.63 (m, 2H), 6.95 (m, 3H), 7.06 (m, 1H), 7.18 (m, 2H), 7.37 (d, 2H), 7.44 (s, 1H), 8.29 (s, 1H), 8.88 (s, 1H).
110	dimethyl amine	RT/18hrs/Nal/ EtOH	25	m/e 485 (M+H) ⁺	(d-6-DMSO, d values) 2.91 (m, 6H), 3.63 (m, 2H), 3.74 (s, 3H), 3.99 (s, 3H), 4.54 (m, 2H), 6.96 (m, 3H), 7.05 (m, 1H), 7.18 (m, 2H), 7.35 (d, 2H), 7.42 (s, 1H), 8.17 (s, 1H), 8.83 (s, 1H).
424		75°C/1hr/thioanisole/TFA	26	m/e 414 (M+H) ⁺	(d-6-DMSO, d values) 3.73 (s, 3H), 3.95 (s, 3H), 6.89 (d, 2H), 6.95 (m, 1H), 7.02 (m, 1H), 7.15 (m, 2H), 7.22 (d, 2H), 7.30 (s, 1H), 7.69 (s, 1H), 8.48 (s, 1H), 9.60 (bs, 1H), 9.94 (bs, 1H).
9		TFA / thioanisole / 90deg / 1.5 hours	27	m/e 414 (M ⁺ +H)	(d-6-DMSO, d values) 3.75 (s, 3H), 3.91 (s, 3H), 6.89 (d, 2H), 6.94 (m, 1H), 7.02 (d, 1H), 7.16 (m, 3H), 7.23 (m, 1H), 7.73 (s, 1H), 8.31 (s, 1H), 9.33 (s, 1H), 10.31 (broad, 1H).
26	2-chloromethylpyridine	RT/96hr/ KOtBu, /DMA	28	m/e 505 (M ⁺ +H)	(d-6-DMSO, d values) 3.74 (s, 3H), 4.01 (s, 3H), 5.39 (s, 2H), 6.95 (m, 3H), 7.04 (d, 1H), 7.17 (m, 2H), 7.41 (m, 3H), 7.50 (s, 1H), 7.63 (d, 1H), 7.93 (m, 1H), 8.34 (s, 1H), 8.61 (d, 1H), 8.97 (s, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
26	2-bromo thiazole	120°C/18hrs/ KOH/DMA	29	m/e 499 (M ⁺ +H)	(d-6-DMSO, d values) 3.73 (s, 3H), 3.99 (s, 3H), 6.96 (m, 4H), 7.05 (m, 1H), 7.18 (m, 2H), 7.36 (m, 3H), 7.41 (s, 1H), 7.96 (s, 1H), 8.85 (s, 1H).
26	2-chloro pyrimidine	100°C/18hrs/ K ₂ C O ₃ /DMA	30	m/e 492 (M ⁺ +H)	(d-6-DMSO, d values) 3.73 (s, 3H), 3.90 (s, 3H), 6.95 (m, 3H), 7.04 (m, 1H), 7.17 (m, 2H), 7.31 (m, 1H), 7.37 (m, 2H), 7.60 (s, 1H), 8.66 (m, 3H), 9.01 (s, 1H).
26	2-bromo pyridine	120°C/18hrs/ C ₂ C O ₃ /DMA	31	m/e 491 (M ⁺ +H)	(d-6-DMSO, d values) 3.71 (s, 3H), 3.89 (s, 3H), 6.93 (m, 3H), 7.03 (m, 1H), 7.16 (m, 3H), 7.36 (d, 2H), 7.51 (s, 1H), 7.89 (m, 1H), 8.05 (m, 1H), 8.35 (s, 1H), 8.93 (s, 1H)
118	morpholine	78°C/3hr/ethanol	33	m/e 541 (M ⁺ +H)	(d-6-DMSO, d values) 2.35 (m, 2H), 3.10 (m, 2H), 3.48 (d, 4H), 3.74 (s, 3H), 3.92 (m, 4H), 3.98 (s, 3H), 4.31 (t, 2H), 6.95 (m, 3H), 7.05 (m, 1H), 7.17 (m, 2H), 7.39 (d, 2H), 7.56 (s, 1H), 8.25 (s, 1H), 8.89 (s, 1H), 11.22 (broad, 1H), 11.26 (broad, 1H)
26		RT/18hr/DMA / KOtBu, /18- crown-6	34	m/e 511 (M ⁺ +H)	(d-6-DMSO, d values) 1.90 (m, 2H), 2.08-2.40 (m, 3H), 3.73 (s, 3H), 3.98 (s, 3H), 4.15 (m, 2H), 6.98 (m, 3H), 7.08 (m, 1H), 7.18 (m, 3H), 7.39 (d, 2H), 7.58 (s, 1H), 7.78 (s, 1H), 8.18 (s, 1H), 8.92 (s, 1H), 11.0 (bs, 1H).


Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
I18	piperazine	EtOH / 80deg / 3.5 hours	35	m/e 540 (M ⁺ +H)	(d-6-DMSO, d values) 2.33 (m, 2H), 3.32 (m, 2H), 3.48 (s, 8H), 3.73 (s, 3H), 3.97 (s, 3H), 4.31 (t, 2H), 6.96 (m, 3H), 7.04 (d, 1H), 7.18 (m, 2H), 7.37 (d, 2H), 7.48 (s, 1H), 8.13 (s, 1H), 8.87 (s, 1H), 11.04 (broad, 1H).
I18	pyrrolidine	EtOH / 80deg / 3.5 hours	36	m/e 525 (M ⁺ +H)	(d-6-DMSO, d values) 1.95 (broad, 2H), 2.28 (m, 2H), 3.03 (broad, 2H), 3.31 (t, 2H), 3.58 (broad, 2H), 3.73 (s, 3H), 3.97 (s, 3H), 4.29 (t, 2H), 6.96 (m, 3H), 7.05 (d, 1H), 7.18 (m, 2H), 7.39 (d, 2H), 7.47 (s, 1H), 8.14 (s, 1H), 8.86 (s, 1H), 11.06 (broad, 1H).
I18	Piperidine	EtOH / 80deg / 3.5 hours	37	m/e 539 (M ⁺ +H)	(d-6-DMSO, d values) 1.74 (m, 4H), 2.30 (m, 2H), 2.44 (m, 2H), 2.90 (m, 2H), 3.20 (t, 2H), 3.47 (m, 2H), 3.72 (s, 3H), 3.95 (s, 3H), 4.28 (t, 2H), 6.94 (m, 3H), 7.04 (d, 1H), 7.17 (m, 2H), 7.38 (d, 2H), 7.49 (s, 1H), 8.11 (s, 1H), 8.84 (s, 1H).
I18	N- (2 hydroxyethyl) piperazine	EtOH / 80deg / 7 hours	38	m/e 584 (M ⁺ +H)	(d-6-DMSO, d values @ 373deg K) 2.27 (m, 2H), 3.18 (m, 4H), 3.43 (s, 4H), 3.53 (s, 4H), 3.77 (s, 3H), 3.82 (t, 2H), 3.98 (s, 3H), 4.33 (t, 2H), 6.97 (m, 3H), 7.04 (d, 1H), 7.16 (m, 2H), 7.35 (d, 2H), 7.56 (s, 1H), 8.09 (s, 1H), 8.67 (s, 1H).

127

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
27	2-chloromethyl-pyridine	RT/48hr/DMSO/ KOtBu ₄ (1M in THF)	39	m/e 505 (M ⁺ +H)	(d-6-DMSO, d values) 3.73 (s, 3H), 4.01 (s, 3H), 5.41 (s, 2H), 6.96 (m, 3H), 7.05 (m, 1H), 7.18 (m, 2H), 7.40 (m, 3H), 7.54 (s, 1H), 7.58 (d, 1H), 7.89 (m, 1H), 8.21 (s, 1H), 8.63 (d, 1H), 8.96 (s, 1H), 11.10 (broad, 1H)
27	3-chloromethyl-pyridine	RT/48hr/DMSO/ KOtBu ₄ (1M in THF)	40	m/e 505 (M ⁺ +H)	(d-6-DMSO, d values) 3.74 (s, 3H), 3.98 (s, 3H), 5.40 (s, 2H), 6.96 (m, 3H), 7.05 (m, 1H), 7.18 (m, 2H), 7.38 (m, 2H), 7.58 (m, 1H), 7.63 (s, 1H), 8.09 (m, 1H), 8.19 (s, 1H), 8.65 (d, 1H), 8.82 (d, 1H), 8.86 (s, 1H), 11.04 (broad, 1H)
27		RT/96hr/ DMSO/ KOtBu ₄ (1M in THF)	41	m/e 511 (M ⁺ +H)	(d-6-DMSO, d values) 1.93 (m, 1H), 2.10 (m, 1H), 2.20 (m, 1H), 2.34 (m, 1H), 3.74 (s, 3H), 3.90 (s, 3H), 3.94 (m, 1H), 4.10 (m, 2H), 6.90 (d, 2H), 6.93 (m, 1H), 7.02 (m, 1H), 7.16 (m, 2H), 7.23 (d, 2H), 7.32 (s, 1H), 7.73 (s, 1H), 7.76 (s, 1H), 8.37 (s, 1H), 9.41 (s, 1H)
27	N-(2-chloroethyl)piperidine	RT/96hr/ powdered KOH/DMSO	44	m/e 525 (M ⁺ +H)	(d-6-DMSO, d values) 1.77 (m, 6H), 3.06 (m, 2H), 3.56 (m, 4H), 3.74 (s, 3H), 3.98 (s, 3H), 4.63 (t, 2H), 6.95 (m, 3H), 7.04 (m, 1H), 7.18 (m, 2H), 7.36 (d, 2H), 7.50 (s, 1H), 8.11 (s, 1H), 8.81 (s, 1H), 10.47 (broad, 1H), 10.75 (broad, 1H)

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
27	 WO 9965867	RT/120hr/DM SO KOtBu ₄ (1M in THF)	48	m/e 611 (M ⁺ +H)	(d-6-DMSO, d values) 1.23 (m, 2H), 1.40 (s, 9H), 1.78 (m, 2H), 2.02 (broad, 1H), 2.77 (m, 2H), 3.75 (s, 3H), 3.91 (s, 3H), 4.00 (m, 4H), 6.91 (m, 3H), 7.02 (m, 1H), 7.15 (m, 2H), 7.23 (d, 2H), 7.30 (s, 1H), 7.75 (s, 1H), 8.36 (s, 1H), 9.38 (s, 1H)
26	F ₃ CCH ₂ O-S (O) ₂ .CH ₃	120°C/20hr/ DMA/ KOtBu, /18-crown-6	50	m/e 496.1 (M ⁺ +H)	(CDCl ₃ , d values) 3.76 (s, 3H), 3.94 (s, 3H), 4.07 (q, 2H), 6.78 (s, 1H), 6.84-7.12 (m, 9H), 7.30 (s, 1H), 8.52 (s, 1H). Intermediate 1. M461666
26	CH ₂ CHCH ₂ - Br	23°C/20hr/DM A/ KOtBu, /18-crown-6	51	m/e 454 (M ⁺ +H)	(d-6-DMSO, d values) 3.74 (s, 3H), 3.93 (s, 3H), 4.8 (d, 2H), 5.29 (d, 1H), 5.44 (d, 1H), 6.03-6.2 (m, 1H), 6.86-7.26 (m, 8H), 7.3 (s, 1H), 7.77 (s, 1H), 8.37 (s, 1H), 9.36 (s, 1H).
49		RT/18hrs/NaO H/MeOH/wate r	52	m/e 472 (M+H) ⁺	(d-6-DMSO, d values) 3.77 (s, 3H), 3.97 (s, 3H), 4.85 (s, 2H), 6.92 (d, 2H), 6.96 (m, 1H), 7.03 (m, 1H), 7.18 (m, 2H), 7.25 (d, 2H), 7.33 (s, 1H), 7.77 (s, 1H), 8.39 (s, 1H), 9.44 (s, 1H).
26		RT/4hrs/ KOtBu / 18-C-6/n- Bu ₄ NI/DMA	53	m/e 511 (M+H) ⁺	(d-6-DMSO, d values) 1.91 (m, 2H), 2.11-2.30 (m, 3H), 3.76 (s, 3H), 4.00 (s, 3H), 4.12 (m, 2H), 6.99 (m, 3H), 7.08 (m, 1H), 7.21 (m, 3H), 7.40 (d, 2H), 7.80 (s, 1H), 8.20 (s, 1H), 8.92 (s, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
26	$\text{CH}\equiv\text{CCH}_2\text{Br}$	$23^\circ\text{C}/20\text{hr}/$ DMA/ KOtBu , /18-crown-6	54	m/e 452 ($\text{M}^+ + \text{H}$)	d-6-DMSO, d values) 3.6 (t, 1H), 3.73 (s, 3H), 3.94 (s, 3H), 4.92 (d, 2H), 6.84-7.3 (m, 8H), 7.33 (s, 1H), 7.84 (s, 1H), 8.38 (s, 1H), 9.38 (s, 1H).
26	CH_3OCH_2 CH_2Br	$23^\circ\text{C}/20\text{hr}/\text{DM}$ A/ KOtBu , /18-crown-6	55	m/e 472 ($\text{M}^+ + \text{H}$)	(d-6-DMSO, d values) 3.32 (s, 3H), 3.71 (t, 2H), 3.73 (s, 3H), 3.93 (s, 3H), 4.21 (t, 2H), 6.85-7.28 (m, 8H), 7.3 (s, 1H), 7.75 (s, 1H), 8.36 (s, 1H), 9.36 (s, 1H).
52	morpholine	RT/64hrs/ EDC/DMAP/ DCM	57	m/e 541 ($\text{M} + \text{H}$) ⁺	(d-6-DMSO, d values) 3.48 (m, 4H), 3.61 (m, 4H), 3.76 (s, 3H), 4.01 (s, 3H), 5.11 (s, 2H), 6.96 (m, 3H), 7.08 (m, 1H), 7.21 (m, 2H), 7.40 (m, 3H), 8.15 (s, 1H), 8.89 (s, 1H).
52	N-methyl piperazine	RT/18hrs /EDC/DMAP/ DCM	58	m/e 552 ($\text{M} + \text{H}$) ⁺	(d-6-DMSO, d values) 2.80 (bs, 3H), 3.00-3.60 (m, 8H (under H_2O peak)), 3.75 (s, 3H), 4.01 (s, 3H), 5.18 (s, 2H), 6.95 (m, 3H), 7.05 (m, 1H), 7.18 (m, 2H), 7.39 (m, 3H), 7.70 (bs, 1H), 8.33 (bs, 1H), 8.78 (bs, 1H).
52	allylamine	RT/18hrs/EDC /DMAP/DCM	59	m/e 511 ($\text{M} + \text{H}$) ⁺	(d-6-DMSO, d values) 3.74 (s, 3H), 3.78 (m, 2H), 4.02 (s, 3H), 4.77 (s, 2H), 5.09 (m, 2H), 5.80 (m, 1H), 6.97 (m, 3H), 7.08 (m, 1H), 7.20 (m, 2H), 7.37 (d, 2H), 7.44 (s, 1H), 8.21 (s, 1H), 8.16 (m, 1H), 8.85 (bs, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
52	methylamine	RT/18hrs/THF/ EDC/DMAP/D CM	60	m/e 485 (M+H) ⁺	(d-6-DMSO, d values) 2.68 (m, 3H), 3.78 (s, 3H), 4.03 (s, 3H), 4.70 (s, 2H), 6.96 (m, 3H), 7.06 (m, 1H), 7.19 (m, 2H), 7.36 (s, 1H), 7.40 (m, 2H), 7.89 (bs, 1H), 8.08 (s, 1H), 8.86 (s, 1H), 10.68 (bs, 1H).
52	methoxy ethanolamine	RT/18hrs/EDC /DMAP/DCM	61	m/e 529 (M+H) ⁺	(d-6-DMSO, d values) 3.25 (s, 3H), 3.75 (s, 3H), 4.01 (s, 3H), 4.73 (s, 2H), 6.96 (m, 3H), 7.05 (m, 1H), 7.20 (m, 2H), 7.37 (s, 1H), 7.40 (m, 2H), 7.95 (bs, 1H), 8.07 (s, 1H), 8.86 (s, 1H), 10.70 (bs, 1H).
48		95°C/18hr/HC HO(aq.)/HCO OH	63	m/e 525 (M ⁺ +H)	(d-6-DMSO, d values) 1.73 (m, 2H), 2.04 (m, 2H), 2.20 (m, 1H), 2.78 (s, 3H), 3.06 (m, 2H), 3.44 (m, 2H), 3.77 (s, 3H), 3.96 (s, 3H), 4.14 (d, 2H), 6.96 (m, 3H), 7.03 (m, 1H), 7.16 (m, 2H), 7.29 (m, 2H), 7.44 (s, 1H), 7.89 (s, 1H), 8.58 (s, 1H)
27		55°C/30hr/DM SO/KOtBu(1M in THF)	64	m/e 511 (M ⁺ +H)	(d-6-DMSO, d values) 1.93 (m, 1H), 2.10 (m, 1H), 2.20 (m, 1H), 2.34 (m, 1H), 3.74 (s, 3H), 3.90 (s, 3H), 3.94 (m, 1H), 4.10 (m, 2H), 6.90 (d, 2H), 6.93 (m, 1H), 7.02 (m, 1H), 7.16 (m, 2H), 7.23 (d, 2H), 7.32 (s, 1H), 7.73 (s, 1H), 7.76 (s, 1H), 8.37 (s, 1H), 9.41 (s, 1H).
65		RT/18hrs/ NaOH/MeOH/ water	66	m/e 472 (M+H) ⁺	(d-6-DMSO, d values) 3.74 (s, 3H), 3.92 (s, 3H), 4.88 (s, 2H), 6.89 (d, 2H), 6.95 (m, 1H), 7.01 (m, 1H), 7.16 (m, 3H), 7.24 (d, 2H), 7.76 (s, 1H), 8.35 (s, 1H), 9.43 (bs, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
27	$\text{CH}\equiv\text{CCH}_2\text{Br}$	23°C/20hr/DM A/KOtBu	67	m/e 452.2 (M ⁺ +H)	(d-6-DMSO, d values) 3.63 (t, 1H), 3.75 (s, 3H), 3.9 (s, 3H), 5.0(d, 2H), 6.84-7.04 (m, 4H), 7.1-7.28 (m, 4H), 7.4 (s, 1H), 7.78 (s, 1H), 8.37(s, 1H), 9.42(s, 1H).
66	cyclopropyl amine	RT/18hrs/EDC / DMAP/DCM	68	m/e 511 (M+H) ⁺	(d-6-DMSO, d values) 0.47 (m, 2H), 0.63 (m, 2H), 2.66 (m, 1H), 3.74 (s, 3H), 3.96 (s, 3H), 4.70 (s, 2H), 6.95 (m, 3H), 7.05 (m, 1H), 7.17 (m, 2H), 7.21 (m, 1H), 7.37 (d, 2H), 8.03 (s, 1H), 8.29 (m, 1H), 8.81 (s, 1H).
62	Chloropropyl morpholine	60°C/18hrs/ KO ^t Bu/Bu ₄ NI/ 18-C-6/DMA	70	m/e 553.6 (M-H) ⁻	(d-6-DMSO, d values) 2.33 (m, 2H), 3.12 (m, 2H), 3.29 (m, 2H), 3.49 (m, 2H), 3.83 (m, 2H), 3.83 (m, 2H), 3.95 (m, 2H), 4.00 (s, 3H), 4.03 (m, 2H), 4.30 (m, 2H), 6.96 (m, 3H), 7.05 (m, 1H), 7.17 (m, 2H), 7.38 (m, 2H), 7.54 (s, 1H), 8.18 (s, 1H), 8.88 (s, 1H).
102	diphenyl phosphoryl azide	100°C/18hrs/N Et ₃ /t-BuOH	71	m/e 513 (M+H) ⁺	(d-6-DMSO, d values) (broadened due to rotamers) 1.48 (bs, 9H), 3.71 (bs, 3H), 3.99 (bs, 3H), 6.92 (bm, 2H), 6.95 (bm, 3H), 7.03 (bm, 1H), 7.15 (bs, 1H), 7.40 (bm, 3H), 8.66 (bs, 1H), 8.80 (bs, 1H), 8.91 (bs, 1H), 10.93 (bs, 1H).
71		RT/2hrs/Et ₃ Si H/TFA	72	m/e 413 (M+H) ⁺	(d-6-DMSO, d values) 3.74 (s, 3H), 3.95 (s, 3H), 5.39 (bs, 2H), 6.85 (d, 2H), 6.91 (m, 1H), 6.96 (m, 1H), 7.10 (m, 3H), 7.20 (s, 1H), 7.29 (s, 1H), 8.24 (s, 1H), 9.06 (s, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
27	$\text{CH}\equiv\text{CCH}_2\text{Br}$	$23^\circ\text{C}/20\text{hr}/\text{DM A/KOtBu}$	67	m/e 452.2 (M^++H)	(d-6-DMSO, d values) 3.63 (t, 1H), 3.75 (s, 3H), 3.9 (s, 3H), 5.0(d, 2H), 6.84-7.04 (m, 4H), 7.1-7.28 (m, 4H), 7.4 (s, 1H), 7.78 (s, 1H), 8.37(s, 1H), 9.42(s, 1H).
66	cyclopropylamine	$\text{RT}/18\text{hrs}/\text{EDC}$ / DMAP/DCM	68	m/e 511 ($\text{M}+\text{H}$) ⁺	(d-6-DMSO, d values) 0.47 (m, 2H), 0.63 (m, 2H), 2.66 (m, 1H), 3.74 (s, 3H), 3.96 (s, 3H), 4.70 (s, 2H), 6.95 (m, 3H), 7.05 (m, 1H), 7.17 (m, 2H), 7.21 (m, 1H), 7.37 (d, 2H), 8.03 (s, 1H), 8.29 (m, 1H), 8.81 (s, 1H).
62	Chloropropyl morpholine	$60^\circ\text{C}/18\text{hrs}/\text{KO}^t\text{Bu}/\text{Bu}_4\text{NI}/18\text{-C-6}/\text{DMA}$	70	m/e 553.6 ($\text{M}-\text{H}^+$) ⁻	(d-6-DMSO, d values) 2.33 (m, 2H), 3.12 (m, 2H), 3.29 (m, 2H), 3.49 (m, 2H), 3.83 (m, 2H), 3.83 (m, 2H), 3.95 (m, 2H), 4.00 (s, 3H), 4.03 (m, 2H), 4.30 (m, 2H), 6.96 (m, 3H), 7.05 (m, 1H), 7.17 (m, 2H), 7.38 (m, 2H), 7.54 (s, 1H), 8.18 (s, 1H), 8.88 (s, 1H).
102	diphenylphosphorylazide	$100^\circ\text{C}/18\text{hrs}/\text{N Et}_3/t\text{-BuOH}$	71	m/e 513 ($\text{M}+\text{H}$) ⁺	(d-6-DMSO, d values) (broadened due to rotamers) 1.48 (bs, 9H), 3.71 (bs, 3H), 3.99 (bs, 3H), 6.92 (bm, 2H), 6.95 (bm, 3H), 7.03 (bm, 1H), 7.15 (bs, 1H), 7.40 (bm, 3H), 8.66 (bs, 1H), 8.80 (bs, 1H), 8.91 (bs, 1H), 10.93 (bs, 1H).
71		$\text{RT}/2\text{hrs}/\text{Et}_3\text{Si H}/\text{TFA}$	72	m/e 413 ($\text{M}+\text{H}$) ⁺	(d-6-DMSO, d values) 3.74 (s, 3H), 3.95 (s, 3H), 5.39 (bs, 2H), 6.85 (d, 2H), 6.91 (m, 1H), 6.96 (m, 1H), 7.10 (m, 3H), 7.20 (s, 1H), 7.29 (s, 1H), 8.24 (s, 1H), 9.06 (s, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
72	MeSO ₂ Cl	70°C/12hrs/ pyridine	73	m/e 490 (M+H) ⁺	(d-6-DMSO, d values) 3.06 (s, 3H), 3.74 (s, 3H), 3.99 (s, 3H), 6.89 (d, 2H), 6.95 (m, 1H), 7.01 (m, 1H), 7.13 (m, 2H), 7.22 (d, 2H), 7.37 (s, 1H), 8.21 (s, 1H), 8.44 (s, 1H), 9.24 (bs, 1H), 9.65 (bs, 1H).
425		RT/3days/NaO H/MeOH/ water	102	m/e 442 (M+H) ⁺	(d-6-DMSO, d values) 3.74 (s, 3H), 3.95 (s, 3H), 6.89 (d, 2H), 6.95 (m, 1H), 7.01 (m, 1H), 7.14 (m, 2H), 7.25 (d, 2H), 7.37 (s, 1H), 8.49 (s, 1H), 8.78 (s, 1H), 9.89 (bs, 1H).
111	1-Methyl piperazine	60°C/3hr/NaI	114	m/e 579 (M+H) ⁺	(d-6-DMSO, δ values) 2.25 (m, 2H), 2.80 (s, 3H), 3.24 - 3.53 (m under H ₂ O, 10H), 3.56 (m, 1H), 3.99 (s, 3H), 4.30 (m, 2H), 4.80 (d, 2H), 6.96 - 7.05 (m, 4H), 7.16 - 7.28 (m, 2H), 7.40 (m, 2H), 7.46 (s, 1H), 8.22 (s, 1H), 8.91 (s, 1H).
108	N-(3chloro- propyl) morpholine	RT/15min/KOt Bu/DMA then 60°C/4hr/nBu ₄ NI/18 crown 6	115		(d-6-DMSO, δ values) 2.23 - 2.36 (m, 2H), 3.03 - 3.16 (m, 2H), 3.24 - 3.34 (m, 2H), 3.42 - 3.51 (m, 2H), 3.71 - 3.83 (m, 2H), 3.92 - 4.03 (m, 5H), 4.35 (t, 2H), 6.75 (tt,), 6.90 (s, 1H), 7.00 - 7.06 (m, 2H), 7.21 - 7.28 (d, 2H), 7.46 - 7.56 (m, 4H), 8.31 (s, 1H), 8.92 (s, 1H).
112	1-Methyl- piperazine	60°C/3hr/NaI	118	m/e 640 (M+H) ⁺	(d-6-DMSO, δ values) 2.23 - 2.37 (m, 2H), 2.80 (s, 3H), 3.39 - 3.78 (m under H ₂ O, 10H), 4.00 (s, 3H), 4.35 (t, 2H), 6.76 (tt, 1H), 6.90 (m, 1H), 7.02 (dd, 1H), 7.24 (d, 2H), 7.45 (d, 1H), 7.50 - 7.56 (m, 3H), 8.37 (s, 1H), 8.93 (s, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
I12	1-Methyl-piperazine	60°C/3hr/NaI	119	m/e 585 (M+H) ⁺	(d-6-DMSO, δ values) 2.20 - 2.30 (m, 2H), 2.81 (s, 6H), 3.25 (m, 2H), 4.00 (s, 3H), 4.32 (t, 2H), 6.74 (tt, 1H), 6.90 (m, 1H), 7.02 (m, 1H), 7.24 (d, 2H), 7.44 - 7.56 (m, 4H), 8.27 (s, 1H), 8.95 (s, 1H).
I11	Morpholine	60°C/3hr/NaI	120	m/e 527 (M+H) ⁺	(d-6-DMSO, δ values) 1.87 - 2.00 (m, 2H), 2.32 - 2.40 (m, 2H), 3.50 - 3.59 (m, 4H), 3.77 - 3.88 (m, 4H), 3.94 (s, 3H), 4.13 (t, 2H), 6.78 (m, 1H), 6.87 - 7.02 (m, 5H), 7.22 (m, 2H), 7.30 (s, 1H), 7.75 (s, 1H), 8.36 (s, 1H), 9.39 (s, 1H), 9.49 (s, 1H).
I11	Dimethylamine	60°C/3hr/NaI/ MeOH	121	m/e 485 (M+H) ⁺	(d-6-DMSO, δ values) 2.07 (m, 2H), 2.54 (s, 6H), 2.86 (m, 2H), 3.93 (s, 3H), 4.15 (t, 2H), 6.78 (m, 1H), 6.89 - 7.00 (m, 5H), 7.22 (d, 2H), 7.31 (s, 1H), 7.75 (s, 1H), 8.38 (s, 1H), 9.38 (s, 1H), 9.48 (bs, 1H).
I13	1-Methyl-piperazine	80°C/3hr/NaI	127	m/e 580 (M+H) ⁺	(d-6-DMSO, δ values) 2.21 - 2.32 (m, 2H), 2.79 (s, 3H), 3.19 - 3.65 (m under H ₂ O, 10H), 4.00 (s, 3H), 4.32 (t, 2H), 4.57 (d, 2H), 5.16 (d, 1H), 5.28 (d, 1H), 5.86 - 6.00 (m, 1H), 6.93 - 7.00 (m, 3H), 7.06 (d, 1H), 7.17 (d, 2H), 7.39 (d, 2H), 7.52 (s, 1H), 8.32 (s, 1H), 8.91 (s, 1H), 9.70 (s, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
113	Dimethyl amine	80°C/3hr/NaI/ MeOH	128	m/e 525.4 (M+H) ⁺	(d-6-DMSO, δ values) 2.20 - 2.30 (m, 2H), 2.77 (s, 3H), 2.79 (s, 3H), 3.16 - 3.31 (m under H ₂ O, 2H), 3.99 (s, 3H), 4.30 (t, 2H), 4.57 (d, 2H), 5.17 (d, 1H), 5.29 (d, 1H), 5.86 - 6.00 (m, 1H), 6.92 - 7.00 (m, 3H), 7.06 (d, 1H), 7.16 (d, 2H), 7.39 (d, 2H), 7.49 (s, 1H), 8.29 (s, 1H), 8.89 (s, 1H).
110	CH ₃ CCH ₂ Br	23°C/20hr/DM A/KOtBu	131	m/e 447.2 (M ⁺ +H)	(d-6-DMSO, δ values) 3.61(t, 1H), 3.94 (s, 3H), 4.93(d, 2H), 6.92 (d, 1H), 7.2-7.3 (m, 3H), 7.35 (s, 1H), 7.38 (d, 2H), 7.62(t, 1H) 7.88(s, 1H), 7.9 (d, 1H), 8.43 (s, 1H), 9.52 (s, 1H),
130	CH ₃ CCH ₂ Br	23°C/20hr/DM A/KOtBu	132	m/e 447 (M ⁺ +H)	(d-6-DMSO, δ values) 3.64(t, 1H), 3.92 (s, 3H), 5.0(d, 2H), 6.93 (d, 1H), 7.2-7.3 (m, 3H), 7.4 (d, 2H), 7.42 (s, 1H), 7.6(t, 1H) 7.8(s, 1H), 7.89 (d, 1H), 8.42 (s, 1H), 9.6 (s, 1H),
114	Morpholine	RT/18hrs/NaI	137	m/e 570 (M+H) ⁺	(d-6-DMSO, δ values) 2.29 (m, 2H), 3.10 (m, 2H), 3.29 (m, 2H), 3.47 (m, 2H), 3.60 (m, 2H), 3.79 (m, 2H), 4.00 (m, 5H), 4.32 (m, 2H), 6.95 (m, 4H), 7.14 (m, 2H), 7.41 (m, 2H), 7.47 (s, 1H), 8.28 (s, 1H), 8.95 (s, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
115	Morpholine	RT/18hrs/NaI	138	m/e 571 (M+H) ⁺	(d-6-DMSO, d values) 2.34 (m, 2H), 3.08 (m, 2H), 3.29 (m, 2H), 3.49 (m, 2H), 3.62 (m, 2H), 3.84 (m, 2H), 3.93 (m, 1H), 4.01 (m, 5H), 4.31 (m, 2H), 6.97 (m, 4H), 7.17 (m, 2H), 7.41 (m, 2H), 7.57 (s, 1H), 8.25 (s, 1H), 8.93 (s, 1H).
130	N-(3-chloropropyl)-morpholine	80°C/4hr/ KOtBu/tetrabu tylammonium iodide/18- crown-6	139	m/e 536.04 (M ⁺ +H)	(d-6-DMSO, d values) 1.94 (m, 2H), 2.3-2.5 (m, 4H), 3.29 (m, 2H), 3.57 (m, 4H), 3.92 (s, 3H), 4.2 (t, 2H), 6.93 (d, 1H), 7.13 (d, 2H), 7.16 (s, 1H), 7.2 (s, 1H), 7.29(d, 2H) 7.62 (t, 1H), 7.76 (s, 1H), 7.89 (d, 1H), 8.4 (s, 1H), 9.54(s, 1H).
427	Chloropropyl morpholine	60°C/18hrs/K O'Bu/Bu ₄ NI/18 -C-6/DMA	142	m/e 596 (M-H) ⁺	(d-6-DMSO, d values) 2.34 (m, 2H), 2.61 (m, 3H), 3.11 (m, 2H), 3.26 (m, 2H), 3.47 (m, 2H), 3.82 (m, 2H), 3.96 (m, 2H), 4.01 (m, 3H), 4.30 (m, 2H), 4.47 (s, 2H), 7.05 (m, 6H), 7.29 (m, 1H), 7.43 (m, 2H), 7.54 (m, 2H), 8.21 (s, 1H), 8.92 (s, 1H).
129	Chloropropyl morpholine	60°C/18hrs/K O'Bu/Bu ₄ NI/18 -C-6/DMA	143	m/e 596 (M-H) ⁺	(d-6-DMSO, d values) 2.31 (m, 2H), 2.62 (m, 3H), 3.13 (m, 2H), 3.29 (m, 2H), 3.46 (m, 2H), 3.82 (m, 2H), 3.92 (m, 2H), 3.99 (m, 3H), 4.34 (m, 2H), 4.47 (s, 2H), 7.06 (m, 6H), 7.43 (m, 2H), 7.54 (m, 2H), 8.37 (s, 1H), 8.93 (s, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
113	Chloropropyl morpholine	60°C/18hrs/K O'Bu/Bu ₄ NI/18 -C-6/DMA	154	m/e 625.5 (M-H) ⁺	(d-6-DMSO, d values) 2.33 (m, 2H), 3.11 (m, 2H), 3.29 (m, 2H), 3.35 (m, 2H), 3.82 (m, 2H), 3.96 (m, 2H), 3.98 (s, 3H), 4.32 (m, 2H), 6.76 (tt, 1H), 7.04 (m, 2H), 7.24 (m, 2H), 7.49 (m, 1H), 7.54 (m, 3H), 8.21 (s, 1H), 8.89 (s, 1H).
219		75°C/1.5hr/TF A/ Thioanisole	218	m/e 417.4 (M+H) ⁺	(d-6-DMSO, δ values) 2.24 (s, 3H), 3.94 (s, 3H), 7.21 - 7.39 (m, 6H), 7.77 (s, 1H), 8.00 (s, 1H), 8.25 (s, 1H), 9.20 (s, 1H), 10.34 (bs, 1H).
17	Piperidine	RT/18hrs/NaI	170	m/e 581.5 (M-H) ⁺	(d-6-DMSO, d values) 1.51 (m, 2H), 1.71 (m, 4H), 2.18 (m, 2H), 3.08 (m, 6H), 3.91 (s, 3H), 4.24 (m, 2H), 4.66 (s, 2H), 7.03 (m, 6H), 7.24 (m, 2H), 7.33 (s, 1H), 7.76 (s, 1H), 8.37 (s, 1H), 9.46 (s, 1H).
14 (Ex. 7)	morpholine	23°C/24hr/NaI	175	m/e 628.58 (M ⁺ +H)	(d-6-DMSO, d values) 2.32 (m, 2H), 3.0-3.64 (m, 10H), 3.8 (t, 2H), 3.96 (m, 2H), 3.98 (s, 3H), 4.28 (t, 2H), 4.48(s, 2H), 6.94-7.21 (m, 6H), 7.4 (d, 2H), 7.52 (s, 1H), 7.6 (t, 1H), 8.11 (s, 1H), 8.85 (s, 1H).
14	pyrrolidine	23°C/24hr/NaI	176	m/e 612.56 (M ⁺ +H)	(d-6-DMSO, d values) 1.89 (m, 2H), 2.0 (m, 2H), 2.28 (m, 2H), 3.02 (m, 2H), 3.15 (q, 2H), 3.2-3.7 (m, 6H), 3.98 (s, 3H), 4.29 (t, 2H), 4.48(s, 2H), 6.95-7.21 (m, 6H), 7.4 (d, 2H), 7.5 (s, 1H), 7.6 (t, 1H), 8.16 (s, 1H), 8.86 (s, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
14	dimethyl-morpholine	23°C/24hr/NaI	177	m/e 656.6 (M ⁺ +H)	(d-6-DMSO, d values) 1.14 (d, 6H), 2.22-2.74 (m, 4H), 3.1-3.62 (m, 8H), 3.9-4.09 (m, 5H), 4.3 (t, 2H), 4.48 (s, 2H), 6.95-7.21 (m, 6H), 7.4 (d, 2H), 7.47 (s, 1H), 7.6 (t, 1H), 8.1 (s, 1H), 8.85 (s, 1H).
14	1-acetyl-piperazine	23°C/24hr/NaI	194	m/e 669.59 (M ⁺ +H)	(d-6-DMSO, d values) 2.0 (s, 3H), 2.32 (m, 2H), 2.84-3.7 (m, 14H), 3.99 (s, 3H), 4.3 (t, 2H), 4.42 (br.d., 1H), 4.48 (s, 2H), 6.95-7.22 (m, 6H), 7.4 (d, 2H), 7.46 (s, 1H), 7.6 (t, 1H), 8.18 (s, 1H), 8.84 (s, 1H), 9.2 (br.s., 1H).
27	4-chloromethyl-pyridine	RT/48hr/DMS O/ KOtBu(1M in THF)	197	m/e 505 (M ⁺ +H)	(d-6-DMSO, d values) 3.75 (s, 3H), 4.03 (s, 3H), 5.53 (s, 2H), 6.96 (m, 3H), 7.05 (m, 1H), 7.18 (m, 2H), 7.38 (d, 2H), 7.53 (s, 1H), 7.74 (d, 2H), 8.20 (s, 1H), 8.74 (d, 2H), 8.79 (s, 1H), 10.93 (broad, 1H)
116	1-Methyl-piperazine	60°C/16hr/NaI	204	m/e 555 (M+H) ⁺	(d-6-DMSO, δ values) 2.24 - 2.37 (m, 2H), 2.78 (s, 3H), 3.19 - 3.62 (m under H ₂ O, 10H), 3.67 (s, 3H), 3.99 (s, 3H), 4.36 (t, 2H), 6.98 (t, 1H), 7.08 - 7.15 (m, 3H), 7.21 (t, 1H), 7.55 (s, 1H), 7.90 (dd, 1H), 8.19 (d, 1H), 8.41 (m, 1H), 8.95 (s, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
I17	1-Methyl-piperazine	60°C/16hr/NaI	205	m/e 555 (M+H) ⁺	(d-6-DMSO, δ values) 2.26 - 2.38 (m, 2H), 2.80 (s, 3H), 3.20 - 3.66 (m under H ₂ O, 10H), 3.70 (s, 3H), 4.01 (s, 3H), 4.31 (t, 2H), 6.98 (t, 1H), 7.09 - 7.16 (m, 3H), 7.21 (m, 1H), 7.52 (s, 1H), 7.92 (dd, 1H), 8.19 (d, 1H), 8.25 (s, 1H), 8.92 (s, 1H), 9.62 (bs, 1H).
I16	Dimethyl amine	60°C/16hr/NaI/ MeOH	206	m/e 500 (M+H) ⁺	(d-6-DMSO, δ values) 2.19 - 2.30 (m, 2H), 2.79 (s, 3H), 2.80 (s, 3H), 3.16 - 3.28 (m, 2H), 3.68 (s, 3H), 3.99 (s, 3H), 4.32 (t, 2H), 6.98 (t, 1H), 7.08 - 7.16 (m, 3H), 7.21 (m, 1H), 7.48 (s, 1H), 7.89 (d, 1H), 8.17 (d, 1H), 8.34 (s, 1H), 8.88 (s, 1H).
I17	Dimethyl amine	60°C/16hr/NaI/ MeOH	207	m/e 500 (M+H) ⁺	(d-6-DMSO, δ values) 2.18 - 2.28 (m, 2H), 2.76 (s, 6H), 3.16 - 3.22 (m, 2H), 3.68 (s, 3H), 3.95 (s, 3H), 4.26 (t, 2H), 6.96 (t, 1H), 7.00 (d, 1H), 7.11 (d, 2H), 7.19 (m, 1H), 7.32 (s, 1H), 7.76 (dd, 1H), 7.92 (s, 1H), 8.06 (d, 1H), 8.38 (s, 1H), 9.73 (s, 1H).
220	N-(3-chloro-propyl) morpholine	RT/15min/ KOtBu/DMA then RT/18hr/ nBu ₄ Ni/18-crown-6	208		(d-6-DMSO, δ values) 2.24 - 2.35 (m, 2H), 3.04 - 3.16 (m, 2H), 3.24 - 3.33 (m, 2H), 3.43 - 3.51 (m, 2H), 3.68 (s, 3H), 3.72 - 3.83 (m, 2H), 3.91 - 3.98 (m, 2H), 3.99 (s, 3H), 4.34 (t, 2H), 6.98 (t, 1H), 7.08 - 7.14 (m, 3H), 7.21 (m, 1H), 7.47 (s, 1H), 7.90 (dd, 1H), 8.19 (d, 1H), 8.31 (m, 1H), 8.92 (s, 1H).


Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
221	N-(3-chloropropyl) morpholine	i) RT/15min/KOt Bu/DMA then ii) RT/18hr/(2)/ nBu ₄ NI/18- crown-6	209		(d-6-DMSO, δ values) 2.27 - 2.36 (m, 2H), 3.04 - 3.19 (m, 2H), 3.24 - 3.31 (m, 2H), 3.44 - 3.54 (m, 2H), 3.68 (s, 3H), 3.74 - 3.86 (m, 2H), 3.93 - 3.98 (m, 2H), 3.99 (s, 3H), 4.31 (t, 2H), 6.98 (t, 1H), 7.08 - 7.15 (m, 3H), 7.21 (t, 1H), 7.50 (s, 1H), 7.91 (dd, 1H), 8.19 (m, 2H), 8.92 (s, 1H).
203	N-(3-chloropropyl) morpholine	i) RT/15min/KOt Bu/DMA then ii) RT/16hr/(2)/ nBu ₄ NI/18- crown-6	210	m/e 543 (M+H) ⁺	(d-6-DMSO D4 Acetic, δ values) 2.23 - 2.37 (m, 2H), 3.04 - 3.17 (m, 2H), 3.29 (t, 2H), 3.44 - 3.54 (m, 2H), 3.67 (s, 3H), 3.73 - 3.84 (m, 2H), 3.92 - 3.99 (m, 2H), 4.00 (s, 3H), 4.31 (t, 2H), 6.99 (t, 1H), 7.11 - 7.28 (m, 3H), 7.49 (s, 1H), 8.18 (s, 1H), 8.75 (s, 2H), 8.90 (s, 1H).
222		75°C/1.5hr/ TFA/ Thioanisole	211	m/e 429.4 (M+H) ⁺	(d-6-DMSO, δ values) 2.20 (s, 3H), 3.67 (s, 3H), 3.93 (s, 3H), 6.93 - 7.00 (m, 2H), 7.08 - 7.12 (m, 2H), 7.16 - 7.20 (m, 2H), 7.79 (s, 1H), 7.98 (s, 1H), 8.25 (s, 1H), 9.20 (s, 1H), 10.31 (bs, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
I20	pyrrolidine	RT/48hr/NaI	214	m/e 623.5 (M ⁺ +H)	(d-6-DMSO d-4-Acetic, δ values) 0.20 (m, 2H), 0.41 (m, 2H), 0.96 (m, 1H), 1.86 - 2.09 (m, 4H), 2.25 - 2.36 (m, 2H), 3.00 - 3.12 (m, 4H), 3.34 (t, 2H), 3.61 (m, 2H), 4.04 (s, 3H), 4.34 (t, 2H), 4.48 (s, 2H), 6.72 - 6.81 (m, 2H), 6.85 (dd, 1H), 7.22 (d, 1H), 7.35 (t, 1H), 7.53 (s, 1H), 7.99 (dd, 1H), 8.24 (s, 1H), 8.35 (d, 1H), 8.95 (s, 1H).
I6	Morpholine	RT/48hr/NaI	215	m/e 542.5 (M+H) ⁺	(d-6-DMSO, δ values) 1.99 (t, 2H), 2.34 - 2.45 (m, 4H), 3.52 - 3.61 (m, 4H), 3.79 (s, 3H), 3.96 (s, 3H), 4.20 (t, 2H), 7.03 (t, 1H), 7.20 - 7.32 (m, 3H), 7.40 (s, 1H), 7.55 (d, 1H), 7.78 (m, 1H), 8.06 (s, 1H), 8.61 (d, 1H), 9.38 (s, 1H), 9.47 (bs, 1H).
I33	Morpholine	RT/48hr/NaI	216	m/e 599.5 (M+H) ⁺	(d-6-DMSO, δ values) 2.21 (m, 2H), 2.61 (d, 3H), 4.00 (s, 3H), 4.27 (t, 2H), 4.52 (s, 2H), 7.09 (t, 1H), 7.18 (d, 1H), 7.27 (t, 2H), 7.49 (s, 1H), 7.64 (m, 1H), 7.75 (m, 1H), 7.87 (dd, 1H), 8.15 (s, 1H), 8.77 (d, 1H), 9.49 (s, 1H), 9.70 (bs, 1H). HPLC time 6.99, 93.5%

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
I24	dimethyl morpholine	RT/72hr/NaI	223	m/e 653.6 (M ⁺ +H)	(d-6-DMSO d-4-Acetic, δ values) 0.48 (m, 2H), 0.61 (m, 2H), 1.14 (s, 3H), 1.16 (s, 3H), 2.29 - 2.37 (m, 2H), 2.59 - 2.71 (m, 3H), 3.26 (m, 2H), 3.50 (d, 2H), 3.89 - 3.96 (m, 2H), 4.00 (s, 3H), 4.30 (t, 2H), 4.41 (s, 2H), 6.68 - 6.74 (m, 2H), 6.77 (d, 1H), 7.19 (d, 1H), 7.31 (t, 1H), 7.48 (s, 1H), 7.97 (dd, 1H), 8.19 (s, 1H), 8.31 (d, 1H), 8.93 (s, 1H).
I24	pyrrolidine	RT/48hr/NaI	224	m/e 609.5 (M ⁺ +H)	(d-6-DMSO d-4-Acetic, δ values) 0.47 (m, 2H), 0.61 (m, 2H), 1.84 - 2.06 (m, 4H), 2.21 - 2.31 (m, 2H), 2.68 (m, 1H), 2.98 - 3.10 (m, 2H), 3.31 (t, 2H), 3.59 (m, 2H), 4.00 (s, 3H), 4.30 (t, 2H), 4.41 (s, 2H), 6.68 - 6.72 (m, 2H), 6.76 (dd, 1H), 7.18 (d, 1H), 7.31 (t, 1H), 7.48 (s, 1H), 7.95 (dd, 1H), 8.15 (s, 1H), 8.29 (d, 1H), 8.89 (s, 1H).
I20	dimethylmorpholine	RT/72hr/NaI	225	m/e 677.6 (M ⁺ +H)	(d-6-DMSO d-4-Acetic, δ values) 0.20 (m, 2H), 0.31 (m, 2H), 0.96 (m, 1H), 1.15 (s, 3H), 1.19 (s, 3H), 2.36 (m, 2H), 2.70 (m, 2H), 3.04 (d, 2H), 3.20 (t, 2H), 3.55 (d, 2H), 3.94 - 4.02 (m, 2H), 4.04 (s, 3H), 4.34 (m, 2H), 4.50 (s, 2H), 6.73 - 6.80 (m, 2H), 6.85 (dd, 1H), 7.23 (d, 1H), 7.36 (t, 1H), 7.51 (s, 1H), 8.00 (dd, 1H), 8.20 (s, 1H), 8.33 (d, 1H), 8.95 (s, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
272		75°C/2hrs/thio anisole/TFA	273	m/e 391 (M+H) ⁺	(d-6-DMSO, d values) 3.93 (s, 3H), 8.00 (d, 1H), 8.02 (d, 1H), 8.33 (d, 2H), 8.42 (s, 1H), 8.45 (s, 1H), 8.61 (m, 2H), 8.76 (m, 2H).
II	Morpholine	RT/18hrs/NaI	274	m/e 518 (M+H) ⁺	(d-6-DMSO, d values) 2.31 (m, 2H), 3.28 (m, 2H), 3.4-3.6 (m, 4H (under H ₂ O peak)), 3.83 (m, 2H), 3.92 (m, 2H), 3.99 (s, 3H), 4.36 (m, 2H), 7.26 (d, 1H), 7.32 (d, 1H), 7.46 (m, 2H), 7.55 (m, 3H), 8.41 (s, 1H), 8.95 (s, 1H).
II	N-methylpiperidine	RT/18hrs/NaI	275	m/e 531 (M ⁺ +H)	(d-6-DMSO, d values) 2.25 (m, 2H), 2.83 (s, 3H), 3.2-3.7 (m, 10H (under H ₂ O peak)), 3.99 (s, 3H), 4.32 (m, 2H), 7.25 (d, 1H), 7.33 (d, 1H), 7.50 (m, 5H), 8.26 (bs, 1H), 8.92 (s, 1H).
II	pyrrolidine	RT/18hrs/NaI	276	m/e 531 (M ⁺ +H)	(d-6-DMSO, d values) 1.86 (m, 2H), 2.02 (m, 2H), 2.25 (m, 2H), 3.26 (m, 2H), 3.58 (m, 2H), 3.75 (m, 2H), 3.97 (s, 3H), 4.31 (m, 2H), 7.26 (d, 1H), 7.33 (d, 1H), 7.47 (m, 3H), 7.55 (m, 2H), 8.28 (s, 1H), 9.0 (s, 1H).
II	piperidine	RT/18hrs/NaI	277	m/e 516 (M ⁺ +H)	(d-6-DMSO, d values) 1.53 (m, 1H), 1.61 (m, 4H), 1.80 (m, 1H), 2.23 (m, 2H), 2.97 (m, 4H), 3.21 (m, 2H), 3.99 (s, 3H), 4.28 (m, 2H), 7.28 (d, 1H), 7.33 (d, 1H), 7.40 (s, 1H), 7.47 (m, 4H), 8.11 (s, 1H), 8.88 (s, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
I2	morpholine	RT/18hrs/NaI	278	m/e 504 (M ⁺ +H)	(d-6-DMSO, d values) 3.26 (m, 2H), 3.42-3.7 (m, 4H (under H ₂ O peak)), 3.83 (m, 2H), 3.95 (m, 2H), 4.01 (s, 3H), 4.73 (m, 2H), 7.28 (d, 1H), 7.33 (d, 1H), 7.47 (m, 2H), 7.54 (m, 3H), 8.47 (s, 1H), 8.97 (s, 1H).
I2	N-methyl piperidine	RT/36hrs/NaI	279	m/e 531 (M ⁺ +H)	(d-6-DMSO, d values) 3.26 (m, 2H), 3.42-3.7 (m, 4H (under H ₂ O peak)), 3.83 (m, 2H), 3.95 (m, 2H), 4.01 (s, 3H), 4.73 (m, 2H), 7.28 (d, 1H), 7.33 (d, 1H), 7.47 (m, 2H), 7.54 (m, 3H), 8.47 (s, 1H), 8.97 (s, 1H).
I2	piperidine	RT/36hrs/NaI	280	m/e 502 (M ⁺ +H)	(d-6-DMSO, d values) 1.53 (m, 1H), 1.64 (m, 4H), 1.80 (m, 1H), 3.01 (m, 4H), 3.4-3.6 (m, 2H (under H ₂ O peak)), 4.02 (s, 3H), 4.61 (m, 2H), 7.26 (d, 1H), 7.32 (d, 1H), 7.44 (m, 2H), 7.50 (m, 3H), 8.26 (s, 1H), 8.92 (s, 1H).
I2	dimethyl amine	RT/36hrs/NaI/ ethanol	281	m/e 462 (M ⁺ +H)	(d-6-DMSO, d values) 2.91 (d, 6H), 3.5-3.7 (m, 2H (under H ₂ O peak)), 4.00 (s, 3H), 4.68 (m, 2H), 7.28 (d, 1H), 7.33 (d, 1H), 7.46 (m, 2H), 7.54 (m, 3H), 8.53 (s, 1H), 8.95 (s, 1H).
300		75°C/2hrs/ thioanisole/ TFA	282	m/e 391 (M ⁺ +H)	(d-6-DMSO, d values) 3.90 (s, 3H), 7.21 (d, 2H), 7.30 (m, 3H), 7.37 (m, 2H), 7.69 (s, 1H), 8.40 (s, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
I3	morpholine	RT/18hrs/NaI	283	m/e 488 (M ⁺ +H)	(d-6-DMSO, d values) 2.31 (m, 2H), 3.08 (m, 2H), 3.29 (m, 2H), 3.35 (m, 2H), 3.81 (m, 2H), 3.95 (m, 2H), 4.01 (s, 3H), 4.31 (m, 2H), 7.26 (d, 1H), 7.33 (d, 1H), 7.47 (m, 2H), 7.54 (m, 3H), 8.22 (s, 1H), 8.94 (s, 1H).
I3	N-methyl piperazine	RT/18hrs/NaI	284	m/e 531 (M ⁺ +H)	(d-6-DMSO, d values) 2.34 (m, 2H), 2.84 (bs, 3H), 3.25-3.8 (m, 10H (under H ₂ O peak)), 4.02 (s, 3H), 4.31 (m, 2H), 7.26 (d, 1H), 7.33 (d, 1H), 7.47 (m, 2H), 7.55 (m, 3H), 8.26 (s, 1H), 8.96 (s, 1H).
282		RT/18hr/DMA KOtBu/18- crown-6	285	m/e 488 (M ⁺ +H)	(d-6-DMSO, d values) 1.95 (m, 2H), 2.10-2.4 (m, 3H), 3.99 (s, 3H), 4.15 (m, 2H), 7.27 (m, 1H), 7.35 (d, 1H), 7.52 (m, 4H), 7.80 (s, 1H), 8.08 (s, 1H), 8.98 (s, 1H).
I3	dimethyl amine	50°C/18hrs/ NaI/ ethanol	286	m/e 476 (M ⁺ +H)	(d-6-DMSO, d values) 2.28 (m, 2H), 2.82 (m, 6H), 3.24 (m, 2H), 3.97 (s, 3H), 4.28 (m, 2H), 7.28 (d, 1H), 7.34 (d, 1H), 7.45 (s, 1H), 7.50 (m, 4H), 8.09 (s, 1H), 8.88 (s, 1H), 9.95 (bs, 1H).
288		RT/18hrs/NaO H/MeOH/ water	289	m/e 449 (M+H) ⁺	(d-6-DMSO, d values) 3.92 (s, 3H), 4.90 (s, 2H), 7.21 (m, 2H), 7.30 (d, 1H), 7.34 (m, 4H), 7.74 (s, 1H), 8.45 (s, 1H), 9.51 (bs, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
289	methylamine	RT/18hrs/ THF/EDC/DM AP/DCM	291	m/e 462 (M ⁺ +H) ⁺	(d-6-DMSO, d values) 2.66 (d, 3H), 3.99 (s, 3H), 4.74 (s, 2H), 7.26 (m, 2H), 7.31 (d, 1H), 7.45 (m, 4H), 7.97 (s, 1H), 8.06 (bs, 1H), 8.76 (s, 1H).
301		75°C/2hrs/ Et ₃ SiH/ TFA	293	m/e 476 (M ⁺ +H)	(d-6-DMSO, d values) 4.03 (s, 3H), 7.26 (m, 2H), 7.32 (d, 1H), 7.45 (m, 4H), 7.50 (m, 1H), 8.81 (s, 1H).
289	cyclopropyl- amine	RT/1 week/EDC/ DMAP/DMA	302	m/e 488 (M ⁺ +H)	d-6-DMSO, d values) 0.47 (m, 2H), 0.64 (m, 2H), 2.70 (m, 1H), 3.97 (s, 3H), 4.68 (s, 2H), 7.26 (m, 2H), 7.32 (m, 1H), 7.46 (m, 4H), 8.03 (s, 1H), 8.29 (m, 1H), 8.84 (s, 1H).
13	cyclopropyl- amine	RT/18hr/NaI	319	m/e 488 (M ⁺ +H)	(d-6-DMSO, d values) 1.13 (d, 6H), 2.34 (m, 2H), 2.56 (d, 3H), 2.61 (m, 2H), 3.24 (m, 2H), 3.50 (m, 2H), 3.58 (s, 2H), 3.98 (m, 5H), 4.29 (2H, m), 6.20 (m, 1H), 6.26 (m, 1H), 6.33 (m, 1H), 7.05 (m, 3H), 7.45 (d, 2H), 7.54 (s, 1H), 7.81 (m, 1H), 8.26 (s, 1H), 8.93 (s, 1H).
13	dimethyl- morpholine	RT/18hr/NaI	260	m/e 546 (M ⁺ +H)	(d-6-DMSO, d values) 1.13 (d, 6H), 2.31 (m, 2H), 2.66 (m, 2H), 3.24 (m, 2H), 3.97 (bs, 5H), 4.28 (m, 2H), 7.26 (d, 1H), 7.32 (d, 1H), 7.49 (m, 5H), 8.13 (s, 1H), 8.89 (s, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
I19	2,6-dimethylmorpholine	5 days	448		(d-6-DMSO, δ values) 1.02 (d, 6H), 1.58 (t, 2H), 1.94 (t, 3H), 2.42 (m, 2H), 2.56 (d, 3H), 2.75 (d, 2H), 3.53 (m, 2H), 3.69 (d, 2H), 3.91 (s, 3H), 4.17 (t, 2H), 4.53 (s, 2H), 7.0 (m, 4H), 7.11 (m, 2H), 7.22 (s, 1H), 7.28 (d, 2H), 7.74 (m, 2H), 7.88 (t, 1H), 8.35 (s, 1H), 9.4 (s, 1H).
I22	dimethylmorpholine	RT/72hr/NaI	449	m/e 666.5 (M^+ +H)	(d-6-DMSO d-4-Acetic, δ values) 0.20 (m, 2H), 0.43 (m, 2H), 0.96 (m, 1H), 1.17 (s, 3H), 1.19 (s, 3H), 2.32 - 2.42 (m, 2H), 2.60 (m, 2H), 3.04 (d, 2H), 3.20 (t, 2H), 3.55 (d, 2H), 3.93 - 4.15 (m, 5H), 4.34 (t, 2H), 4.48 (s, 2H), 6.62 - 6.70 (m, 2H), 7.19 (d, 2H), 7.32 (t, 1H), 7.47 - 7.53 (m, 3H), 8.14 (s, 1H), 8.88 (s, 1H).
I21	dimethylmorpholine	RT/72hr/NaI	450	m/e 666.5 (M^+ +H)	(d-6-DMSO d-4-Acetic, δ values) 1.14 (s, 3H), 1.16 (s, 3H), 1.62 (m, 2H), 1.96 (m, 2H), 2.12 (m, 2H), 2.34 (m, 2H), 2.67 (t, 2H), 3.27 (t, 2H), 3.50 (d, 2H), 3.89 - 4.01 (m, 5H), 4.18 - 4.26 (m, 1H), 4.30 (t, 2H), 4.39 (s, 2H), 6.59 - 6.65 (m, 2H), 6.72 (dd, 1H), 7.16 (d, 2H), 7.27 (t, 1H), 7.45 - 7.52 (m, 3H), 8.14 (s, 1H), 8.89 (s, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
I22	pyrrolidine	RT/48hr/NaI	451	m/e 622.5 (M ⁺ +H)	(d-6-DMSO d-4-Acetic, δ values) 0.19 (m, 2H), 0.41 (m, 2H), 0.95 (m, 1H), 1.88 - 2.10 (m, 2H), 2.15 - 2.36 (m, 2H), 3.02 (d, 2H), 3.07 - 3.14 (m, 2H), 3.34 (t, 2H), 3.61 (m, 2H), 4.02 (s, 3H), 4.33 (t, 2H), 4.47 (s, 2H), 6.62 - 6.70 (dd, 1H), 7.18 (d, 2H), 7.31 (t, 1H), 7.46 - 7.56 (m, 3H), 8.24 (s, 1H), 8.89 (s, 1H).
I21	pyrrolidine	RT/48hr/NaI	452	m/e 622.5 (M ⁺ +H)	(d-6-DMSO d-4-Acetic, δ values) 1.60 (m, 2H), 1.84 - 2.03 (m, 6H), 2.13 (m, 2H), 2.29 (m, 2H), 3.05 (m, 2H), 3.30 (t, 2H), 3.56 (m, 2H), 4.00 (s, 3H), 4.19 - 4.26 (m, 1H), 4.30 (t, 2H), 4.39 (s, 2H), 6.59 - 6.63 (m, 2H), 6.71 (d, 1H), 7.14 (d, 2H), 7.28 (t, 1H), 7.46 (d, 2H), 7.52 (s, 1H), 8.18 (s, 1H), 8.81 (s, 1H).
I23	pyrrolidine	RT/48hr/NaI	453	m/e 608.5 (M ⁺ +H)	(d-6-DMSO d-4-Acetic, δ values) 0.46 (m, 2H), 0.60 (m, 2H), 1.83 - 2.06 (m, 4H), 2.28 (m, 2H), 2.66 (m, 1H), 2.95 - 3.05 (m, 2H), 3.30 (t, 2H), 3.56 (m, 2H), 3.99 (s, 3H), 4.28 (t, 2H), 4.39 (s, 2H), 6.58 - 6.62 (m, 2H), 6.68 (dd, 1H), 7.13 (d, 2H), 7.26 (t, 1H), 7.47 (d, 2H), 7.54 (s, 1H), 8.22 (s, 1H), 8.87 (s, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
I23	dimethylmorpholine	RT/72hr/NaI	454	m/e 652.5 (M ⁺ +H)	(d-6-DMSO d-4-Acetic, δ values) 0.46 (m, 2H), 0.61 (m, 2H), 1.11 (s, 3H), 1.14 (s, 3H), 2.34 (m, 2H), 2.59 - 2.72 (m, 3H), 3.26 (t, 2H), 3.50 (d, 2H), 3.89 - 4.01 (m, 5H), 4.30 (t, 2H), 4.39 (s, 2H), 6.58 - 6.62 (m, 2H), 6.769(d, 1H), 7.15 (d, 2H), 7.27 (t, 1H), 7.44 - 7.50 (m, 3H), 8.14 (s, 1H), 8.90 (s, 1H).
I25	dimethylmorpholine	RT/72hr/NaI	455	m/e 596.5 (M ⁺ +H)	(d-6-DMSO d-4-Acetic, δ values) 1.13 (s, 3H), 1.15 (s, 3H), 2.32 (m, 2H), 2.65 (t, 2H), 2.75 (s, 3H), 3.26 (m, 2H), 3.50 (d, 2H), 3.89 - 3.95 (m, 2H), 3.96 (s, 3H), 4.28 (t, 2H), 7.15 (d, 3H), 7.40 - 7.49 (m, 5H), 7.59 (d, 1H), 8.08 (s, 1H), 8.80 (s, 1H).
I25	pyrrolidine	RT/48hr/NaI	456	m/e 552.5 (M ⁺ +H)	(d-6-DMSO d-4-Acetic, δ values) 1.81 - 2.05 (m, 4H), 2.28 (m, 2H), 2.76 (s, 3H), 3.04 (m, 2H), 3.31 (t, 2H), 3.57 (m, 2H), 3.99 (s, 3H), 4.30 (t, 2H), 7.12 - 7.20 (m, 3H), 7.42 - 7.52 (m, 5H), 7.59 (d, 1H), 8.16 (s, 1H), 8.94 (s, 1H).
I26	dimethylmorpholine	RT/72hr/NaI	457	m/e 570.5 (M ⁺ +H)	(d-6-DMSO d-4-Acetic, δ values) 1.12 (s, 3H), 1.15 (s, 3H), 2.34 (m, 2H), 2.66 (t, 2H), 3.25 (t, 2H), 3.51 (d, 2H), 3.72 (s, 3H), 3.91 - 3.99 (m, 2H), 4.00 (s, 3H), 4.30 (t, 2H), 6.69 (m, 2H), 6.77 (dd, 1H), 7.19 (d, 1H), 7.30 (t, 1H), 7.50 (s, 1H), 7.97 (dd, 1H), 8.21 (s, 1H), 8.32 (d, 1H), 8.92 (s, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
127	morpholine	RT/18hr/NaI	458	m/e 623 (M ⁺ +H)	(d-6-DMSO, d values) 0.39 (m, 2H), 0.59 (m, 2H), 2.33 (m, 2H), 2.63 (m, 1H), 3.28 (m, 2H), 3.49 (m, 2H), 3.56 (s, 2H), 3.82 (2H, m), 3.94 (m, 2H), 3.99 (s, 3H), 4.30 (2H, m), 6.20 (m, 1H), 6.26 (m, 1H), 6.34 (m, 1H), 7.07 (m, 4H), 7.44 (d, 2H), 7.52 (s, 1H), 8.20 (m, 1H), 8.92 (s, 1H).
127	dimethylmorpholine	RT/18hr/NaI	459	m/e 651 (M ⁺ +H)	(d-6-DMSO, d values) 0.39 (m, 2H), 0.59 (m, 2H), 1.13 (d, 6H), 2.36 (m, 2H), 2.65 (m, 3H), 3.26 (m, 2H), 3.53 (m, 4H), 3.99 (5H, m), 4.31 (m, 2H), 6.20 (m, 1H), 6.27 (m, 1H), 6.35 (m, 1H), 7.07 (m, 3H), 7.45 (d, 2H), 7.52 (s, 1H), 8.18 (m, 1H), 8.97 (s, 1H).
127	pyrrolidine	RT/18hr/NaI	460	m/e 607 (M ⁺ +H)	(d-6-DMSO, d values) 0.38 (m, 2H), 0.60 (m, 2H), 1.89 (m, 2H), 2.01 (m, 2H), 2.37 (m, 2H), 2.64 (m, 1H), 3.03 (m, 2H), 3.31 (m, 2H), 3.57 (m, 4H), 4.00 (s, 3H), 4.30 (m, 2H), 6.21 (m, 1H), 6.27 (m, 1H), 6.34 (m, 1H), 7.08 (m, 3H), 7.46 (d, 2H), 7.52 (s, 1H), 7.96 (m, 1H), 8.21 (s, 1H), 8.94 (s, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
I28	morpholine	RT/18hr/NaI	461	m/e 637 (M ⁺ +H)	(d-6-DMSO, d values) 0.20 (m, 2H), 0.45 (m, 2H), 0.96 (m, 1H), 2.42 (m, 2H), 3.17 (m, 2H), 3.37 (m, 2H), 3.57 (m, 2H), 3.70 (s, 2H), 3.91 (m, 2H), 4.07 (m, 5H), 4.40 (2H, m), 6.30 (m, 1H), 6.38 (m, 1H), 6.46 (m, 1H), 7.14 (m, 3H), 7.53 (d, 2H), 7.61 (s, 1H), 8.01 (m, 1H), 8.30 (s, 1H), 9.01 (s, 1H).
I28	dimethylmorpholine	RT/18hr/NaI	462	m/e 665 (M ⁺ +H)	(d-6-DMSO, d values) 0.17 (m, 2H), 0.41 (m, 2H), 0.93 (m, 1H), 1.20 (d, 6H), 2.42 (m, 2H), 2.71 (m, 2H), 3.30 (m, 2H), 3.56 (m, 2H), 3.66 (s, 2H), 3.80 (m, 2H), 4.05 (m, 5H), 4.37 (2H, m), 6.27 (m, 1H), 6.34 (m, 1H), 6.42 (m, 1H), 7.15 (m, 3H), 7.51 (d, 2H), 7.58 (s, 1H), 7.97 (m, 1H), 8.27 (s, 1H), 8.98 (s, 1H).
I28	pyrrolidine	RT/18hr/NaI	463	m/e 621 (M ⁺ +H)	(d-6-DMSO, d values) 0.11 (m, 2H), 0.36 (m, 2H), 0.87 (m, 1H), 1.87 (m, 2H), 2.00 (m, 2H), 2.29 (m, 2H), 2.96 (m, 2H), 3.02 (m, 2H), 3.31 (m, 2H), 3.56 (m, 2H), 3.61 (s, 2H), 4.00 (s, 3H), 4.29 (2H, m), 6.23 (m, 1H), 6.30 (m, 1H), 6.38 (m, 1H), 7.11 (m, 3H), 7.45 (d, 2H), 7.56 (s, 1H), 7.95 (m, 1H), 8.28 (s, 1H), 8.96 (s, 1H).



Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
I29	morpholine	RT/18hr/NaI	464	m/e 597 (M ⁺ +H)	(d-6-DMSO, d values) 2.31 (m, 2H), 3.26 (m, 2H), 3.46 (m, 2H), 3.58 (s, 2H), 3.85 (m, 2H), 3.95 (m, 2H), 4.00 (s, 3H), 4.26 (2H, m), 6.21 (m, 1H), 6.26 (m, 1H), 6.34 (m, 1H), 7.08 (m, 3H), 7.45 (d, 2H), 7.58 (s, 1H), 7.81 (m, 1H), 8.30 (s, 1H), 8.93 (s, 1H).
I29	dimethylmorpholine	RT/18hr/NaI	465	m/e 625 (M ⁺ +H)	(d-6-DMSO, d values) 1.13 (d, 6H), 2.34 (m, 2H), 2.56 (d, 3H), 2.61 (m, 2H), 3.24 (m, 2H), 3.50 (m, 2H), 3.58 (s, 2H), 3.98 (m, 5H), 4.29 (2H, m), 6.20 (m, 1H), 6.26 (m, 1H), 6.33 (m, 1H), 7.05 (m, 3H), 7.45 (d, 2H), 7.54 (s, 1H), 7.81 (m, 1H), 8.26 (s, 1H), 8.93 (s, 1H).
I30	dimethylmorpholine	RT/4 days/NaI	466	m/e 651.6 (M ⁺ +H)	(d-6-DMSO, (d-4Acetic) d values) 0.45 (m, 2H), 0.64 (m, 2H), 1.17 (d, 6H), 2.37 (m, 2H), 2.68 (m, 3H), 3.29 (t, 2H), 3.54 (d, 2H), 4.01 (m, 5H), 4.33 (t, 3H), 4.46 (s, 2H), 7.05 (m, 5H), 7.18 (m, 1H), 7.45 (d, 2H), 7.51 (s, 1H), 8.17 (m, 1H), 8.95 (s, 1H).
I31	morpholine	RT/18hr/NaI	467	m/e 626.4 (M ⁺ +H)	(d-6-DMSO, d values) 1.05 (d, 6H), 2.33 (m, 2H), 3.09 (m, 2H), 3.29 (m, 2H), 3.47 (m, 2H), 3.84 (m, 3H), 3.97 (m, 5H), 4.28 (t, 2H), 4.42 (s, 2H), 7.07 (m, 6H), 7.42 (m, 4H), 8.10 (s, 1H), 8.84 (s, 1H).

Start Comp	Reagent	Conditions	Prod	Mass spec.	Nmr
I30	pyrrolidine	RT/18hr/Nal	468	m/e 608.6 (M ⁺ +H)	(d-6-DMSO, d values) 0.47 (m, 2H), 0.67 (m, 2H), 1.92 (m, 2H), 2.05 (m, 2H), 2.30 (m, 2H), 2.67 (m, 1H), 3.06 (m, 2H), 3.34 (m, 2H), 3.59 (m, 2H), 4.03 (s, 3H), 4.32 (t, 2H), 4.47 (s, 2H), 7.06 (m, 5H), 7.19 (m, 1H), 7.46 (d, 2H), 7.55 (s, 1H), 7.83 (m, 1H), 8.19 (s, 1H), 8.92 (s, 1H).
I32	dimethylmorpholine	RT/96hr/Nal	469	m/e 626.5 (M ⁺ +H)	(d-6-DMSO(d4-Acetic), d values) 1.16 (d, 6H), 2.37 (m, 2H), 2.63 (s, 3H), 2.70 (m, 2H), 3.29 (m, 2H), 3.56 (d, 2H), 3.99 (m, 5H), 4.30 (t, 2H), 4.47 (s, 2H), 7.09 (m, 6H), 7.44 (m, 3H), 8.16 (s, 1H), 8.98 (s, 1H).
I32	pyrrolidine	RT/96hr/Nal	470	m/e 582.5 (M ⁺ +H)	(d-6-DMSO(d4-Acetic), d values) 1.16 (d, 6H), 2.37 (m, 2H), 2.63 (s, 3H), 2.70 (m, 2H), 3.29 (m, 2H), 3.56 (d, 2H), 3.99 (m, 5H), 4.30 (t, 2H), 4.47 (s, 2H), 7.09 (m, 6H), 7.44 (m, 3H), 8.16 (s, 1H), 8.98 (s, 1H).
I34	Dimethylmorpholine	RT/48hr/Nal	481	m/e 626.6 (M ⁺ +H) ⁺	(d-6-DMSO, δ values) 1.00 (s, 3H), 1.04 (s, 3H), 1.56 (t, 2H), 1.95 (m, 2H), 2.42 (t, 2H), 2.64 (d, 3H), 2.76 (d, 2H), 3.55 (m, 2H), 3.90 (s, 3H), 4.19 (t, 2H), 4.41 (s, 2H), 6.56 - 6.62 (m, 2H), 6.70 (d, 1H), 7.09 (d, 2H), 7.22 - 7.37 (m, 4H), 7.28 (s, 1H), 8.00 (bs, 1H), 8.40 (s, 1H), 9.50 (s, 1H).

Intermediate Table 9


Start No.	Reagent	Conditions	Int.	Mass spec	structure
273	dichloro propane	70°C/2hr//KOt Bu/DMA	I1	m/e 467, 469 (M+H) ⁺ .	
273	dichloro -ethane	70°C/2hr//KOt Bu/DMA	I2	m/e 453,455 (M ⁺ +H)	
282	bromo chloro propane	RT/18hrs/ /KOtBu/18-C-6/DMA	I3	m/e 467, 469 (M+H) ⁺ .	
26	3-bromo-1-propanol	RT/18hr/ PPh ₃ / DEAD/THF	I8	m/e 533,535 (M+H) ⁺ .	
26	1,3-dichloro -propane	70°C/4hr/ KOtBu/DMA	I9	m/e 490, 492 (M+H) ⁺ .	
26	dichloro -ethane	85°C/4hr/ KOtBu/DMA	I10	m/e 476,478 (M+H) ⁺ .	

Start No.	Reagent	Conditions	Int.	Mass spec	structure
109	1-Bromo-3-chloro-propane	RT/ nBu ₄ NI/ 18crown6	I11	nmr obtained	
108	1-Bromo-3-chloro-propane	RT/ nBu ₄ NI/ 18crown6	I12	nmr obtained	
126	1-Bromo-3-chloro-propane	RT/ nBu ₄ NI/ 18crown6	I13	nmr obtained	
123	1-Bromo-3-chloro propane	RT/ nBu ₄ NI/ DMA 18crown6/18h	I14	m/e 520 (M+H) ⁺	
125	1-Bromo-3-chloro-propane	RT/ nBu ₄ NI/ DMA 18crown6/ 8hr	I15	m/e 520 (M+H) ⁺	
220	1-Bromo-3-chloro-propane	RT/15min/ KOtBu/DMA then RT/16hr/ /nBu ₄ NI/18-Crown-6	I16	nmr available	

Start No.	Reagent	Conditions	Int.	Mass spec	structure
221	1-Bromo-3-chloro-propane	RT/15min/ KOTBu/DMA then RT/16hr /nBu ₄ NI 18-Crown-6	I17	nmr available	
27	1-chloro-3-bromo-propane	RT/18hr/ KO ^t Bu(1.0M in THF) / DMSO	I18	m/e 490 (M ⁺ +H)	

Example 7

In the above Table I4 is a compound of structure



- 5 which had been prepared by a method analogous to that described in Example 1, but using reaction conditions of 100°C/2hr/1-PrOH.





Mass Spectrum m/e 577.45, 579.46 (M⁺+H).






- 10 NMR Spectrum (d-6-DMSO, d values) 2.28 (m, 2H), 3.16 (q, 2H), 3.4 (t, 2H), 3.82 (t, 2H), 3.98 (s, 3H), 4.3 (t, 2H), 4.48(s, 2H), 6.95-7.22 (m, 6H), 7.4 (d, 2H), 7.46 (s, 1H), 7.6 (t, 1H), 8.09 (s, 1H), 8.9 (s, 1H), 11.07 (br.s, 1H).







The chloropropoxyquinoline intermediate (Mass Spectrum m/e 311.2 (M+H)⁺) was prepared by reacting the corresponding hydroxy quinoline with 1-bromo-3-chloropropane at room temperature for 16hr in the presence of nBu₄NI/18-crown-6

- 15 The following haloalkoxy quinolines were prepared by analogous routes:




Table 10

I No.	reaction conditions	mass spec.	structure
I5	100°C/18hr s/n-PrOH	m/e 548.5 (M+H) ⁺	
I6			
I19	100°C/2hr/1 -PrOH	m/e 604.44 (M ⁺ +H).	
I20	100°C/3.5hr /1-PrOH	m/e 604.44 (M ⁺ +H)	


I No.	reaction conditions	mass spec.	structure
I21	100°C/3.5hr /1-PrOH	m/e 587.5 (M ⁺ +H)	
I22	100°C/2hr/1 -PrOH	m/e 587.5 (M ⁺ +H)	
I23	100°C/2hr/1 -PrOH	m/e 573.4 (M ⁺ +H)	
I24	100°C/3.5hr /1-PrOH	m/e 574.4 (M ⁺ +H)	
I25	100°C/3.5hr /1-PrOH	m/e 517.3 (M ⁺ +H)	

I No.	reaction conditions	mass spec.	structure
I26	100°C/2hr/1 -PrOH	m/e 570.5 (M ⁺ +H)	
I27	100°C/4hr/1 -PrOH	m/e 572, 574 (M ⁺ +H)	
I28	100°C/4hr/1 -PrOH	m/e 586, 588 (M ⁺ +H)	
I29	100°C/4hr/1 -PrOH	m/e 546, 548 (M ⁺ +H)	
I30	100°C/18hr/ 1-PrOH	m/e 573.5 (M ⁺ +H)	
I31	100°C/18hr/ 1-PrOH	m/e 575.5 (M ⁺ +H)	

160

I No.	reaction conditions	mass spec.	structure
I32	100°C/18hr/ 1-PrOH	m/e 547.5 (M ⁺ +H)	
I33	RT/15min/ NaH/DMA then RT/2hr/(2)		
I34	100°C/2hr/ n-PrOH		

In addition I5 was converted to I7



(17)

5


using the following reaction conditions: RT/3hrs/LiOH.H₂O/MeOH/H₂O

Mass Spectrum m/e 534.5 (M+H)⁺

NMR Spectrum (d-6-DMSO, d values) 2.26 (m, 2H), 3.82 (m, 2H), 3.93 (s, 3H), 4.26 (t, 2H), 4.68 (s, 2H), 7.04 (m, 6H), 7.29 (m, 2H), 7.39 (s, 1H), 7.93 (s, 1H), 8.55 (s, 1H).

Example 8Preparation of Compound No. 312

- 5 In this example, an intermediate nitro compound of formula (2) was reacted in situ with a chloroquinoline intermediate to produce compound 312, (a compound of formula (I)) directly in accordance with the following scheme:



- 10 The reaction conditions were: Cyclohexene, 1-propanol, Pd/C, filter then add quinoline to obtain the desired product

Mass Spectrum m/e 452 ($M^+ + H$)

NMR Spectrum ($CDCl_3$, d values) 2.70 (m 2H), 3.15 (m 2H), 3.75 (s, 3H), 4.00 (s, 3H), 6.70 (d, 1H), 6.80 (broad s, 1H), 6.95 (s, 1H), 7.05 (d, 2H), 7.15 (d, 2H), 7.15 (m, 1H),

- 15 7.35 (s, 1H), 7.45 (t, 1H), 8.60 (s, 1H).

Quinoline SM: WO 9843960

The reaction conditions used to obtain Intermediate labelled (2) was KOtBu, DMA.

Mass Spectrum m/e 270 ($M^+ + H$)

20

Using an analogous method, the following compounds were also produced

Table 11

No.	Mass spec	N.M.R
313	m/e 429 (M ⁺ +H)	(CDCl ₃ , d values) 3.70 (s, 3H), 4.00 (s, 3H), 6.85 (broad s, 1H), 6.90 (m, 2H), 7.10 (d, 2H), 7.15 (d, 2H), 7.35 (m, 3H), 8.00 (s, 1H), 8.60 (s, 1H).
314	m/e 453 (M ⁺ +H)	(d-6-DMSO@373K, d values) 3.60 (s, 3H), 3.95 (s, 3H), 4.00 (s, 3H), 6.90 (d, 1H), 7.15 (d, 2H), 7.25 (t, 1H), 7.40 (m, 3H), 7.45 (s, 1H), 8.00 (s, 1H), 8.70 (s, 1H).
315	m/e 438 (M ⁺ +H)	(d-6-DMSO, d values) 4.00 (s, 3H), 4.00 (s, 3H), 6.75 (d, 1H), 6.85 (d, 1H), 7.20 (d, 2H), 7.30 (t, 1H), 7.40 (d, 1H), 7.50 (d, 2H), 7.50 (s, 1H), 7.95 (d, 1H), 8.20 (s, 1H), 8.95 (s, 1H), 11.30 (broad s, 1H).

Example 9Preparation of Compounds 136 and 140 in Table 1

- 5 Compound 85 prepared as described above, was dissolved in trichloromethane and reacted with oxone in the presence of wet alumina to yield the title compounds.

Compound 136

Mass Spectrum m/e 460 (M⁺+H)

- 10 NMR Spectrum (d-6-DMSO, d values) 2.80 (s, 3H), 3.90 (s, 3H), 3.95 (s, 3H), 6.85 (d, 1H), 7.20 (d, 2H), 7.35 (m, 4H), 7.45 (m, 1H), 7.75 (m, 2H), 8.40 (s, 1H), 9.55 (broad s, 1H).

Compound 140

Mass spec m/e 476 (M⁺+H)

- 15 NMR Spectrum (d-6-DMSO, d values) 3.40 (s, 3H), 3.95 (s, 3H), 4.00 (s, 3H), 6.95 (d, 1H), 7.20 (d, 2H), 7.35 (m, 2H), 7.40 (d, 2H), 7.65 (m, 1H), 7.80 (s, 1H), 7.90 (dd, 1H), 8.45 (s, 1H), 9.65 (broad s, 1H).

Example 10

Preparation of Compound 168 in Table 1


Compound 173 in Table 1 was reacted with methylamine for 18 hours at room temperature in the presence of HCl, EDC, NMM and DCM to yield the desired amide.

Mass spec. m/e 582 ($M+H$)⁺.

- 5 NMR Spectrum (d-6-DMSO, δ values) 2.33 (m, 2H), 2.55 (d, 3H), 3.12 (m, 2H), 3.22-3.45 (m, 4H (under H₂O signal)), 3.43 (s, 2H), 3.78 (m, 2H), 3.97 (m, 5H), 4.28 (m, 2H), 6.83 (d, 1H), 7.05 (d, 2H), 7.10 (m, 1H), 7.21 (m, 1H), 7.33 (m, 1H), 7.41 (d, 2H), 7.47 (s, 1H), 7.75 (m, 1H), 8.12 (s, 1H), 8.81 (s, 1H).

10 Example 11Preparation of Compound 301 in Table 3

This compound was prepared using the following scheme:



- 15 Reaction conditions: 100°C/4hrs/NEt₃/Diphenylphosphorylazide/t-BuOH

Chromatography: yes

Mass Spectrum m/e 490 ($M+H$)⁺.

NMR Spectrum (d-6-DMSO, δ values) 1.48 (s, 9H), 4.01 (s, 3H), 7.26 (d, 1H), 7.33 (d, 1H), 7.45 (m, 1H), 7.49 (m, 2H), 7.53 (d, 2H), 8.70 (s, 1H), 8.82 (s, 1H), 8.97 (s, 1H).

- 20 Intermediate (3)

Reaction conditions: 100°C/18hrs/n-PrOH

Mass Spectrum m/e 433 (M+H)⁺.

Intermediate (4)

Reaction conditions: RT/36hrs/LiOH/MeOH/water

Mass Spectrum m/e 418 (M+H)⁺.

5

Example 12

Preparation of Compound 183 in Table 1

Intermediate I7 in Table 1 was reacted with cyclopropylamine and N-methylmorpholine at room temperature for 48hours in the presence of DMAP, EDC and DCM to yield the
10 desired product.


Mass Spectrum m/e 624.5 (M+H)⁺

NMR Spectrum (d-6-DMSO, d values) 0.42 (m, 2H), 0.61 (m, 2H), 2.30 (m, 2H), 2.63 (m, 1H), 3.11 (m, 2H), 3.35 (2H under H₂O peak), 3.49 (m, 2H), 3.79 (m, 2H), 3.97 (m, 5H), 4.30 (m, 2H), 7.08 (m, 7H), 7.40 (d, 2H), 7.45 (s, 1H), 7.78 (s, 1H), 8.08 (s, 1H),
15 8.84 (s, 1H).

Example 13

Preparation of Compound No 430 in Table 1

This compound was prepared using the following scheme:



100°C/18hrs/n-PrOH

Chromatography: yes

Mass Spectrum m/e 525 (M+H)⁺

- 5 NMR Spectrum (d-6-DMSO, d values) 0.182 (m, 2H), 0.41 (m, 2H), 0.94 (m, 1H), 3.02 (t, 2H), 4.00 (m, 6H), 4.52 (s, 2H), 7.14 (m, 6H), 7.47 (m, 3H), 7.70 (t, 1H), 8.16 (s, 1H), 8.94 (s, 1H).

The aniline starting material (1) was prepared as described above in relation to


- 10 Intermediate I5.

This was converted to Intermediate (2) above by reaction with cyclopropanemethylamine in methanol at room temperature for 18hrs.

Mass Spectrum m/e 313.5 (M+H)⁺

- 15 Example 14










Using a method analogous to that of Example 13, the R⁷ group was modified to form a different group R⁷ in the anilines used as starting materials in accordance with the following general scheme:



- 20

prior to conversion to the corresponding compound of formula (I) as summarised in the following Table 12.

Table 12

Starting aniline R ⁹⁰		R ⁹¹	Reagent/conditions	R ⁹²	Final aniline R ⁹³	Final Product
O(CH ₂) ₃ Br	H		RT/5days/cyclopropyl amine/NaI/MeOH		H	437
	H		RT/5days/cyclopropyl amine/NaI/MeOH		H	438
H			RT/5days/Me- amine/NaI/MeOH	H		439
	H		methylamine/ethanol		H	444
	H		methylamine/ethanol		H	445

167

Starting aniline		Reagent/conditions	Final aniline		Final Product
R ⁹⁰	R ⁹¹		R ⁹²	R ⁹³	
	H	cyclopropylamine/ ethanol		H	447

Example 15

In the preparation of other compounds of formula (I) the R⁷ group was modified to form a different group R⁷ in the nitrobenzyl compounds of formula (VII) used as starting materials in accordance with the following general scheme:


















Table 13

Starting 4-phenoxy nitrobenzene R ⁹⁴		R ⁹⁵	Reagent/conditions	Final 4-phenoxy nitrobenzene R ⁹⁶ R ⁹⁷		Final Product
NH ₂		H	3-bromopropionyl chloride, triethylamine, DMA; then dimethyl morpholine		H	433
NH ₂		H	3-bromopropionyl chloride, triethylamine, DMA; then piperidine		H	434
NH ₂		H	3-bromopropionyl chloride, triethylamine, DMA; then methylamine in methanol		H	435
NH ₂		H	3-bromopropionyl chloride, triethylamine, DMA; then dimethylamine in methanol		H	435

Starting 4-phenoxy nitrobenzene R ⁹⁴		Reagent/conditions	Final 4-phenoxy nitrobenzene R ⁹⁶		Final Product
R ⁹⁵			R ⁹⁷		
H	NH ₂	80°C/6hrs/ethylbromoacetate/NaOAc/EtOH	H		439
OCH ₂ COOH	H	EDC/DMAPI/HOBT/DMA 	H		441
OCH ₂ COOH	H	EDC/DMAPI/HOBT/DMA 	H		442
OCH ₂ COOH	H	EDC/DMAPI/HOBT/DMA 	H		443

170

Starting 4-phenoxy nitrobenzene R ⁹⁴		R ⁹⁵	Reagent/conditions	Final 4-phenoxy nitrobenzene R ⁹⁶		R ⁹⁷	Final Product
OCH ₂ COOH		H	EDC/DMAP/HOBT/DMA 		H		447* intermediat e(see also Ex 15)
O(CH ₂) ₂ NH ₂		H	RT/48hrs/Succinamic acid/EDC/DEAD/NMM/ DCM 	O(CH ₂) ₂ NHC(O)(CH ₂) ₂ - CN	H		472
O(CH ₂) ₂ NH ₂		H	RT/18hrs/acetylchloride/ DCM iPr ₂ N(CH ₂ CH ₃) ₂	O(CH ₂) ₂ NHC(O)CH ₃	H		474
O(CH ₂) ₂ NH ₂		H	RT/18hrs/ iPr ₂ N(CH ₂ CH ₃) ₂ /DCM allylchloroformate	O(CH ₂) ₂ NHC(O)OCH ₂ - CH=CH ₂	H		475
H		OCH ₂ C(O)OCH ₂ C H ₃	RT/2hrs/ methylamine/ MeOH	H	OCH ₂ C(O)NH- CH ₃		477

171

Starting 4-phenoxy nitrobenzene R ⁹⁴		R ⁹⁵	Reagent/conditions	Final 4-phenoxy nitrobenzene R ⁹⁶		R ⁹⁷	Final Product
H		OH	65°C/1.5hr/K ₂ CO ₃ /Ethyl/br omoacetate/Acetone	H		OCH ₂ C(O)O- CH ₂ CH ₃	477
H		OCH ₃	195°C/2hr/Pyridine.HCl	H		OH	477
OCH ₂ C(O)OH		H	RT/18hrs/isopropylamine/ EDC/DEAD/NMM/DCM	OCH ₂ C(O)NHCH(CH ₃) ₂		H	482

Biological Data

Assay for inhibitors of the MAP kinase pathway

To evaluate inhibitors of the MAPK pathway a coupled assay was carried out which measures phosphorylation of serine/threonine residues present in the substrate in the presence or absence of inhibitor. Recombinant glutathione S-transferase fusion protein containing human p45MEK1 (GST-MEK) was activated by c-raf (Sf9 insect cell lysate from triple baculoviral infection with c-raf/ras/lck) and used for the assay. Active GST-MEK was first used to activate a recombinant glutathione S-transferase fusion protein containing p44MAP kinase (GST-MAPK) in the presence of ATP and Mg^{2+} for 60min at room temperature in the presence or absence of potential inhibitors. The activated GST-MAPK was then incubated with myelin basic protein (MBP) as substrate for 10min at room temperature in the presence of ATP, Mg^{2+} and ^{33}P -ATP. The reaction was stopped by addition of 20% v/v phosphoric acid. Incorporation of ^{33}P into the myelin basic protein was determined by capture of the substrate on a filter mat, washing and counting using scintillation methods. The extent of inhibition was determined by comparison with untreated controls.

The final assay solution contained 10mM Tris, pH 7.5, 0.05mM EGTA, 8.33 μ M [γ - ^{33}P]ATP, 8.33mM Mg(OAc)₂, 0.5mM sodium orthovanadate, 0.05%w/v BSA, 6.5ng GST-MEK, 1 μ g GST-MAPK and 16.5 μ g MBP in a reaction volume of 60 μ l.

Compounds tested of the present invention had IC₅₀ results typically less than 0.5 μ M. For example, Compound No 252 gave an IC₅₀ of 0.15 μ M.

In vitro MAP kinase assay

To determine whether compounds were inhibiting GST-MEK or GST-MAPK, a direct assay of MAPK activity was employed. GST-MAPK was activated by a constitutively active GST-MEK fusion protein containing two point mutations (S217E, S221E) and used for the assay in the presence and absence of potential inhibitors. The activated GST-MAPK was incubated with substrate (MBP) for 60min at room temperature in the presence of ATP, Mg^{2+} and ^{33}P -ATP. The reaction was stopped by addition of 20% v/v phosphoric acid. Incorporation of ^{33}P into the myelin basic protein was determined by capture of the substrate on a filter mat, washing and counting using scintillation methods.

The final assay solution contained 12mM Tris, pH 7.5, 0.06mM EGTA, 30 μ M [γ ³³P]ATP, 10mM Mg(OAc)₂, 0.6mM sodium orthovanadate, 0.06%w/v BSA, 28ng GST-MAPK and 16.5 μ g MBP in a reaction volume of 60 μ l.


Compounds of the invention showed activity in this screen.

5 Cell proliferation assays

Cells were seeded into multi-well plates at 20 000 - 40 000 cells/ml in growth medium containing 5% FCS and incubated overnight at 37°C. The compounds were prepared in fresh medium at an appropriate concentration and added to the wells containing the cells. These were then incubated for a further 72 hours. Cells were then
10 either removed from the wells by incubating with trypsin/EDTA and counted using a Coulter counter, or treated with XTT/PMS in PBSA and optical densities read at 450nm. Compounds tested of the present invention had IC₅₀ results typically less than 30 μ M. For example, Compound No 250 gave an IC₅₀ of 7.76 mM in HT29 human colon tumour cells; Compound No 32 gave an IC₅₀ of 1.5 μ M in HT29 cells and an IC₅₀ of 0.6 μ M in
15 MC26 mouse colon tumour cells.

Claims

1. A compound of formula (I)



(I)

or a pharmaceutically acceptable salt thereof.

wherein:

n is 0-1;

X and Y are independently selected from -NH-, -O-, -S-, or -NR⁸- where R⁸ is alkyl of 1-6 carbon atoms and X may additionally comprise a CH₂ group;

R⁷ is a group (CH₂)_mR⁹ where m is 0, or an integer of from 1-3 and R⁹ is a substituted aryl group, an optionally substituted cycloalkyl ring of up to 10 carbon atoms, or an optionally substituted heterocyclic ring or an N-oxide of any nitrogen containing ring;

R⁶ is a divalent cycloalkyl of 3 to 7 carbon atoms, which may be optionally further

substituted with one or more alkyl of 1 to 6 carbon atom groups; or is a divalent pyridinyl, pyrimidinyl, or phenyl ring; wherein the pyridinyl, pyrimidinyl, or phenyl ring may be optionally further substituted with one or more groups selected from halogen,

alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, azido, hydroxyalkyl of 1-6 carbon atoms, halomethyl, alkoxymethyl of 2-7 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkoxy of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, hydroxy, trifluoromethyl, cyano, nitro, carboxy, carboalkoxy of 2-7 carbon atoms, carboalkyl of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzoyl, benzyl,

amino, alkylamino of 1-6 carbon atoms, dialkylamino of 2 to 12 carbon atoms, phenylamino, benzylamino, alkanoylamino of 1-6 carbon atoms, alkenoylamino of 3-8 carbon atoms, alkynoylamino of 3-8 carbon atoms, and benzoylamino;

- 5 R₁, R₂, R₃ and R₄ are each independently selected from hydrogen, hydroxy, halogeno, cyano, nitro, trifluoromethyl, C₁₋₃alkyl, -NR¹¹R¹² (wherein R¹¹ and R¹², which may be the same or different, each represents hydrogen or C₁₋₃alkyl), or a group R¹³-X¹-(CH₂)_x wherein x is 0 to 3, X¹ represents -O-, -CH₂-, -OCO-, carbonyl, -S-, -SO-, -SO₂-, -NR¹⁴CO-, -CONR¹⁵-, -SO₂NR¹⁶-, -NR¹⁷SO₂- or -NR¹⁸- (wherein R¹⁴, R¹⁵, R¹⁶, R¹⁷ and
- 10 R¹⁸ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R¹³ is selected from one of the following sixteen groups:
 - 1) C₁₋₅alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro and amino;
 - 2) C₁₋₅alkylX²COR¹⁹ (wherein X² represents -O- or -NR²⁰- (wherein R²⁰ represents
 - 15 hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R¹⁹ represents -NR²¹R²²- or -OR²³- (wherein R²¹, R²² and R²³ which may be the same or different each represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl));
 - 3) C₁₋₅alkylX³R²⁴ (wherein X³ represents -O-, -S-, -SO-, -SO₂-, -OCO-, -NR²⁵CO-, -CONR²⁶-, -SO₂NR²⁷-, -NR²⁸SO₂- or -NR²⁹- (wherein R²⁵, R²⁶, R²⁷, R²⁸ and R²⁹ each
 - 20 independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R²⁴ represents hydrogen, C₁₋₃alkyl, cyclopentyl, cyclohexyl or a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which C₁₋₃alkyl group may bear one or two substituents selected from oxo, hydroxy, halogeno and C₁₋₄alkoxy and which cyclic group may bear one or two substituents selected from oxo,
 - 25 hydroxy, halogeno, C₁₋₄alkyl, C₁₋₄hydroxyalkyl and C₁₋₄alkoxy);
 - 4) C₁₋₅alkylX⁴C₁₋₅alkylX⁵R³⁰ (wherein X⁴ and X⁵ which may be the same or different are each -O-, -S-, -SO-, -SO₂-, -NR³¹CO-, -CONR³²-, -SO₂NR³³-, -NR³⁴SO₂- or -NR³⁵- (wherein R³¹, R³², R³³, R³⁴ and R³⁵ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R³⁰ represents hydrogen or C₁₋₃alkyl);
 - 30 5) C₁₋₅alkylR³⁶ (wherein R³⁶ is a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which heterocyclic group

may bear one or two substituents selected from oxo, hydroxy, halogeno, C₁₋₄alkyl, C₁₋₄hydroxyalkyl and C₁₋₄alkoxy);

- 6) (CH₂)_qX⁶R³⁷ (wherein q is an integer from 0 to 5, X⁶ represents a direct bond, -O-, -S-, -SO-, -SO₂-, -NR³⁸CO-, -CONR³⁹-, -SO₂NR⁴⁰-, -NR⁴¹SO₂- or -NR⁴²- (wherein R³⁸, R³⁹, R⁴⁰, R⁴¹ and R⁴² each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R³⁷ is a phenyl group, a pyridone group or a 5 or 6 membered aromatic heterocyclic group with 1 to 3 heteroatoms selected from O, N and S, which phenyl, pyridone or aromatic heterocyclic group may carry up to 5 substituents selected from hydroxy, halogeno, amino, C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄hydroxyalkyl, C₁₋₄hydroxyalkoxy, C₁₋₄aminoalkyl, C₁₋₄alkylamino, carboxy, cyano, -CONR⁴³R⁴⁴ and -NR⁴⁵COR⁴⁶ (wherein R⁴³, R⁴⁴, R⁴⁵ and R⁴⁶, which may be the same or different, each represents hydrogen, C₁₋₄alkyl or C₁₋₃alkoxyC₂₋₃alkyl));
- 7) C₂₋₆alkenylR³⁶ (wherein R³⁶ is as defined hereinbefore);
- 8) C₂₋₆alkynylR³⁶ (wherein R³⁶ is as defined hereinbefore);
- 9) X⁷R⁴⁷ (wherein X⁷ is -SO₂-, -O- or -CONR⁴⁸R⁴⁹- (wherein R⁴⁸ and R⁴⁹, which may be the same or different, each represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R⁴⁷ represents C₁₋₅alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro and amino) with the provisos that when X⁷ is -SO₂-, X¹ is -O-, when X⁷ is -O-, X¹ is carbonyl, when X⁷ is -CONR⁴⁸R⁴⁹-, X¹ is -O- or NR¹⁸ (wherein R⁴⁸, R⁴⁹ and R¹⁸ are as defined hereinbefore);
- 10) C₂₋₆alkenylR³⁷ (wherein R³⁷ is as defined hereinbefore);
- 11) C₂₋₆alkynylR³⁷ (wherein R³⁷ is as defined hereinbefore);
- 12) C₂₋₆alkenylX⁸R³⁷ (wherein X⁸ represents -O-, -S-, -SO-, -SO₂-, -NR⁵⁰CO-, -CONR⁵¹-, -SO₂NR⁵²-, -NR⁵³SO₂- or -NR⁵⁴- (wherein R⁵⁰, R⁵¹, R⁵², R⁵³ and R⁵⁴ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R³⁷ is as defined hereinbefore);
- 13) C₂₋₆alkynylX⁹R³⁷ (wherein X⁹ represents -O-, -S-, -SO-, -SO₂-, -NR⁵⁵CO-, -CONR⁵⁶-, -SO₂NR⁵⁷-, -NR⁵⁸SO₂- or -NR⁵⁹- (wherein R⁵⁵, R⁵⁶, R⁵⁷, R⁵⁸ and R⁵⁹ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R³⁷ is as defined hereinbefore);
- 14) C₁₋₃alkylX¹⁰C₁₋₃alkylR³⁷ (wherein X¹⁰ represents -O-, -S-, -SO-, -SO₂-, -NR⁶⁰CO-, -CONR⁶¹-, -SO₂NR⁶²-, -NR⁶³SO₂- or -NR⁶⁴- (wherein R⁶⁰, R⁶¹, R⁶², R⁶³ and R⁶⁴ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R³⁷ is as defined hereinbefore);

15) R^{36} (wherein R^{36} is as defined hereinbefore); and

16) $C_{1-3}alkylX^{10}C_{1-3}alkylR^{36}$ (wherein X^{10} and R^{36} are as defined hereinbefore).


2. A compound according to claim 1 wherein R^9 is substituted by one or
 5 more groups selected from hydroxy; halo; nitro; cyano; carboxy; $C_{1-6}alkoxy$; $C_{1-6}alkyl$; $C_{2-6}alkenyl$; $C_{2-6}alkynyl$; $C_{2-6}alkenyloxy$; $C_{2-6}alkynyloxy$; $C_{3-6}cycloalkyl$; amino; mono- or di-
 $C_{1-6}alkyl$ amino; heterocyclyl optionally substituted with $C_{1-6}alkyl$ or oxo; $C(O)R^a$,
 $C(O)OR^a$, $S(O)_dR^a$; $NR^aC(O)R^b$; $C(O)NR^aS(O)_dR^b$, $C(O)NR^aR^b$; $NR^aC(O)NR^bR^c$;
 $NR^aS(O)_dR^b$ or $N(S(O)_dR^b)S(O)_dR^c$ where d is 0, 1 or 2 and R^a , R^b and R^c are
 10 independently selected from hydrogen, $C_{1-6}alkyl$, aryl, $C_{3-6}cycloalkyl$ or heterocyclyl, and
 wherein any alkyl, alkenyl or alkynyl group or moiety contained within the substituent one
 R^9 may themselves be optionally substituted with one or more groups selected from
 hydroxy; cyano; nitro; halo; carboxy; carboalkoxy of 2-7 carbon atoms, $C_{3-6}cycloalkyl$,
 heterocyclyl optionally substituted with $C_{1-6}alkyl$ or oxo; $C(O)R^d$, $C(O)OR^d$, NR^dR^e , $S(O)_e$
 15 R^d , $NR^dC(O)R^e$; $C(O)NR^dR^e$; $NR^dC(O)NR^eR^f$; $NR^dS(O)_eR^e$ where e is 0, 1 or 2 and R^d ,
 R^e and R^f are independently selected from hydrogen or $C_{1-6}alkyl$ optionally substituted
 with one or more groups selected from hydroxy; cyano; nitro; halo; carboxy; carboalkoxy
 of 2-7 carbon atoms, $C_{3-6}cycloalkyl$, heterocyclyl optionally substituted with $C_{1-6}alkyl$ or
 oxo; $C(O)R^g$, $C(O)OR^g$, NR^gR^h , $S(O)_eR^g$, $NR^gC(O)R^h$; $C(O)NR^gR^h$; $NR^gC(O)NR^hR^i$;
 20 $NR^gS(O)_eR^h$ where e is as defined above and R^g , R^h and R^i are independently selected
 from hydrogen or $C_{1-6}alkyl$: or two substituents on adjacent atoms may be joined to form
 the second ring of a bicyclic ring system wherein the said second ring is optionally
 substituted with one or more of the groups listed above for R^9 and optionally contains one
 or more heteroatoms.

25

3. A compound according to claim 1 where R^9 is phenyl substituted with an
 optionally substituted alkoxy group.

4. A compound according to claim 1 which is a compound of formula (IA)

178



(IA)

or a pharmaceutically acceptable salt thereof.

wherein:

5 n is 0-1;

X and Y are independently selected from $-NH-$, $-O-$, $-S-$, or $-NR^8-$ where R^8 is alkyl of 1-6 carbon atoms and X may additionally comprise a CH_2 group;

10 R^7 is a group $(CH_2)_m R^9$ where m is 0, or an integer of from 1-3 and R^9 is a substituted aryl or substituted cycloalkyl ring of up to 10 carbon atoms, wherein the substituents comprise at least one alkoxy group of 1-6 carbon atoms and optionally one or more further substituents, or R^9 is a heterocyclic ring containing 1 or 2 oxygen atoms and optionally one or more substituents;

15 R^6 is a divalent cycloalkyl of 3 to 7 carbon atoms, which may be optionally further substituted with one or more alkyl of 1 to 6 carbon atom groups; or is a divalent pyridinyl, pyrimidinyl, or phenyl ring; wherein the pyridinyl, pyrimidinyl, or phenyl ring may be optionally further substituted with one or more groups selected from halogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, azido, hydroxyalkyl of 1-6 carbon atoms, halomethyl, alkoxymethyl of 2-7 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkoxy of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, hydroxy, trifluoromethyl, cyano, nitro, carboxy, carboalkoxy of 2-7 carbon atoms, carboalkyl of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzoyl, benzyl, amino, alkylamino of 1-6 carbon atoms, dialkylamino of 2 to 12 carbon atoms, phenylamino, benzylamino, alkanoylamino of 1-6 carbon atoms, alkenoylamino of 3-8 carbon atoms, alkynoylamino of 3-8 carbon atoms, and benzoylamino;

25

R_1 , R_2 , R_3 and R_4 are each independently selected from hydrogen, hydroxy, halogeno, cyano, nitro, trifluoromethyl, C_{1-3} alkyl, $-NR^{11}R^{12}$ (wherein R^{11} and R^{12} , which may be the

same or different, each represents hydrogen or C_{1-3} alkyl), or a group $R^{13}-X^1-(CH_2)_x$ wherein x is 0 to 3, X^1 represents -O-, -CH₂-, -OCO-, carbonyl, -S-, -SO-, -SO₂-, -NR¹⁴CO-, -SO₂NR¹⁶-, -NR¹⁷SO₂- or -NR¹⁸- (wherein R^{14} , R^{15} , R^{16} , R^{17} and R^{18} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{13} is selected from one of the following sixteen groups:

- 1) C_{1-5} alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro and amino;
- 2) C_{1-5} alkyl X^2 COR¹⁹ (wherein X^2 represents -O- or -NR²⁰- (wherein R^{20} represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{19} represents -NR²¹R²²- or -OR²³- (wherein R^{21} , R^{22} and R^{23} which may be the same or different each represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl));
- 3) C_{1-5} alkyl X^3 R²⁴ (wherein X^3 represents -O-, -S-, -SO-, -SO₂-, -OCO-, -NR²⁵CO-, -CONR²⁶-, -SO₂NR²⁷-, -NR²⁸SO₂- or -NR²⁹- (wherein R^{25} , R^{26} , R^{27} , R^{28} and R^{29} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{24} represents hydrogen, C_{1-3} alkyl, cyclopentyl, cyclohexyl or a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which C_{1-3} alkyl group may bear one or two substituents selected from oxo, hydroxy, halogeno and C_{1-4} alkoxy and which cyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, C_{1-4} alkyl, C_{1-4} hydroxyalkyl and C_{1-4} alkoxy);
- 4) C_{1-5} alkyl X^4 C_{1-5} alkyl X^5 R³⁰ (wherein X^4 and X^5 which may be the same or different are each -O-, -S-, -SO-, -SO₂-, -NR³¹CO-, -CONR³²-, -SO₂NR³³-, -NR³⁴SO₂- or -NR³⁵- (wherein R^{31} , R^{32} , R^{33} , R^{34} and R^{35} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{30} represents hydrogen or C_{1-3} alkyl);
- 5) C_{1-5} alkylR³⁶ (wherein R^{36} is a 5 or 6 membered saturated heterocyclic group with one or two heteroatoms, selected independently from O, S and N, which heterocyclic group may bear one or two substituents selected from oxo, hydroxy, halogeno, C_{1-4} alkyl, C_{1-4} hydroxyalkyl and C_{1-4} alkoxy);
- 6) $(CH_2)_qX^6R^{37}$ (wherein q is an integer from 0 to 5, X^6 represents a direct bond, -O-, -S-, -SO-, -SO₂-, -NR³⁸CO-, -CONR³⁹-, -SO₂NR⁴⁰-, -NR⁴¹SO₂- or -NR⁴²- (wherein R^{38} , R^{39} , R^{40} , R^{41} and R^{42} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl) and R^{37} is a phenyl group, a pyridone group or a 5 or 6 membered aromatic heterocyclic group with 1 to 3 heteroatoms selected from O, N and S, which phenyl, pyridone or

aromatic heterocyclic group may carry up to 5 substituents selected from hydroxy, halogeno, amino, C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄hydroxyalkyl, C₁₋₄hydroxyalkoxy, C₁₋₄aminoalkyl, C₁₋₄alkylamino, carboxy, cyano, -CONR⁴³R⁴⁴ and -NR⁴⁵COR⁴⁶ (wherein R⁴³, R⁴⁴, R⁴⁵ and R⁴⁶, which may be the same or different, each represents hydrogen, C₁₋₄alkyl or C₁₋₃alkoxyC₂₋₃alkyl));

7) C₂₋₆alkenylR³⁶ (wherein R³⁶ is as defined hereinbefore);

8) C₂₋₆alkynylR³⁶ (wherein R³⁶ is as defined hereinbefore);

9) X⁷R⁴⁷ (wherein X⁷ is -SO₂-, -O- or -CONR⁴⁸R⁴⁹- (wherein R⁴⁸ and R⁴⁹, which may be the same or different, each represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R⁴⁷ represents C₁₋₅alkyl which may be unsubstituted or which may be substituted with one or more groups selected from hydroxy, fluoro and amino) with the provisos that when X⁷ is -SO₂-, X¹ is -O-, when X⁷ is -O-, X¹ is carbonyl, when X⁷ is -CONR⁴⁸R⁴⁹-, X¹ is -O- or NR¹⁸ (wherein R⁴⁸, R⁴⁹ and R¹⁸ are as defined hereinbefore);

10) C₂₋₆alkenylR³⁷ (wherein R³⁷ is as defined hereinbefore);

11) C₂₋₆alkynylR³⁷ (wherein R³⁷ is as defined hereinbefore);

12) C₂₋₆alkenylX⁸R³⁷ (wherein X⁸ represents -O-, -S-, -SO-, -SO₂-, -NR⁵⁰CO-, -CONR⁵¹-, -SO₂NR⁵²-, -NR⁵³SO₂- or -NR⁵⁴- (wherein R⁵⁰, R⁵¹, R⁵², R⁵³ and R⁵⁴ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R³⁷ is as defined hereinbefore);


13) C₂₋₆alkynylX⁹R³⁷ (wherein X⁹ represents -O-, -S-, -SO-, -SO₂-, -NR⁵⁵CO-, -CONR⁵⁶-, -SO₂NR⁵⁷-, -NR⁵⁸SO₂- or -NR⁵⁹- (wherein R⁵⁵, R⁵⁶, R⁵⁷, R⁵⁸ and R⁵⁹ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R³⁷ is as defined hereinbefore);

14) C₁₋₃alkylX¹⁰C₁₋₃alkylR³⁷ (wherein X¹⁰ represents -O-, -S-, -SO-, -SO₂-, -NR⁶⁰CO-, -CONR⁶¹-, -SO₂NR⁶²-, -NR⁶³SO₂- or -NR⁶⁴- (wherein R⁶⁰, R⁶¹, R⁶², R⁶³ and R⁶⁴ each independently represents hydrogen, C₁₋₃alkyl or C₁₋₃alkoxyC₂₋₃alkyl) and R³⁷ is as defined hereinbefore);

15) R³⁶ (wherein R³⁶ is as defined hereinbefore); and

16) C₁₋₃alkylX¹⁰C₁₋₃alkylR³⁶ (wherein X¹⁰ and R³⁶ are as defined hereinbefore).

5. A compound according to claim 1 of formula (II)



5


where R^1 , R^2 , R^3 and R^4 are as defined in claim 1, R^{66} is an optionally substituted C_{1-6} alkyl and R^{67} is selected from hydrogen, halogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, azido, hydroxyalkyl of 1-6 carbon atoms, halomethyl, alkoxymethyl of 2-7 carbon atoms, alkanoyloxymethyl of 2-7 carbon atoms, alkoxy of 1-6 carbon atoms, alkylthio of 1-6 carbon atoms, hydroxy, trifluoromethyl, cyano, nitro, carboxy, carboalkoxy of 2-7 carbon atoms, carboalkyl of 2-7 carbon atoms, phenoxy, phenyl, thiophenoxy, benzoyl, benzyl, amino, alkylamino of 1-6 carbon atoms, dialkylamino of 2 to 12 carbon atoms, phenylamino, benzylamino, alkanoylamino of 1-6 carbon atoms, alkenoylamino of 3-8 carbon atoms, alkynoylamino of 3-8 carbon atoms, and benzoylamino.

10

15

182

6. A compound of formula (IB)




(IB)

5 where Y, n, R⁶, X and R⁷ are as defined in claim 1 and at least one of R^{1''}, R^{2''}, R^{3''} or R^{4''} is a group R^{13'}-X¹-(CH₂)_x wherein X¹ and x are as defined in claim 1 and R^{13'} is alkyl substituted by chloro or bromo; and the remainder are groups R¹, R², R³ and R⁴ respectively.

10 7. A pharmaceutical composition comprising a compound of formula (I) as defined in claim 1 in combination with a pharmaceutically acceptable carrier or excipient.


8. A method of preparing a compound of formula (I) as defined in claim 1 which method comprises either (a) reacting a compound of formula (III)



(III)

where R^{1'}, R^{2'}, R^{3'}, R^{4'} represent R¹, R², R³ and R⁴ respectively as defined in relation to formula (I) or a precursor thereof, and Z' is a leaving group, with a compound of formula

(IV)




183

(IV)

where R^6 , Y, X, and n are as defined in relation to formula (I), and $R^{7'}$ is a group R^7 or a precursor thereof; or

(b) reacting a compound of formula (V)



(V)

where $R^{1'}$, $R^{2'}$, $R^{3'}$, $R^{4'}$ are as defined in relation to formula (III) R^6 , X, Y and n are as defined in relation to formula (I), with a compound of formula (VI)

$$R^{7'}-Z'' \quad (VI)$$

where R^7 is as defined in relation to formula (IV) and Z'' is a leaving group;

and thereafter if necessary or desired converting precursor groups $R^{1'}$, $R^{2'}$, $R^{3'}$, $R^{4'}$ and $R^{7'}$ to groups of formula R^1 , R^2 , R^3 , R^4 and R^7 respectively, or converting a group R^1 , R^2 , R^3 , R^4 and R^7 to a different such group.

9. A compound for use in therapy comprising a compound of formula (I) as defined in claim 1.

10. The use of a compound of formula (I) as defined in claim 1 in the preparation of a medicament for use in the inhibition of MEK enzymes.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 00/01697

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D215/54 A61K31/47 A61P43/00 C07D405/12 C07D401/12
C07D417/12 C07D413/12 C07D409/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 43960 A (AMERICAN CYANAMID COMPANY) 8 October 1998 (1998-10-08) cited in the application page 2, line 23 - line 26; claim 1 -----	1,7,10
A	WO 99 01426 A (WARNER-LAMBERT COMPANY) 14 January 1999 (1999-01-14) page 3, line 10 - line 15; claim 1 -----	1,7,10
P,X	WO 00 18761 A (AMERICAN CYANAMID COMPANY) 6 April 2000 (2000-04-06) page 3, line 2 - line 5; claim 1 page 139 -page 142 -----	1,7,10

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

7 September 2000

Date of mailing of the international search report

19/09/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 00/01697

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9843960	A	08-10-1998	AU	6877798 A	22-10-1998
			EP	0973746 A	26-01-2000
			NO	994798 A	24-11-1999
			PL	335999 A	05-06-2000
WO 9901426	A	14-01-1999	AU	8262798 A	25-01-1999
			EP	0993439 A	19-04-2000
			HR	980368 A	30-04-1999
			NO	996491 A	29-12-1999
			ZA	9805728 A	27-01-1999
WO 0018761	A	06-04-2000	AU	6159399 A	17-04-2000